


Geologica Macedonica

Journal of the Geological Institute at the Faculty of Natural and
Technical Sciences, University "Goce Delčev"-Štip, R. Macedonia

Microcline-Belutče



<i>Geologica Macedonica</i>	Vol.	29	No	1	pp.	1-100	Štip	2015
<i>Geologica Macedonica</i>	Год.		Број		стр.		Штип	

UDC 55
CODEN – GEOME 2

In print: ISSN 0352–1206
On line: ISSN 1857–8586

GEOLOGICA MACEDONICA

<i>Geologica Macedonica</i>	Vol.	29	No	1	pp.	1–100	Štip	2015
<i>Geologica Macedonica</i>	Год.		Број		стр.		Штип	

<i>Geologica Macedonica</i>	Vol.	29	No	1	pp.	1–100	Štip	2015
<i>Geologica Macedonica</i>	Год.		Број		стр.		Штип	

GEOLOGICA MACEDONICA

Published by: – Издава:

The "Goce Delčev" University, Faculty of Natural and Technical Sciences, Štip, Republic of Macedonia
Универзитет „Гоце Делчев“, Факултет за природни и технички науки, Штип, Република Македонија

EDITORIAL BOARD

Todor Serafimovski (R. Macedonia, *Editor in Chief*), **Prof. Blažo Boev** (R. Macedonia, *Editor*),
David Alderton (UK), Tadej Dolenc (R. Slovenia), Ivan Zagorchev (R. Bulgaria), Wolfgang Todt (Germany),
acad. Nikolay S. Bortnikov (Russia), Clark Burchfiel (USA), Thierry Augé (France), Todor Delipetrov (R. Macedonia),
Vlado Bermanec (Croatia), Milorad Jovanovski (R. Macedonia), Spomenko Mihajlović (Serbia),
Dragan Milovanović (Serbia), Dejan Prelević (Germany), Albrecht von Quadt (Switzerland)

УРЕДУВАЧКИ ОДБОР

Тодор Серафимовски (Р. Македонија, *главен уредник*), **Блажо Боев** (Р. Македонија, *уредник*),
Дејвид Олдертон (В. Британија), Тадеј Доленец (Р. Словенија), Иван Загорчев (Р. Бугарија),
Волфганг Тод (Германија), акад. Николај С. Бортников (Русија), Кларк Барчфил (САД), Тиери Оже (Франција),
Тодор Делипетров (Р. Македонија), Владо Берманец (Хрватска), Милорад Јовановски (Р. Македонија),
Споменко Михајловиќ (Србија), Драган Миловановиќ (Србија), Дејан Прелевиќ (Германија),
Албрехт фон Квад (Швајцарија)

Language editor	Лектура
Marijana Kroteva	Маријана Кротева
(English)	(англиски)
Georgi Georgievski, PhD.	д-р Георги Георгиевски
(Macedonian)	(македонски)

Technical editor	Технички уредник
Blagoja Bogatinoski	Благоја Богатиноски
Proof-reader	Коректор
Alena Georgievska	Алена Георгиевска

Address	Адреса
GEOLOGICA MACEDONICA	GEOLOGICA MACEDONICA
EDITORIAL BOARD	РЕДАКЦИЈА
Faculty of Natural and Technical Sciences	Факултет за природни и технички науки
P. O. Box 96	пошт. факс 96
МК-2000 Štip, Republic of Macedonia	МК-2000 Штип, Република Македонија
Tel. ++ 389 032 550 575	Тел. 032 550 575
E-mail: todor.serafimovski@ugd.edu.mk	

400 copies	Тираж: 400
Published twice yearly	Излегува двапати годишно

Printed by:	Печати:
2 nd August – Štip	2 nd Август – Штип

Price: 500 den	Цена: 500 ден.
The edition is published in June 2015	Бројот е отпечатен во јуни 2015

<i>Geologica Macedonica</i>	Vol.	29	No	1	pp.	1–100	Štip	2015
<i>Geologica Macedonica</i>	Год.		Број		стр.		ШТИП	

TABLE OF CONTENTS

285. Tena Šijakova-Ivanova, Vesna Ambarkova Geochemical assessment of some natural waters from eastern and south-eastern Macedonia	5–14
286. Vojo Mirčovski, Blažo Boev, Zlatko Efremoski, Ajka Šorša, Đorđi Dimov Hydrochemical data for the ground waters in the Bitola's part of the Pelagonia valley, Republic of Macedonia	15–24
287. Dafinka Dimitrova, Ivan Boev, Vesna Zajkova Panova Heavy and toxic metals in water and sediments of the river of Vardar in Skopje	25–46
288. Orce Spasovski, Daniel Spasovski Possibilities for using travertine from the Nabigori deposit village of Velmej (western Macedonia) as architectural stone	47–52
289. Biljana Balabanova, Trajče Stafilov, Robert Šajn, Claudiu Tănăselia Lead distribution in soil due to lithogenic and anthropogenic factors in the Bregalnica river basin.....	53–61
290. Igor Ivanovski, Dalibor Serafimovski, Goran Tasev, Todor Serafimovski 3D modelling of the Plavica Au-Cu polymetallic deposit, Republic of Macedonia	63–74
291. Igor Peševski, Milorad Jovanovski, Jovan Br. Papić, Biljana Abolmasov Model for GIS landslide database establishment and operation in Republic of Macedonia	75–86
292. Arianit A. Reka, Blagoj Pavlovski, Todor Anovski, Slobodan Bogoevski, Boško Boškovski Phase transformations of amorphous SiO ₂ in diatomite at temperature range of 1000–1200°C	87–92
293. Vladimir Manevski, Todor Delipetrov, Marjan Delipetrov Geoelectrical measurements in environment with presence of underground water	93–98
Instructions to authors	99–100

<i>Geologica Macedonica</i>	Vol.	29	No	1	pp.	1–100	Štip	2015
<i>Geologica Macedonica</i>	Год.		Број		стр.		ШТИП	

СОДРЖИНА

285. Тена Шијакова-Иванова, Весна Амбаркова Геохемиска процена на некои природни води од источна и југоисточна Македонија	5–14
286. Војо Мирчовски, Блажо Боев, Златко Ефремоски, Ајка Šorša, Ѓорги Димов Хидрохемиски податоци за подземните води на битолскиот дел од Пелагониската Котлина, Република Македонија	15–24
287. Дафинка Димитрова, Иван Боев, Весна Зајкова Панова Тешки и токсични метали во водата и седиментите на реката Вардар во Скопје	25–46
288. Орце Спасовски, Даниел Спасовски Можности за користење на травертинот од наоѓалиштето Набигори, с. Велмеј (западна Македонија) како архитектонски камен	47–52
289. Biljana Balabanova, Trajče Stafilov, Robert Šajn, Claudiu Tănăselia Дистрибуција на олово во почвите како резултат на литогените и антропогените фактори во регионот на сливот на реката Брегалница.....	53–61
290. Игор Ивановски, Далибор Серафимовски, Горан Тасев, Тодор Серафимовски 3D моделирање на полиметаличното Au-Cu наоѓалиште Пластица, Република Македонија.....	63–74
291. Игор Пешевски, Милорад Јовановски, Јован Бр. Папиќ, Биљана Аболмасов Индекс за воспоставување и управување на банка на податоци за свлечишта во Република Македонија.....	75–86
292. Аринит А. Река, Благој Павловски, Тодор Ановски, Слободан Богоевски, Бошко Бошковски Фазни трансформации на аморфен SiO ₂ во дијатомит во температурен интервал 1000–1200 °C.....	87–92
293. Владимир Маневски, Тодор Делипетров, Марјан Делипетров Геоелектрични мерења во средина со присуство на подземна вода	93–98
Упатство за авторите	99–100

3D MODELLING OF THE PLAVICA Au-Cu POLYMETALLIC DEPOSIT, REPUBLIC OF MACEDONIA

Igor Ivanovski¹, Dalibor Serafimovski², Goran Tasev², Todor Serafimovski²

¹*Genesis Resources International DOOEL Skopje, Republic of Macedonia*

²*Faculty of Natural and Technical Sciences, Institute of Geology, University “Goce Delčev” in Štip,
Blvd. Goce Delčev no.89, Štip, Republic of Macedonia
icestip@yahoo.com*

A b s t r a c t: The latest exploration and study of the high sulfidation epithermal deposit of Plavica displayed significant progress in understanding the geology and definition of the particular ore body. The results from boreholes made by Genesis Resources International DOOEL Skopje, Republic of Macedonia, in 2011, 2012 and 2013 were: 51 m interval with 3.9 g/t Au in oxidation zone, 65 m interval with 3.1 g/t Au in oxidation zone, 51 m interval with 3.7 g/t Au, 2.8% Cu and 16 g/t Ag in sulfide zone. For this 3D model all 195 boreholes made by the Genesis Resources International were used. The boreholes were made in the period of 2011–2013, with total length of 47295.8 m of which 74 boreholes were performed by diamond core method and total length of 23908.3 m while the other 121 boreholes were performed by rotary circulation (RC) with total length of 23387.5 m. Using professional ArcGIS software were prepared 3D models of the ore body for gold and copper, at respective resolution of 30 m and 10 m, for each element, which have defined mineralization extent from 1305 m to 600 m of the hypsometric level. These models should improve our understanding of the ore mineralization and complement the geological data of the Plavica ore deposit.

Key words: 3D model; Plavica; gold; copper; epithermal

INTRODUCTION

Polymetallic gold, copper and silver ore deposit Plavica is part of Kratovo–Zletovo volcanic area in the NE part of the Republic of Macedonia. It occupies the central part of the Plavica mountain (1290 m) and is located between cities of Probištip (6 km distance) and Kratovo (4 km distance). The Plavica deposit is well known to the local and regional community. Its exploration and exploitation history (in the central parts known as Zlatica) reaches back to Roman times when were undertaken large-scale mining and processing of copper from pyrite-enargite veins (sporadically limited exploitation of silver and lead). Limited mining took place here during and after the so called ‘Turkish’ era (XV–XVII centuries) when gold was extracted for production of coins. The British company Selection Mines Ltd carried out some exploration in the area and further exploitation of enargite veins between 1934 and 1936 (with more than 2800 m of underground exploration workings). Later on an intense programme of exploration and drilling was undertaken by the Zletovo

mine and the Geological Survey in the decades following the Second World War (1945–1952; 1961–1962 and 1963–1974). Based on those explorations an extensive programme prepared was by the Geological Survey and RIK “Sileks”, realized in the period 1976–1986 (with respectable 30 325 m of exploration drilling). From the late 1980 s, there have been several, further, exploration programmes for Cu and Au by foreign mining companies (e.g. Cominco, Cyprus Amax, Minorco, Goldfields, Rio Tinto, European Minerals) with the emphasis more on Au associated with the peripheral silica bodies. The latest exploration in the Plavica concession area was performed during the period 2011–2014 by the Australian company Genesis Resources International Ltd. All those explorations contributed to an intensive professional and scientific database where we would like to point out the workings of Pendžerkovski et al. (1960), Mijalković and Pešić (1966), Janković (1967), Marković (1971), Pantić et al. (1972), Ristić and Klajn (1973), Stojanov (1974, 1980), Bili-

bajkić and Bilibajkić (1978), Petković and Romić (1977), Ivanov and Denkovski (1978, 1980), Rakić (1978, 1982), Janković et al. (1980), Mudrinić (1982, 1986), Drovenik et al. (1983), Bogoevski and Rakić (1985), Stojanov and Serafimovski (1990), Serafimovski (1990, 1993), Stojanov et al. (1995), Tomson et al. (1998), Serafimovski and Rakić (1998, 1999), Serafimovski and Tasev (2003), Stefanova (2005), Alderton and Serafimovski (2007), Stefanova et al. (2007), Volkov et al. (2010), Stefanova et al. (2013) and Zlatkov et al. (2014).

During last two decades the 3D geologic modelling has evolved from the application of

computer aided design (CAD) software into a matured Earth science discipline that covers geomathematics, geophysics and geology. In that direction 3D and 4D modelling are the new exploration tools that can help the mineral explorers to visualize, interpolate and interpret geological data, are a critical time and money-saving methods. This particular model was built in Target 3D for ArcGIS system, and using this as a basis for modelling of the certain Au-Cu ore body. A better understanding of the geometry and intensity in 3D may lead to new discoveries within the deposit area.

METHODOLOGY

The database with drill core collar, survey, assay and all the data have been provided by Genesis Resources International DOOEL Skopje. The database contains information from 195 boreholes with individual length of up to 594.1 m. Although exploration concession is for 17.41 km², the latest explorations displayed within this paper covered an area of 0.63 km², which have been estimated by geological experts as the most promising one. Drillholes have been spaced 50 to 100 m between each other and sampled at 1 m intervals, in that direction the total database consists of 43965 sampling results (multi-elemental). All of the drill holes were located using GPS, and these coordinates and elevations were recorded. The database provides information about the location of drillholes, their depth and orientation, lithology, mineralogy, and geochemistry (Atomic Adsorption Spectrometry – [AAS] and Inductively Coupled Plasma Atomic Emission Spectrometry – [ICP-AES], analyses of 36 chemical elements). Cut of

grade for gold was calculated at 0.2 g/t Au while for copper it used to be at 0.2% Cu.

For the construction of the Plavica deposit 3D model was used Target 3D, which was developed for the Target for ArcGIS system. This particular system enables to display drillhole, surface and other data types in an interactive three-dimensional environment. The *Target 3D Viewer* enables three-dimensional viewing of our data. Drillholes are displayed in their "true" three-dimensional location and can have up to two different data types plotted along their trace. Grids created in "sectional" views (e.g. from Target or Interactiv IP applications) can be displayed directly into the 3D view, in their correct orientation, using Geosoft's "on-the-fly" technology. Other grids and images (including bitmaps and jpegs) can be opened and easily located in any specified orthogonal plane. 3D "Voxel grids" can also be displayed in a 3D drillhole map and modified using the Target 3D Tool. 3D Voxels can be created using the Voxels/Grid Voxel menu item, on the Target 3D Toolbar.

GEOLOGICAL FEATURES AND MINERALOGY OF THE DEPOSIT

The Plavica deposit is a part of well known Kratovo–Zletovo ore area that in general is characterized with a complex composition of mainly Tertiary volcanic-sedimentary complex imposed on the foundation of the Serbian–Macedonian massif, which represents constituting part of the western Tethyan belt. K/Ar radiometric data (Karamata et al., 1992) yield an age of 32–16 Ma (late Oligocene to middle Miocene). In the geological setting of the Plavica deposit participate mainly vol-

canic and volcano-sedimentary rocks, which exact determination is complex due to intensive presence of hydrothermal alterations (Figure 1).

However, with numerous lithostratigraphic and petrographic studies it was confirmed that in the geological setting of this deposit participate ignimbrite, stratified volcanic tuff and breccia, dacito-andesite and their pyroclasts as well as quartzlatites that the most often occur as break-

troughs. Definition of these rocks is hard because of intensive hydrothermal alterations as are: kaolinization, silification, alunization, seritization and other. The central part of Plavica is made of different kind of volcanic rocks, hydrothermally altered,

pyritized, and mineralized with ore mineralization (Cu, Au, Ag, Pb, Zn). According the research made on Plavica mainly based on boreholes data, the depth of the ore mineralization and intensive hydrothermal alteration exceeds 1000 m.




Fig. 1. Simplified geological map of the Plavica deposit area (Ivanovski, 2014)

The silica ledges seen on Plavica represent the level of paleo-regional ground water table. Due to their silica content both areas have resisted erosion and through topographic inversion have formed prominent landforms (Figure 2). The extend of erosional landform processes is illustrated in the surrounding region by the widespread occurrence of large scale block slides, slumps and smaller scale mud slides. Landform failure appears to have occurred along steeply dipping contacts between individual sub-aerial volcanic flow-air fall

units and regions of subaqueous reworked volcanic detritus. The contacts between these units quite commonly show quartz-sericite clay alteration after pyrite thus contributing to the loss in rock strength and consequent landform failure. The volcanic units of Plavica are protected from erosion by the widely distributed silica ledges (Figure 2) and these units dip towards the SE (Cuthberston, 1998). The Plavica area is underlain by a thick section (nearly 500 m) of intercalated volcanic flows, tuffs, debris flows and subvolcanic intrusions.




Fig. 2. Plavica conceptual cross-section (Stevanović, 2014)

Flows and massive tuffs ranging in composition from andesite to latite, comprise the basal portion of the volcanic sequence and are overlain by a series of debris flows with intercalated water-lain tuffs, tuffs and volcano-clastic sediments. In addition, the extensively developed hydrothermal alteration, faulting, and large amounts of alluvial cover complicate attempts to make detailed stratigraphic determinations (Percival, 1992). The geology of Plavica consists of a package of a high level intermediate volcanics of mixed sub-aerial and sub-marine facies. The package varies from coarse grained crystal and lithic tuffs to finely laminated sub-aqueous epiclasts that dip approximately 30° to the SSW. This volcanic package is cut by apparently sub-vertical vuggy silica bodies of narrow extend (100–150 m) that, due to erosion and topog-

raphic inversion have become topographic highs (Fish, 1998; Fish, et al., 2000).

The Eocene–Miocene igneous rocks associated with the area are mostly of an intermediate, calc-alkaline/high K-composition (especially granodiorite, quartz, monzonite, quartz diorite, andesite and dacite). The present level of erosion is such that the igneous rocks currently exposed are mostly high level volcanics and pyroclastics (Alderton and Serafimovski, 2007).

Tertiary rocks are the dominant lithology at the Plavica area, review of these numerous and different types of rocks will be present according their geochronological order defined on basis of the found fauna superpositional relations and absolute age of certain types of volcanic rocks (Serafimovski et al., 2014)

Ignimbrite of dacite-andesite composition are present on NE slopes of the Plavica (Trnjak, Golak, Ramna Niva) and in one wide band envelope southern slopes of the Plavica. They are represented by grey-green to reddish color and composition variable at short distances (andesite-dacite-quartzlatite). Due to their structural setting and inclusion of some older rocks some authors named them as ignimbrites. These rocks are known for the presence of numerous lead-zinc ore veins. This quite well fits Plavica deposit into the explanation that although the most high-sulfidation deposits are generated in calc-alkaline andesitic-dacitic arcs characterized by near neutral stress states or mild extension, few major deposits may also occur in compressive arcs characterized by the suppression of volcanic activity (Sillitoe and Hedenquist, 2003), as it is shown in the text that follows, too.

Volcano-sedimentary rocks (stratified tuff) are well spread on the northern margins of the Plavica in an elongated belt starting from Šlegovo village and ending to the east, near the Siroka Padina locality. Their presence points out that some of the volcanic stages occurred in submarine or sublacustrine conditions. These rocks are sometimes hydrothermally altered, mainly, along the cracks and fissures.

Propylitized dacite-andesite rocks are the rock counterparts of former composition strongly hydrothermally altered and spread over northern slopes of the Plavica (Rajkovac, Golak, Trnjak) and southern part of the Maričanski Rid. Hydrothermally altered andesite rocks were determined in western and southwestern margins of the Plavica (saved fenocrystals and some colored compounds that easily can be classified as biotite-hornblende andesite).

Quartzlatites were determined at several locations (Karac, Dogandziski Kamen, Plavički Potok). In general they occur as neck, stock and dyke forms that break-through the adjacent dacite-andesite or their tuff counterparts. These rocks are built of cryptocrystalline quartz, K-feldspar with phenocrysts of sanidine, biotite and quartz and characterized by porphyry structure and massive texture.

Dykes are common at several places, especially on the northern margin of the Plavica area, where they break-through the dacite-andesite and pyroclasts.

Silex-secondary quartzite in the central part of the Plavica have a significant distribution. They build the highest parts of the Plavica and stick out of the ground in the form of chops and occur in

two elongated zones: northern and southern. Towards the east these silex ore bodies contain significant amounts of alunite that gradually turns into alunite quartzite. It is interesting to note that the silex-secondary quartzite except traces of sulphides contained in itself and some quantities of gold (0.1–3.0 g/t Au). Their transition towards the surrounding rocks are usually not sharp and they gradually transform into the surrounding rocks in form of more or less pronounced silification.

At the highest parts of the Plavica at several locations there are increase quantities of diluvial sediments (Beglički Livadi and Dolno Kadiski) in streams of northern slopes and remains of the ancient river terraces.

In regards to the ore mineralization within the Plavica deposit, we would like to stress out that it has been deposited in empty spaces in fault-crack systems, veins, veinlets and metasomatically has been impregnated in adjacent hydrothermally altered volcanic rocks. Mineralization of impregnated type is the most common near the cracking systems. The mineralization occurs in four distinct settings: (i) stockwork and disseminated Cu–Au (+Mo, Ag) mineralization occurs in the central and deepest part of the system; (ii) veins of quartz, pyrite, sphalerite and enargite (\pm gold) occur at intermediate levels and appear to be superimposed on the stockwork; (iii) silica bodies occur peripheral to the central zone (ore bodies consist of quartz plus some opal and are sub-vertical in orientation); (iv) small veins with enrichments in Pb and Zn (up to 1% combined Pb–Zn), similar to the larger Pb–Zn veins worked in the nearby Zletovo mine, but much smaller. We would like to point out that the detailed exploration performed by the Genesis Resources International DOOEL Skopje, which study was concentrated mostly in the northeastern part of the Plavica deposit (mineralized area known as Plavica–Zlatica) and hill top where dominate secondary quartzites. Those extensive exploration activities defined one new Cu–Au mineralization type where dominate epithermal gold of high sulfidation with representative elongated lens-like ore bodies, which follows the structural controlling forms (Figure 1 and Figure 2).

The mineralogy of the Plavica deposit is complex and several paragenetic stages have been suggested (Ivanov and Denkovski, 1978). The latest data about the mineralogical composition of the Plavica deposit can be found in detailed electron microprobe study of Serafimovski (1993), where the mineral presence and their association has been

brought to an higher level of knowledge. Also, later findings of Serafimovski and Rakić (1998), Serafimovski and Rakić (1999), Alderton and Serafimovski (2007), Stefanova et al. (2013), Serafimovski and Tasev (2013) and Zlatkov et al., (2014), contributed to a better understanding of the mineral composition of the Plavica mineral deposit. With up to date studies has been confirmed that within the Plavica–Zlatica deposit can be found the following ore minerals: pyrite, pyrrotite, chalcopyrite, magnetite, rutile, scheelite, hematite, molybdenite, galena, sphalerite, bornite, enargite, native gold, melnikovite, tetrahedrite, tennantite, chalcocite, digenite, covellite, proustite, native copper, malachite, azurite, and some other. The latest study have confirmed presence of seligmannite, luzonite, famatinite, petzite, bogdanovite, bezsmertnovite, sylvanite, pearceite, bilibinskite etc. (Serafimovski and Tasev, 2013). Of the non-metallic minerals the most common were quartz, calcite, siderite, oligonite, chalcedony etc. The common presence of copper minerals, especially

the high-sulfidation state sulfosalts enargite-luzonite (White et al., 1995), as well as relatively high-sulfidation state minerals tennantite-tetrahedrite (Barton and Skinner, 1979), was the factors leading us to classify deposit as high-sulfidation type. Also, only scarce sphalerite and arsenopyrite confirmed our conclusion about the type of the deposit. High-sulfidation mineralization in the Plavica deposit is characterized by the presence of high-sulfidation – state sulfide assemblages dominated by pyrite, sulfosalts (enargite ± luzonite), digenite, bornite and lesser covellite, chalcocite, chalcopyrite, tennantite-tetrahedrite and molybdenite. Traces of colusite (Cu-Sn-V-As-Fe sulfide), sphalerite and galena have also been reported (Serafimovski et al., 2014) quite similar to genetically same deposits worldwide (Madera, 2000). The most common copper sulfides display a very distinctive replacement sequence from bornite → digenite → chalcocite → exsolution covellite (Zlatkov et al., 2014).

3D MODELLING, RESULTS AND DISCUSSION

A comprehensive and in many ways detailed study conducted at the Plavica, beside geochemistry and geophysics, included respectable array of 195 exploration boreholes totaling of 47295.8 m. One part of them, 74 boreholes, were with diamond core and total length of 23908.3 m with average depth of 400 m. The rest, 121 reverse circulation (RC) boreholes, were with total length of 23387.5 m and average depth of 200 m. All boreholes are with different azimuths and dips, but most of them are with azimuth of 360° and dip of –60°. We used all the boreholes, including the negative one so we can get more representative situation in the 3D modelling. The study results have shown interesting gold and copper concentrations in certain mineralized zones, which represents the basis for construction of ours 3D model with use of professional computer software package Target 3D for ArcGIS.

Using the Target 3D Tools enables us to interactively control the transparency of individual items, enabling data to be displayed with a cumulative (light-table) effect as it was already mentioned elsewhere (Ligovski et al., 2014). Data such as MapInfo tables and 2D DXF files can be imported directly into the 3D environment and drawn on any surface displayed in the current 3D view.

The Target system automatically detects the type of data that is being imported, including: drillhole (collar) locations with their unique Hole ID, Easting (X), Northing (Y), Elevation (Relative Level) and Total Depth (EOH) where the X, Y and Depth must all be in the same units. The Azimuth and Dip are also included in the collar file. Geochemical assay data are typically acquired by obtaining core or rotary drill samples over specific depth ranges (from-to ranges) and sending samples to an assay laboratory. Numerical results are typically returned from the laboratory in electronic format and can be imported quickly into the system. Before import, we must make sure that our data files contain the following information – HOLE ID, FROM, TO and a series of ASSAY results. On the DH-Data menu, click Import and then select Text File. Drill Hole menu>Ascii Import Wizard > Browse > KZM_ASSAYS (see Ligovski et al., 2014).

To start the work with Target 3D we selected an area that we would like to concentrate on from the Hole Selection Tool. This particular step enables us to display only the area of the project that is of interest. Target Drillhole > Hole Selection Tool dialog appears > Select using polygon tool to define a polygon around the holes required. However we had to create a Target 3D Voxel Grid,

which selects the holes we would like included in our Target 3D Voxel grid, using one of the Hole Selection Tool options, too.

From the Target Drillhole toolbar, we have to select Voxels > 3D Gridding from Current Geosoft Database > and 3D Gridding dialog is displayed. To Create a Target 3D Map On the Target Drillhole toolbar > Generate 3D Drillhole Plot b > 3D Map Parameters dialog, Page Layout tab, on the 3D Map Parameters dialog, is displayed by default. The following page layout parameters can be controlled from this tab. Background Colour of the 3D view Axis Colour and Font used for annotating the 3D view. Plot Legend (right side of map), including company Logo (image file) and specifying the map Titles. Select the Hole Traces tab >. This tab dialog includes the following drillhole trace parameters: > Colour of the Hole Trace Hole Labels including location, annotations, text size, colour and font .Depth Ticks (annotations etc.) along the hole trace Select the Data tab. The Data and Plot types are selected from drop-down lists, just as you would select data for your Plans or Sections to be plotted. The three available gridding methods are Minimum curvature, Kriging and a TIN based technique. To define the Gridded data parameters> Define button. Plot topography box to enable the topography parameters using the Browse buttons, locate the Topography grid M01GGRD file and the Overlay grid on topography file. 3D block models were used to produce vertical and horizontal slices at required depth levels. "Chair clipping" view was created to visualize 3D distribution within the investigated area. Threshold attribute values have been used to create the isosurfaces or threshold block models.

An element grade distribution block model was generated via ordinary kriging method. Ordinary kriging was developed by Matheron (1971) in the early sixties and plays a special role because it is compatible with a stationary model, only involves a variogram and it is the form of kriging used most in resource exploration (Chile's and Delfiner, 1999). Ordinary kriging estimates based on moving average of the variable of the interest satisfying different dispersion forms of data (Goovaerts, 1997). Although mineralization estimation of domains was modeled based only on metal grade, with definition of cut-off grade and top cut grade values for the Plavica deposit, the ore body was 3D modelled, with respect to intersecting fault zones, the overall geological setting and the compositional distribution of metal grades (Figures

3 and 4). Setting up the complete 3D geological model of the Plavica ore deposit all available data and sources of information were compared and integrated as it was case with 3D deposits elsewhere (Arvanitidis et al., 2013).


When plotted in 3D, the majority of the geochemical groupings had strong association with the lithologies of the deposit sequence (Figure 3a, b). The biggest advantages of constructing 3D volumes instead of 2D cross sections was the possibility to define ore grade volume gains and losses of the entire mineralized systems (Chmielowski et al., 2013; Christidis et al., 2014). In this direction we have proved that even with relatively slight changes in resolution (going from blocks formed at 30 m to blocks formed at 10 m), which have shown significant volume gains or losses of the entire mineralized, both for gold and copper (Figure 3a, b and Figure 4a, b), similarly to stated above.

Also, the contours of the ore bodies, both for gold and copper and both resolutions, have shown strong correlation to the fault systems in the area (being of N-S, NW-SE and sporadically SW-NE direction; see Figure 1) and respectively reflecting their spatial position.


Following sound geological reasoning during the creation of the model, a considerable amount of knowledge regarding the mineralized area and the ore body was specifically updated (mineralization extending up to 300 m at depth); some new N-S striking and cross-cutting fault structures were interpreted (Figures 1, 2, 3 and 4), but further investigations are required to determine their metallogenic relationship to any of the mineralization processes.

The most striking mineralization forms were determined, both for gold and copper, in southern-southeastern parts of the concession area towards so-called Maričanski Rid locality, as the southernmost position there.

We would like to stress out that the actual 3D model of the Plavica polymetallic deposit we can see that it strongly reflects the geochemical Au-Cu pattern already seen at the geochemical cross sections (Serafimovski et al., 2014). The present exploration at the Plavica deposit continues in the western and northwestern parts where exist positive field, structural-morphostructural, geological and geochemical indications for certain mineralization extension. Exploration continues with shallow exploration drill holes (in average up to 100 m).




a)




b)

Fig. 3. 3D model of the Plavica deposit, gold mineralization, resolution scale a) 30 m and b) 10 m.



a)



b)

Fig. 4. 3D model of the Plavica deposit, copper mineralization, precision scale a) 30 m and b) 10 m

CONCLUSION

Latest complex, geological, geochemical and geophysical explorations defined interesting gold and copper mineralization at the Plavica locality followed by significant presence of silver, zinc, lead and some other occasional metals. Direct implications of gold and copper geochemical anomalies, structural features and metallogenetic setting were confirmed by exploration drill holes, which confirmed that mainly gold and copper mineralizations are related to crushing zones, stockworks and disseminations as well as, quartz (massive and

'vuggy' silica) and quartz vein structures that directly reflect structural-tectonic setting of the area.

The gold-copper mineralization of polymetallic character with construction of the 3D model have revealed its isometric to lens/lensoid morphology, which swerves and distorts as depth increases.

It has been proved with the 3D model spatial position of gold and copper mineralization within the Plavica deposit between hypsometric levels 1305 and 600 m.

REFERENCES

- [1] Alderton, H. M. D. and Serafimovski, T.: The geology and genesis of the Plavica copper-gold deposit, Macedonia. *Applied Earth Science (Trans. Inst. Min. Metall. B)* 2007 Vol. **116**, No. 2, p. 94–105 (2007).
- [2] Arvanitidis, N. D., Michael, C., Christidis, C., Weihed, P., Gaal, G., Royer, J. J., Perantonis, G., Bakalis, V. and Ballas, D.: Using 3D/4D modelling tools in exploration for epithermal gold potential areas in Eastern Rhodope zone (Western Trace, NE Greece) – Mineral deposit research for a high-tech world, *12th SGA Biennial Meeting* 2013. *Proceedings*, Volume **1**, Sweden, pp 58–61.
- [3] Barton, P. B., Jr., and Skinner, B. J.: Sulfide mineral stabilities. In: Barnes, H. L., ed.: *Geochemistry of hydrothermal ore deposits*, New York, Holt, Rinehart and Winston, p. 236–333, 1967.
- [4] Bilibajkić, R. and Bilibajkić, D.: Report of gravimetric-geomagnetic studies in the wider zone of the Zlatica-Plavica locality. *Professional fund of the MIC "Sileks" – Kratovo and "Geozavod" – Belgrade*, 1978 (in Macedonian).
- [5] Bogoevski, K and Rakić, S.: Geological features of the Zlatica–Plavica deposit. Геолошке карактеристике лежишта Златица–Планица. *Professional fund of the MIC "Sileks" – Kratovo*, 1985 (in Macedonian).
- [6] Chile's, J. P. and Delfiner, P.: *Geostatistics: Modeling spatial uncertainty*. Ciley, New York: 695, 1999.
- [7] Chmielowski, M. R., Jansson, N., Persson, M. F., Fagerström, P. and Weihed, P.: 3D geochemical modelling of hydrothermal alteration related to 1.89 Ga VHMS-type-deposits, Kristineberg area, Skellefte District, Sweden. *Proceedings of the 12th Biennial SGA Meeting*, Uppsala, Sweden, 12th–16th August 2013, pp. 66–69, 2013.
- [8] Christidis Christosm, Serafimovski Todor, Arvanitidis Nikos, Michael C.: 3D modeling tools jointly applied on Gerakario (Greece) and Kadiica (FYROM) porphyry copper mineralizations. *Proceedings of the XX Congress of Carpathian Balkan Geological Association*, Special Issue, **1** (1). pp. 355–358, 2014.. ISSN 0254-5276; 2306–9600
- [9] Cuthberston, A. S.: The Plavica and Borovic Projects, Macedonia. *Rio Tinto Exploration*, Germany, Report No 80289, 1998.
- [10] Drovenik, M., Pezdic, J. and Rakić, S.: The sulfur isotopic composition of pyrite from borehole ZB-24 Plavica, Zletovo–Kratovo volcanic area. *Rudarsko-metalurški Zbornik*, Vol. **30**, pp 69–76, 1983 (in Slovenian, with English summary).
- [11] Fish, J.: Report on exploration activities, Plavica, R. Macedonia, *Rio Tinto Mining and Exploration*, UK, December 1998.
- [12] Fish, J., Saint, T. and Boast, M.: Report on exploration activities, Plavica, R. Macedonia, *Rio Tinto Mining and Exploration*, UK, March 2000.
- [13] Goovaerts, P.: *Geostatistics for Natural Resources Evaluation*. Oxford University Press, New York, 1997, 496 p.
- [14] Ivanov, T. and Denkovski, G.: Geology of the Plavica–Zlatica deposit, Kratovo–Zletovo. *Proc. 9th Yugoslav Geological Congress*, Sarajevo, Yugoslavia, October 1978, 767–777 (in Macedonian).
- [15] Ivanov, T. and Denkovski, Đ.: Hydrothermal alterations in the Plavica–Zlatica porphyry copper deposit in the Kratovo–Zletovo volcanic area. *Symposia of alteration of rocks and minerals, A 100 years of geological school and science in Serbia*, Faculty of Mining and Geology, University in Belgrade, pp. 139–162, 1980 (in Macedonian).
- [16] Ivanovski, I.: *Preparation of geological map on Plavica using GIS software*. Seminar work, University of Goce Delčev – Štip, 2014, 12 p.
- [17] Janković, S.: *Metallogenic epochs and ore-bearing areas in the Yugoslavia*. Publisher Mining Institute, Belgrade, 1967 (in Serbian).
- [18] Janković, S., Petković, M., Tomson, I. N. and Krasov, V.: Porphyry Copper deposits in the Serbo-Macedonian province, southeastern Europe. In: *European Copper Deposits* by S. Janković and R. H. Sillitoe, SGA-Spec. publ. No. 1, UNESCO-IGCP Projects No. 169 and 60, 1980.
- [19] Karamata, S., Krstić, B., and Stojanov, R.: Terranes from the Adriatic to the Moesian Massif in the Central part of the Balkan Peninsula. *Terra Nova*, **4**, Abstract supplement, **2**, 78, 1992.
- [20] Ligoovski, M., Serafimovski, T., Tasev, G. and Stefanova, V.: Geochemical study and 3D modelling of the Kazan

- Dol copper deposit, Republic of Macedonia. *Geologica Macedonica*, **28** (1). pp. 7–17 (2014). ISSN 0352–1206.
- [21] Madera, A.: *Geology and mineral resource potential within the Cumbio FTAA of WMC Philippines*, Inc, unpublished WMC internal report, 2000, 25 p.
- [22] Marković, M.: Contribution to the knowledge of volcanic morphology of the Kratovo–Zletovo area. *Geological Annals of the Balkan Peninsula (Annales géologiques de la Péninsule balkanique)*, Belgrade, 1971, 36 p. (in Serbian).
- [23] Matheron, G.: *The Theory of Regionalized Variables and its Applications*. Ecole des Mines, Fontainebleau, France, 1971.
- [24] Mijalković, N. and Pešić, D.: Geological and petrological features of the NE part of the Kratovo–Zletovo area. *Papers of the VI counseling of geologists of Yugoslavia, Ohrid*, part II, pp.158–170, 1966 (in Serbian).
- [25] Mudrinić, Č.: Concentration and distribution of rare and precious metals in the Plavica–Zlatica deposit. Study, *Professional fund of MIC “Sileks”– Kparamovo*, 1982 (in Serbian).
- [26] Mudrinić, Č.: Rare and precious metals in the Cu-poly-metallic deposit Plavica–Zlatica (Eastern Macedonia). *XI Congress of geologists of Yugoslavia, Tara*, 1986 (in Serbian).
- [27] Pantić, R., Simić, V., Mijalković, N., Koprivica, D., Lolević, S. and Radusinović, D.: Kratovo–Zletovo exploration and potentiality (Final Report). *Professional Documentation Fund of Geoinstitute Belgrade*, 1972 (in Serbian).
- [28] Pendžerkovski, J., Đuzelkovski, D. and Gavrilova, N.: Geological-petrographic report of the field adjacent to the Zletovo mine. *Professional Documentation Fund of Geozavod, Skopje*, 1960 (in Macedonian).
- [29] Percival, T. J.: *Preliminary geological report on the Plavica property*, Yugoslavia – Nassau Limited. USA, 1992.
- [30] Petković, M. and Romić, K.: Structural-volcanological study with determination of erosion level “Plavica”. *Professional fund MIC “Sileks” – Kratovo*, 1977 (in Macedonian).
- [31] Rakić, S.: Some structural and genetic features of the polymetallic mineralization within the Zletovo ore field. *IX Congress of geologists of Yugoslavia, Sarajevo*, pp. 611–619, 1978 (in Serbian).
- [32] Rakić, S.: Secondary quartzites of the central part of the Kratovo–Zletovo area-important raw resources related to important occurrences of polymetallic and raw mineralization. *X Jubilee Congress of geologists of Yugoslavia, Budva*, 1982 (in Serbian).
- [33] Ristić, M. and Klajn, G.: Geological, geochemical and metallogenetic features of ore occurrences at Plavica–Zlatica in SR Macedonia. *Papers of Geological Survey of SRM*, Vol. **16**, pp. 213–229, 1973, Skopje (in Serbian).
- [34] Serafimovski, T.: *Metallogeny of the Lece-Chalkidiki zone*. Doctoral thesis, Faculty of Mining and Geology Štip, 380 p. 1990 (in Macedonian).
- [35] Serafimovski, T.: *Structural-metallogenic features of the Lece-Chalkidiki zone: Types of deposits and reonization*. Faculty of Mining and Geology, Special Issue No. 2, 1993, 235 p. (in Macedonian).
- [36] Serafimovski, T. and Rakić, S.: The secondary quartzites hosting gold mineralization in the Crni Vrv–Plavica volcanic area. *Geologica Macedonica*, Vol. **12**, p. 9–21 (1998).
- [37] Serafimovski, T. and Rakić, S.: New Geochemical data concerning gold related to silification zones in the Plavica volcanic structure, Eastern Macedonia. *Mineral Deposits: Processes to Processing*, Stanley et al. (eds). 1999 Balkema, Rotterdam, p. 585–588 1999..
- [38] Serafimovski, T., and Tasev, G., 2003. The Plavica epithermal polymetallic system: mineralogical and geochemical data. *Mineral Exploration and Sustainable Development*, Eliopoulos et al. (eds), **1** (1). pp. 523–526. ISSN 90 77017 77 1.
- [39] Serafimovski, T. and Tasev, G.: *Ore-microscopic study of samples from the Plavica deposit*. Department of Mineral deposits, Faculty of Natural and Technical sciences, University “Soce Delčev”– Štip, 411 p. 2013 (in Macedonian).
- [40] Serafimovski, T., Aleksandrov, M., Tasev, G., Stefanova, V., Paterson, J., Stevanović, M., Ristovski, V. and Ivanovski, I.: *Elaborate of detailed geological study, with calculation of geological ore reserves, of copper and gold in the Plavica, Kratovo locality*. Department of Mineral deposits at the Faculty of Natural and Technica Sciences, University “Goce Delcev” for Genesis Resources International DOOEL, Skopje, 431 p. 2014 (in Macedonian).
- [41] Sillitoe, H. R. and Hedenquist, W. J.: Linkages between Volcanotectonic Settings, Ore-Fluid Compositions, and Epithermal Precious Metal Deposits. *Society of Economic Geologists*, Special Publication 10, pp. 315–343 (2003).
- [42] Stefanova, V.: *Eluvial-deluvial gold occurrences related to the Tertiary magmatism in the Republic of Macedonia*. Doctoral thesis, University “Ss. Cyril and Methodius”, Skopje, 198 p., 2005 (in Macedonian).
- [43] Stefanova, V., Serafimovski, T. and Nedelkov, R.: Mineralogical and chemical characteristics of the most important minerals in the volcanic rocks of the Kratovo–Zletovo volcanic area. *Geologica Macedonica*, Vol. **21**, pp. 1–10 (2007).
- [44] Stefanova, V., Volkov, V. A., Serafimovski, T. and Sidorov, A. A.: Native Gold from the Plavica Epithermal Deposit, Republic of Macedonia. *Doklady Earth Sciences*, Vol. **451**, Part 2, pp. 818–823 (2013). © Pleiades Publishing, Ltd., (ISSN 1028–334X).
- [45] Stevanović, M.: *Plavica high sulfidation epithermal system*. Internal report, Genesis Resources International DOOEL Skopje, 2014, 24 p.
- [46] Stojanov, R.: Mineralogical composition of alterations at the Plavica in the Kratovo–Zletovo area. *VIII Yugoslavian geological Congress, Ljubljana*, Vol. **5**, pp. 42–54 (1974).
- [47] Stojanov, R.: Volcanic cupola Plavica and polymetallic mineralizations and alterations within. *Symposia of alteration of rocks and minerals, a 100 years of geological school and science in Serbia*, Faculty of Mining and Geology, University in Belgrade, pp. 139–162 (1980) (in Macedonian).

- [48] Stojanov, R., Serafimovski, T.: Volcanism in the Kratovo–Zletovo area. *XII Congress of geologists of Yugoslavia, Ohrid*, Vol. 2, pp. 424–441 (1990) (in Macedonian).
- [49] Stojanov, R., Serafimovski, T., Boev, B., Rakić, S.: Magmatism and Metalogeny in the Northern Parts of the Zletovo–Kratovo ore District. *International Workshop, UNESCO-IGCP Project 356, 5–7 April, 1995, Štip*, p. 41–44.
- [50] Tomson, I. N., Serafimovski, T. and Kochneva, N.: Cenozoic metallogeny in the Eastern Macedonia. *Geology of Ore Deposits*, Moscow, Vol. 40, No. 3, pp. 195–204 (1998) (in Russian).
- [51] Volkov, V. A., Serafimovski, T., Stefanova, V. and Sidorov, A. A.: Formation Mechanism of Dwarfish Cu–Au Porphyry Deposits of Macedonia. *Doklady Earth Sciences*, 2010, Vol. 431, Part 2, pp. 429–434. (Original Russian Text: A. V. Volkov, T. Serafimovski, V. Stefanova, A. A. Sidorov, 2010, published in *Doklady Akademii Nauk*, 2010, Vol. 431, No. 5, pp. 649–655) 2010.
- [52] White, N. C., Leake, M. J., McCaughey, S. N., and Parris, B. W.: Epithermal deposits of the southwest Pacific: *Journal of Geochemical Exploration*, V. 54, p. 87–136 (1995).
- [53] Zlatkov, G., Tasev, G., Stefanova, V., Bogdanov, K. and Serafimovski, T.: Composition of some major mineral phases from the Plavica epithermal gold deposit, Eastern Macedonia. *Geologica Macedonica*, 28 (2). pp. 149–163 (2014). ISSN 0352–1206.

Резиме

3D МОДЕЛИРАЊЕ НА Au-Cu ПОЛИМЕТАЛИЧНОТО НАОЃАЛИШТЕ ПЛАВИЦА,
РЕПУБЛИКА МАКЕДОНИЈАИгор Ивановски¹, Далибор Серафимовски², Горан Тасев², Тодор Серафимовски¹Генезис Ресурси Интернешнл ДООЕЛ Скопје, Р. Македонија²Факултет за природни и технички науки, Институт за геологија, Универзитет „Гоце Делчев“,
Бул. Гоце Делчев, 89, 2000 Штип, Република Македонија
icestip@yahoo.com**Клучни зборови:** 3D модел; Плавица; злато, бакар; епитермално

Најновите истражувања и проучувања на епитермалното наоѓалиште на висока-сулфидизација Плавица покажаа значителен напредок во разбирањето на геологијата и дефинирањето на конкретното рудно тело. Резултатите од дупчотините направени од Генезис Ресурси Интернешнл ДООЕЛ Скопје, Република Македонија во 2011, 2012 и 2013 покажаа дека постојат: 51 метарски интервал со 3,9 g/t Au во зоната на оксидација, 65 метарски интервал со 3,1 g/t Au повторно во зоната на оксидација, како и 51 метарски интервал со 3,7 g/t Au, 2,8% Cu и 16 g/t Ag во сулфидната зона. За конструирањето на овој 3D модел беа користени сите 195 дупчотини направени од Генезис Ресурси Интернешнл. Дупчотините биле направени во пери-

одот 2011–2013, во вкупна должина од 47295,8 m од кои 74 дупчотини биле направени со методата на дијамантско јадро и вкупна должина од 23908,3 m додека останатите 121 дупчотини биле изработени со методата на реверсна циркулација (РЦ) во вкупна должина од 23387,5 m. Со употреба на професионалниот софтвер ArcGIS беа изработени 3D модели на рудното тело за злато и бакар, во соодветни резолуции од 30 m и 10 m, за секој елемент, а што ни го дефинираше простирањето на минерализацијата од 1305 m до 600 m хипсометриско ниво. Овие модели би требало да го подобрат нашето разбирање за минерализацијата и да ги надополнат геохемиските податоци за наоѓалиштето Плавица.