

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/277332756>

# Prognostic value of normal myocardial perfusion imaging in asymptomatic diabetic patients with moderate and high calcium score SCORE

CONFERENCE PAPER · MAY 2015

DOI: 10.1093/ehjci/jev05

---

DOWNLOADS

11

---

VIEWS

18

6 AUTHORS, INCLUDING:



**Marijan Bosevski**

Ss. Cyril and Methodius University

77 PUBLICATIONS 113 CITATIONS

SEE PROFILE



**Marija Vavlukis**

Ss. Cyril and Methodius University

66 PUBLICATIONS 37 CITATIONS

SEE PROFILE

## European Heart Journal - Cardiovascular Imaging

[About](#)
[Submit](#)
[Subscribe](#)
[Advertise](#)
[ESC journals](#)
[Jobs](#)


Editor-In-Chief  
G. Maurer

Published on behalf of

IMPACT FACTOR 3.669

5 YEAR IMPACT FACTOR 3.669

SI: CARDIAC &  
CARDIOVASCULAR  
SYSTEMS

37 out of 125



[Advanced](#)

[Browse all](#)
[16: suppl 1](#)
[16:7](#)
[Current](#)
[Advance](#)
[access](#)
[◀ Previous article](#)
[Next article ▶](#)

#### Article

[Abstract](#)
[Abstract](#)
[Abstract](#)
[Abstract](#)
[Abstract](#)
[Abstract](#)
[Abstract](#)

## Poster Session 3

P344End-stage renal failure: prevalence and predictors of left

# Prognostic value of normal myocardial perfusion imaging in asymptomatic diabetic patients with moderate and high calcium score SCORE

## Abstract

**Background:** To evaluate the intermediate prognostic value of normal myocardial perfusion imaging in asymptomatic diabetic patients with high cardiovascular (CV) risk with intermediate and high coronary calcium score (CAC).

**Methods:** 115 asymptomatic patients with high CV risk without known coronary artery disease (CAD), underwent SPECT myocardial perfusion imaging (MPI) after multislice computer tomography with coronary calcium score (CAC) assessment for detection of suspected CAD was. Presence of traditional risk factors, diabetes control and presence of albuminuria were evaluated. 75 patients with normal MPI results were included in the study. 17 segment model for myocardial perfusion and function analysis was used. Patients were divided into three groups: I gr- 20 patients with diabetes duration between 1-5 years; II gr- 24 patients with diabetes duration 5-10 years; III gr- 31 patients with diabetes duration >10 years. End points (cardiac death, non fatal myocardial infarction, heart failure, new angina, revascularization) at 6 months, 1 and 2 years were followed.

**Results:** All patients have normal rest left ventricular function with EF >50% and normal myocardial perfusion scans. 40 patients had moderate calcification of the coronary arteries with average CAC 290+/-95. 35 patients had severe coronary calcification with average CAC 568+/-67. There was correlation of diabetes duration with CAC severity (r=0.62 for diabetes duration over 10 years). Only 3 cardiac events were registered at 24 month follow up in the II and III group (new angina with percutaneous coronary (PCI) revascularization). Hard events (cardiac death, nonfatal myocardial infarction STEMI/NSTEMI) were 0% in gr I, 4.2% in gr II and 6.4% in Gr III. Overall rate of CV at 24 month follow up was 4.0%.

**Conclusions:** MPI and CAC are valuable techniques for preclinical assessment of CAD in asymptomatic high risk patients, which help guide optimal treatment decision and prognosis. Diabetic patients even with normal MPI have increased intermediate risk for CV events.