

*Multivariate extraction of dominant geochemical markers for deposition of 69 elements in the Bregalnica River basin, Republic of Macedonia (moss biomonitoring)*

**Biljana Balabanova, Trajče Stafilov,  
Robert Šajn & Claudiu Tănăselia**

**Environmental Science and Pollution  
Research**

ISSN 0944-1344

Volume 23

Number 22

Environ Sci Pollut Res (2016)

23:22852–22870

DOI 10.1007/s11356-016-7502-7



 Springer

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".



RESEARCH ARTICLE

# Multivariate extraction of dominant geochemical markers for deposition of 69 elements in the Bregalnica River basin, Republic of Macedonia (moss biomonitoring)

Biljana Balabanova<sup>1</sup> · Trajče Staflov<sup>2</sup> · Robert Šajn<sup>3</sup> · Claudiu Tănăsescu<sup>4</sup>

Received: 9 February 2016 / Accepted: 22 August 2016 / Published online: 27 August 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Abstract** Atmospheric deposition was investigated using the terrestrial moss species *Hypnum cupressiforme* (Hedw.) and *Homothecium lutescens* (Hedw.) in the Bregalnica River basin, Republic of Macedonia. Long-term emission occurs in this area due to the hydrothermal exploitation of Pb–Zn deposits (Sasa and Zletovo mines) and copper ore exploitation and floatation (Bučim mine). Determination of the chemical elements was conducted using atomic emission spectrometry with inductively coupled plasma (ICP-AES) and mass spectrometry with inductively coupled plasma (ICP-MS). A combination of multivariate techniques (PCA, FA and CA) was applied for data processing and identification of element association with lithogenic/anthropogenic origin. Seven dominant factors were extracted from the total of 69 analysed elements. Spatial distribution maps were constructed for the determination and localisation of smaller areas with higher contents of certain anthropogenic elements. In this way, the influences of

selected human activities on local air pollution can be determined. The summarised data show quantification of the element distributions. This not only allows the determination of the distribution of hazardous elements but also presents complete characterisation of element deposition in the environs of mines.

**Keywords** Air pollution · Moss · Biomonitoring · Atomic emission spectrometry with inductively coupled plasma · Mass spectrometry with inductively coupled plasma · Multivariate assessment · Spatial distribution · Bregalnica River basin

## Introduction

Environmental pollution at hazardous levels for living organisms presents a global problem and a challenge for a macro-scale monitoring. Some sub-disciplines have been developed over time in order to consider realistic environmental conditions. Arguably, the understanding of atmospheric pollution is one of the most emergent areas of environmental science (Fernández et al. 2015). Atmospheric pollution represents solutions or suspensions of minute amounts of harmful compounds in the air (Vallero 2014). The degree and extent of environmental changes over the last decades have given a new urgency and relevance to the detection and understanding of environmental changes due to human activities, which have altered global biogeochemical cycling of heavy metals and other pollutants (Athar and Vohra 1995; Acton 2013). Monitoring of toxic air pollutant is needed in order to understand their spatial and temporal distribution and ultimately to minimise their harmful effects. In addition, to direct physical and chemical methods of air pollution monitoring, bioindication has also been used to evaluate air pollution risk (Abol et al. 2010; Ares et al. 2012; Vallero, 2014).

Responsible editor: Céline Guéguen

**Electronic supplementary material** The online version of this article (doi:10.1007/s11356-016-7502-7) contains supplementary material, which is available to authorized users.

✉ Trajče Staflov  
tmajest@pmfukim.mk

<sup>1</sup> Faculty of Agriculture, University “Goce Delčev”, Kite Mirov  
bb, Štip, Republic of Macedonia

<sup>2</sup> Institute of Chemistry, Faculty of Natural Sciences and Mathematics,  
St. Cyril and Methodius University, POB 162,  
1000 Skopje, Republic of Macedonia

<sup>3</sup> Geological Survey of Slovenia, Dimičeva ulica 14,  
1000 Ljubljana, Slovenia

<sup>4</sup> INCDO-INOE 2000 Research Institute for Analytical  
Instrumentation (ICIA), Cluj-Napoca, Romania