Българско геологическо дружество Bulgarian Geological Society



НАЦИОНАЛНА КОНФЕРЕНЦИЯ

с международно участие

NATIONAL CONFERENCE

with international participation

ГЕОНАУКИ GEOSCIENCES 2016

SHORT COMMUNICATIONS НАУЧНИ СЪОБЩЕНИЯ

Българско геологическо дружество Bulgarian Geological Society Sofia • 2016

ГЕОНАУКИ 2016

Рецензирани научни съобщения от Националната конференция е международно участие

Редакционна колегия: Йодо Янев (главен редактор),

Елеонора Балканска (секретар), Михаил Тарасов, Василка Младенова, Борис Вълчев, Страшимир Страшимиров, Радослав Наков, Димитър Синьовски, Елена Колева-Рекалова, Пламен Иванов, Стефан Шанов, Златка Чернева, Евгения Тарасова

Технически редактор и компютърно оформление – Яна Цветанова Коректор – Надежда Велинова

Организационният комитет благодари на Б. Каменов, А. Бендерев, М. Янева, В. Христов, Р. Недялков, М. Йосифова, П. Паклишина, Д. Димитрова и З. Милаковска за помощта при рецензирането на изучните съобщения.

GEOSCIENCES 2016

Peer reviewed short communications of the National Conference with international participation

Editorial board: Yotzo Yanev (Editor-in-Chief),

Eleonora Balkanska (Secretary), Mihail Tarassov, Vassilka Mladenova, Boris Valchev, Strashimir Strashimirov, Radoslav Nakov, Dimitar Sinnyovsky, Elena Koleva-Rekalova, Plamen Ivanov, Stefan Shanov, Zlatka Chemeva, Eugenia Tarassova

Technical and design editor: Yana Tzvetanova

Corrector: Nadezhda Velinova

Издателетво Българско геологическо дружество, 2016 г.
Отвечатаво в БУЛГЕД ООД



Nannofossil assemblages from the Paleogene Nemanjici section, Republic of Macedonia

Нанофосилни асоциации в палеогенския профил Неманици, Република Македония

Violeta Stojanova, Goše Petrov Виолета Стоянова, Гоше Петров

University "Goce Delcev", Faculty of Natural and Technical Sciences, Štip, Republic of Macedonia; E-mails: violeta.stojanova@ugd.edu.mk; gose.petrov@ugd.edu.mk

Keywords: foraminifers, nannofossils, Paleogene sediments, Ovče Pole Basin.

Introduction

Nemanjici section belongs to the Ovče Pole Paleogene Basin, the Republic of Macedonia. This basin is a large Paleogene sedimentary mass with NW-SE trend that is superimposed on varied rocks from the eastern part of the Vardar Zone in the territory of the Republic of Macedonia (Fig. 1).

According to the current knowledge, the Paleogene sediments of the Ovče Pole basin are nearly 3.5 km thick. Four lithostratigraphical units (basal unit, lower flysch unit, unit of yellow sandstones, upper flysh unit) have been recognized, as their lithology, thickness and spatial relationships were discussed in several works (Maksimovič et al., 1954; Stojanova, 2008; Stojanova et al., 2011, 2012; Stojanova, Petrov, 2014,




Fig. 1. Distribution of Paleogene sediments in the Ovče Pole basin, Republic of Macedonia: 1, Paleogene sediments; 2, Ovče Pole basin; 3, tectonic boundary; SMM, Serbian-Macedonian Massif; VZ, Vardar Zone; WMZ, Western Macedonian Zone; 4, Paleogene cross-section

2016). In the Nemanjici section benthic and planktonic foraminifers have been recently found (Stojanova, 2008; Stojanova et al., 2013; Valchev et al., 2013). Based on benthic foraminifera Bolivina antegressa Subzone of the Planulina costata Zone was divided (Stojanova, 2008), and later on Early Oligocene age for the Nemanjici section was determined on the base of planktonic foraminifers (Juranov in: Stojanova et al., 2013).

Nemanjici section is located 6.5 km NE of the town of Sveti Nikole, and it comprises 90–100 m of the upper flysch unit, represented by greenish calcareous clay alternating with thin sandstone beds. 18 samples (N1 to N18) were taken from the section, and the sampling was performed in marly-clay layers at a distance of 2–3 m between the samples. The overall procedure for allocating nannofossils was performed with standard processing methods, which include making microscopic preparations with Canada balsam. Paleontological determination were made under JENAPOL-d light microscope with magnification x 2000.

Results and discussion

Nannofossil research revealed that the samples contain well preserved specimens of 15 species including: Blackites spinosus (Deflandre & Fert), Braarudosphaera bigelowii (Gran & Braarud), Chiasmolithus grandis (Bramlette & Riedel), Coccolithus formosus (Kamptner), Coccolithus pelagicus (Wallich), Cyclicargolithus floridanus (Roth & Hav), Helicosphaera compacta Bramlette & Wilcoxon, Lanthernites minutus Stradner, Pontosphaera multipora (Deflandre & Fert), Pontosphaera plana (Bramlette & Sullivan), Reticulofenestra bisecta (Hay, Mohler and Wade), Reticulofenestra reticulata (Gartner & Smith), Reticulofenestra stavensis (Levin & Joerger), Zygrhablithus bijugatus (Deflandre and Fert). By analyzing the stratigraphic range of individual species, composing the nannofossil association, Nemanjici

section is referred to NP 18 nannofossil biozone (Martini, 1971), and thus the sediments belong to the Upper Eocene (Priabonian). In samples 13, 16 and 17 *Watznaueria barnesiae* (Black & Barnes) from the Upper Cretaceous was found. Most probably it was resedimented from the surrounding Cretaceous cliffs, during the processes of their erosion and the deposition of the Paleogene sediments.

References

- Maksimovič, B., B. Sikošek, O. Markovič, M. Veselinovič. 1954. Geologic composition and tectonic structure of a part of Ovče Pole and Tikveš with paleontological documentation. – Geological Survey Papers, 4. R. Macedonia, 1–177 (in Macedonian).
- Martini, E. 1971. Standard Tertiary and Quaternary calcareous nanoplankton zonation. – In: *Proc. II Plankt. Conf.*, 2. Roma, 739–785.
- Stojanova, V. 2008. Evolution and Stratigraphy of the Paleogene in the Territory of the Republic of Macedonia. PhD Thesis. Štip, 196 p. (in Macedonian).
- Stojanova, V., G. Petrov. 2014. Biostratigraphic correlation of the Paleogene sections in the Ovče Pole Basin, Republic of Macedonia. – In: *Proceedings of National Conference* "GEOSCIENCES 2014". Sofia, BGS, 67–68.
- Stojanova, V., G. Petrov. 2016. Foraminiferal biostratigraphy of Upper Eocene sediments from the Ovče Pole basin, Republic of Macedonia. – Geologica Macedonica, 30, 1, 65–77.
- Stojanova, V., G. Petrov, V. Stefanova. 2011. Small foraminifers from the Paleogene basins in the Republic of Macedonia. – In: Proceedings of National Conference "GEOSCIENCES 2011". Sofia, BGS, 93–94.
- Stojanova, V., G. Petrov, V. Stefanova. 2012. Biostratigraphy of the Ovce Pole Paleogene basin, R. Makedonija. *Geologica Macedonica*, 26, 2, 53–63.
- Stojanova, V., B. Valchev, S. Juranov. 2013. Palaeogene planktonic foraminifera of the Republic of Macedonia. *C. R. Acad. Bulg. Sci.*, 66, 5, 717–724.
- Valchev, B., V. Stojanova, S. Juranov. 2013. Paleogene hyaline benthic foraminifera (LAGENINA and ROTALIINA) from the Republic of Macedonia. – Rev. Bulg. Geol. Soc., 74, 1–3, 81–110.