

**GOCE DELCEV UNIVERSITY - STIP
FACULTY OF AGRICULTURE**



JOURNAL OF AGRICULTURE AND PLANT SCIENCES

YEAR 2017

VOLUME 15, Number1/2

**GOCE DELCEV UNIVERSITY - STIP, REPUBLIC OF MACEDONIA
FACULTY OF AGRICULTURE**

UDC 63(058)
ISSN 2545-4447 print
ISSN 2545-4455 on line



**Journal of Agriculture and Plant Sciences, JAPS, Vol 15
Successor of the Yearbook of Faculty of Agriculture of GDU, Vol 14**

YEAR 2017

VOLUME XV, Number 1/2

CONTENT

Emilija Arsov, Galina Ivanova, Sasa Mitrev, Multigene characterization of ' <i>Candidatus phytoplasma solani</i> ' in pepper and tomato plants in the Republic of Macedonia	7
Biljana Balabanova, Trajče Stafilov, Robert Šajn, Claudiu Tănăselia Bioindication ability of <i>Hypnum cupressiforme</i> and <i>Homolothecium lutescens</i> for dermination of arsenic distribution in environment	15
Olivera Bicikliski, Krste Tashev, Fidanka Trajkova, Ljupco Mihajlov, Liljana Koleva Gudeva Comparative analysis of capsaicin content in peppers (<i>Capsicum annum</i> L.) grown in conventional and organic agricultural systems	27
Zoran Dimitrovski Inspection of pesticide application equipment	37
Zoran Dimitrovski, Dimitrov Sasko, Kukutanov Risto Condition of air assisted sprayers in Shtip region and possibility of applying European standard EN 13790	45
Violeta Dimovska, Fidanka Ilieva, Sanja Kostadinovic, Ljupco Mihajlov Physical and chemical characteristics of pomegranate fruit (<i>Punica granatum</i> L.), of cv. Karamustafa	53
Sanja Filipovska, Darko Andronikov, Aco Kuzelov Chemical and fatty acid composition in meat of young chickens different hybrid lines	61
Natasa Gunova, Dusan Spasov, Biljana Atanasova, Dragica Spasova, Mite Ilievski Correlation between population dynamics of <i>Tuta absoluta</i> (Lepidoptera: Gelechidae) and climate, at tomato in protected area	69
Verica Ilieva, Natalija Markova Ruzdik, Ilija Karov, Ljupcho Mihajlov, Mite Ilievski, Biljana Kovacevik Genetic variability for yield and some yield-related traits in rice (<i>Oryza sativa</i> L.)	75
Dijana Indzhelieva, Katja Velkova-Jorgova, Darko Andronikov, Aco Kuzelov The influence of starter culture of lactic- acid bacteria and bifid bacteria over the sanitary- hygienic, sensor and physical – chemical indicators on the re – boiled – smoked durable sausage	81
Viktorija Maksimova, Liljana Koleva Gudeva, Rubin Gulaboski, Maja Shishovska, Zorica Arsova Sarafinovska Capsaicin and dihydrocapsaicin variability in <i>Capsicum</i> sp. cultivars from Republic of Macedonia revealed by validated HPLC method	89
Ivana Velesanova, Fidanka Trajkova, Liljana Koleva Gudeva Micropropagation of ornamental species <i>Brassica oleracea</i> cv. Kyoto red given and <i>Ageratum</i> sp.	97



PHYSICAL AND CHEMICAL CHARACTERISTICS OF POMEGRANATE (*Punica granatum* L.), CULTIVAR KARAMUSTAFA

Violeta Dimovska¹, Fidanka Ilieva¹, Sanja Kostadinovic¹, Ljupco Mihajlov¹

¹Faculty of Agriculture, Goce Delcev University, Krste Misirkov, 10-A, 2000 Stip, Republic of Macedonia

violeta.dimovska@ugd.edu.mk; fidanka.ilieva@ugd.edu.mk; sanja.kostadinovic@ugd.edu.mk;

ljupco.mihajlov@ugd.edu.mk

Abstract

Pomegranate is fruit species well-adapted to arid and semi-arid conditions and highly valued crop widely cultivated in Mediterranean and Near Asia countries. The most important pomegranate cultivars in Macedonia are "Bejnarija", "Lifanka" and "Karamustafa". The objective of this study was to evaluate physical and chemical parameters of pomegranate cultivar "Karamustafa", collected from two different locations of the R. Macedonia. The physical parameters include fruit fresh weight, percentage of grain (%), weight of 100 berries (g), and percentage of juice (%). The cultivar "Karamustafa" (239.57- 304.79 g) belongs to a group of large fruits. The yield juice ranged from 47.83 % (Valandovo) to 50.63 % (Josifovo). Based on the content of the total acids (TA), the cultivar "Karamustafa" belongs to a group of sweet cultivar (TA<1).

Based on the results we found significant differences in the weight of the fruit, the content of the total acids and the content of the total polyphenols in the juice.

Key words: *Punica granatum* L., fruits, juice, total acids, total polyphenols

INTRODUCTION

The pomegranate (*Punica granatum* L.) belongs to the *Punicaceae* family, the genus *Punica* L. (Fond Quer, 1979). It is a diploid species whose somatic number is $2n=16$, and haploid chromosomes = 8 (Westwood, 1982) or $2n=16$ or 18 (Mars, 1998).

Pomegranate (*Punica granatum* L.) is an important commercial fruit crop that is extensively cultivated in part of Asia, North Africa and the more Mediterranean regions. All parts of the tree (leaves, flowers, and roots) are used for medical purposes for centuries (Godwa et al., 2009).

The total area of pomegranate cultivation in the world is well above 300,000 ha, which, more than 76% is found in these countries: India, Iran, China, Turkey and the USA (Melgarejo et al., 2012) with production higher than 3,000,000 t. The fruits are consumed fresh or processed into juice, jams, syrup and sauce (Fawole et al., 2011).

Consumption of pomegranate fruit and juice in the world is increasing very fast, because of its health benefits, high antioxidant capacity

and the high content of polyphenols and anthocyanins (Zarei et al., 2010). Pomegranate juice is extremely nutritious, it has medicinal properties that include: sugars, citric and boric acid, pectin, vitamins, iodine, iron and other useful ingredients (Zarei et al., 2011).

Recent studies suggest that pomegranate juice contains anticancer, antimicrobial and antiviral components (Schwartz et al., 2009; Reddy et al., 2007; Kotwal, 2007). Pomegranate fruits' maturity is estimated by the skin colour, juice colour, and juice acidity (Al-Said et al., 2009), all of which vary by cultivar. Pomegranate fruits can be harvested when they reach a suitable size and the colour of skin achieves the desired pigment. The main parameters of maturity are TSS (total soluble solids), TA (total acids) and TSS/TA ratio (Pekmezci and Erkan, 2004). In the production of pomegranate pesticides and fertilizers are not used which means that fruits represent a very healthy food rich in natural nutritional and medicinal properties (Hassan et al., 2012).

In Macedonia, the pomegranate is grown on an area of 50 ha, and it has 700 tons annual production. The most of the area are orchards and some individual trees in the gardens. The pomegranate is traditionally grown in Valandovo region, which is under the influence of Mediterranean climate. In this region, climatic soil conditions (length of 240 vegetation days, annual average temperature > 14,8 C°, the annual temperature sum > 5.200 C°, soils fertile, etc.), enable the successful cultivation of

pomegranate, which is present in this area for centuries.

To the best of our knowledge there are no published results about the determination of physical, morphological and chemical properties of pomegranate fruits, arils and juices in the R. Macedonia. Therefore, the main object of this study was the determination of physical, morphological and chemical parameters of pomegranate cultivar "Karamustafa", grown in two different location of R. Macedonia.

MATERIALS AND METHODS

Pomegranate fruits of "Karamustafa" species were collected from pomegranate orchards located in 10 year old orchard in Valandovo and 20 year old orchard in Josifovo area in R. Macedonia. Trees were planted at the spacing of 2.5 x 3.0 m.

Eight fruits of three different flower levels or a total of 24 per cultivar from Josifovo and Valandovo location were taken for analysis (Tab. 1, Tab. 3; I, II and III repetition).

Physical characteristics

Fruit weight (g) and the weight of 100 arils (g) were measured using the analytical laboratory scale (Mettler Toledo XS). The percentage of peel (%) is calculated as the ratio between the peel weight and fruit weight multiplied by 100. The juice yield (%) is the squeezed juice (ml) of 100 g of fruit. Fruit color was determined by organoleptic method (Koppel et al., 2010). The juice analyzed the pH value, total soluble solids, the total acid content expressed in citric acid, the maturity index and the total phenolic compounds of the juice.

Statistical processing of the results of all

three repetitions was performed. The correlation is represented by the statistical analysis of the results, using the software package for statistical data processing, SPSS 19.

Chemical analyses

Measurements were performed on fresh aril juice. Total soluble solids (°Brix) and salinity (%) were determined by using a digital refractometer (Krüss optronic DR 301-95) at 20°C, calibrated by using water distilled. Titratable acidity was estimated in titration with 0.1 N NaOH to the titration point of pH 8.3, monitored with a pH meter and expressed as citric acids content (g/L). Maturity index was calculated by dividing total soluble solids to titratable acidity. Juice's pH is measured with a pH meter ((100 ATC). Total polyphenolic content was measured by using the Folin-Ciocalteu method (Makkar et al., 1993).

The obtained results are processed by analyzing the variance using the programme package STATVIEW (SAS Institute Version 5.0). The differences between the mean values tested are LSD test for $p \leq 5$.

RESULTS AND DISCUSSION

The physical characteristics of the pomegranate fruit are presented in Table 1. From the results presented in the study, the fruits from the first repetition (I-first level of flowers) of "Karamustafa" cultivar from both locations (Valandovo, Josifovo) were the largest. On the other hand, the lowest fruits were determined from the third repetition (III level of flowers). In general, the average weight of fruits is 239.57 g in Valandovo and 304.79

g in Josifovo. According to the descriptor list for pomegranate (Bellini and Giordani, 1998), they belong to the group of large fruit cultivars (225-375 g). The difference in the weight of the fruit between the location is due to the age structure of the trees. The juice yield (%) varies from 47.83% in location Valandovo to 50.63% in Josifovo location, which is almost identical to some Croatian cultivars, 43.18-53.99% (Radunic and Gadze, 2011).

Table 1. Physical characteristics of pomegranate fruit of cultivar “Karamustafa” at the two locations.

Location	Repetition	Fruit weight (g)	Weight of 100 arils (g)	Juice yield (%)	Percentage of peel (%)	Color of fruit
Valandovo	I	285.43	27.86	48.26	42.14	Reddish yellow
	II	235.00	27.86	46.48	35.56	
	III	198.28	29.28	48.76	37.10	
	I-III	239.57	28.33	47.83	38.27	
Josifovo	I	421.25	38.75	50.36	33.68	Reddish yellow
	II	251.25	36.25	53.68	34.82	
	III	241.88	33.75	47.85	40.83	
	I-III	304.79	34.58	50.63	36.44	

The fruits of “Karamustafa” variety from the two sites were red in color, which is a characteristic of the variety.

The results of the correlation between the weight of the fruit, the weight of 100 arils, the percentage of peel and the juice yield were presented in Table 2.

Table 2. Pearson’s coefficient of correlation between fruit weight and weight of 100 arils, percentage of peel and yield of juice

Pearson’s coefficient of correlation (ρ)					
Valandovo			Josifovo		
Fruit weight / Juice yield	Fruit weight / Percentage of peel	Fruit weight / Weight of 100 arils	Fruit weight / Juice yield	Fruit weight / Percentage of peel	Fruit weight / Weight of 100 arils
-,119	,791	-,817	-,034	,659	-,888

According to the Pearson correlation coefficient, we can conclude that there is a high correlation between the parameters. Negative values denote inverse proportional connection.

Among the physical parameters of the “Karamustafa” variety from the first location (Valandovo) shows a high correlation value. Both cultivars are highly dependent on the weight of the fruit and the percentage of the bark. The cultivar of the Valandovo location, $\rho =$,

791, while in the variety of the Josifovo location $\rho =$, 659. The dependence between the weight of the fruit and the juice yield is the lowest and the same is inversely proportional. The variety of the Valandovo is $\rho =$, 119, while in the variety of the second location $\rho =$, 034. Similarly, there is inverse relationship between the weight of the fruit and the weight of 100 arils. The variety of the Valandovo is $\rho =$, 817, while for the variety of the Josifovo location $\rho =$, 888.



Figure 1. Pomegranate “Karamustafa” (Valandovo).



Figure 2. Pomegranate “Karamustafa” (Josifovo).

Table 3. Chemical characteristics of pomegranate juice.

Location	Repetition	pH	TSS (°Brix)	TA % (citric acid)	MI °Brix/TA	Salinity (%)	Total phenolic (mg/L)	Color of juice
Valandovo	I	3.31 ^a	13.8 ^{b,c}	0.56 ^a	2.49 ^c	12.7 ^{a,b}	978 ^b	Light pink
	II	3.30 ^a	13.5 ^{b,c}	0.55 ^a	2.44 ^c	12.5 ^{a,b}	981 ^b	
	III	3.32 ^a	12.1 ^d	0.51 ^{a,b}	2.20 ^d	12.8 ^{a,b}	980 ^b	
Josifovo	I	3.30 ^a	15.4 ^a	0.35 ^c	4.37 ^a	13.8 ^a	1014 ^a	Pink
	II	3.33 ^a	14.2 ^b	0.35 ^c	4.06 ^a	13.5 ^a	1015 ^a	
	III	3.32 ^a	14.0 ^b	0.35 ^c	3.94 ^a	13.5 ^a	1017 ^a	

Mean values followed by different lower-case letters in each column indicate significant difference among results at $P \leq 0.05$ by LSD test.

TSS - Total soluble solids, TA - titratable acids, MI - maturity index.

The results for the pH, total soluble solids (TSS), titratable acidity, maturity index, total polyphenolic of the pomegranate “Karamustafa” cultivar from the two different locations, are presented in Table 3. Significant differences ($P \leq 0.05$) were revealed among location for pH, total soluble solids, titratable acidity, maturity index and polyphenolic content.

As shown in Table 3, the highest total soluble solids content was in Josifovo, I (15.4 °Brix) and the lowest was in Valandovo, III (12.1 °Brix). The total acids content (equivalent to citric acid) is significantly higher in all fruits on the location Valandovo than Josifovo. Some of the Turkish cultivars range from 14.70°Brix to 17.90°Brix TSS and from 0.5% to 2.4% TA (Ozgen et al., 2008), Croatian autochthonous cultivars contain TSS from 13.0°Brix to 15.55°Brix TSS and from 0.44 to 1.64% total acids (Radunić et al., 2011), while Iranian cultivars from 12.85°Brix to 15.05°Brix TSS and 0.33% 2.44% total acids (Tehraniifar et al., 2010). Based on the total

acid content, the cultivars of pomegranate are divided into groups: sweet cultivars with <1 total acids, sweet-sour cultivars with 1-2% total acids and sour cultivars with > 2% total acids content (Bellini and Giordani, 1998).

The highest level of total phenolic compounds (1014-1017 mg/L gallic acid) was quantified in Josifovo. The level of the total phenolic content in Valandovo varied from 978 to 981 mg/L gallic acid. According to the results from our study, we can indicate that “Karamustafa” cultivar had higher value for total phenolic components than cultivar “Valencia” but significantly lower value than cultivars “Akko”, “Herskovitz” and “Wonderful” (Di Nunzio et al., 2013). However, the total phenolic content in pomegranate juice of “Karamustafa” is higher in Turkish pomegranate juice 500-916 mg/L gallic acid equivalent (Tehraniifar et al., 2010) but lower than in pomegranate juice found in Turkish markets (up to 2000 mg/L) (Tezkan et al., 2009).

CONCLUDING REMARKS

On the basis of the obtained results, we can conclude that the autochthonous cultivar "Karamustafa" belongs to the group of cultivars with large fruits, and according to the content of total acids in the group of sweet cultivars. A higher juice yield and higher content of the total phenols was obtained on the location Josifovo.

This is the first research for autochthonous cultivars pomegranate in the Republic of Macedonia. The research will continue in the direction of determining the potential of this and other pomegranate autochthonous cultivars for fresh consumption and for the production of juices.

REFERENCES

- Al-Said, F.A., Opara, L.U., Al-Yahyai, R.A. (2009). Physico-chemical and textural quality attributes of pomegranate cultivars (*Punica granatum* L.) grown in the Sultanate of Oman. *J. of Food Engineering*, 90, 129-134.
- Bellini, E., Giordani, E. (1998). Descriptor List for Pomegranate (*Punica granatum* L.). Project on "Minor Fruit Tree Species Conservation": RESGEN29. Horticulture Department, University of Florence, Italy.
- Cristofori, V., Caruso, D., Latini, G., Dell'agli, M., Cammilli, C., Rugini, E., Bignami, C., Muleo, R. (2011). Fruit Quality of Italian Pomegranate (*Punica granatum* L.) Autochthonous Varieties. *Eur. Food Res. Technol.*, 232, 397-403.
- Derün, K., Etü, S. (2001). Determination of pollen quality, quantity and effect of cross pollination on the fruit set and quality in the pomegranate. *Turk. J. Agric. For. Nom.*, 25, 169-173.
- Di Nunzio, M., Toselli, M., Verardo, V., Cabonia, M.F., Alessandra, Bordoni A. (2013): Counteraction of oxidative damage by pomegranate juice: influence of the cultivar. *Journal of the science of food and agriculture*, 93, 3565-3573.
- Drogoudi, P., Pantelidis, G., Manganaris, A. (2012). Morphological and physiological characteristics in pomegranate cultivars with different yields. Proceedings of II International symposium on the pomegranate, 67-69.
- Drogoudi, P.D., Tsipouridis, C. (2005). Physical and chemical characteristics of Pomegranates. *HortScience* 40 (5), 1200-1203.
- Fawole, O.A., Opara, U.L., Theron, K.I. (2011). Chemical and phytochemical properties and antioxidant activities of three pomegranate cultivars growing in South Africa. *Food bioprocess technology*, 5 (7), 2934-2940.
- Ferrara, G., Cavoski, I., Pacifico, A., Pacucci, C., Mondelli, D. (2012). A preliminary survey on pomegranate (*Punica granatum* L.) genotypes localized in Apulia region, Southeastern Italy. Proceedings of II International symposium on the pomegranate, 75-77.
- Font Quer P., (1979). Plantas medicinales. El Dioscórides renovado. Labor, S.A. 5ª ed. Barcelona. 1 033 pp.
- Gadže, J., Prlić, M., Bulić, M., Leko, M., Barbarić, M., Vego, D., Raguž, M. (2011). Physical and chemical characteristics and sensory evaluation of pomegranate fruit of cv. "Glavaš" (*Punica granatum* L.). *Pomologia Croatica*, 17, 87-97.
- Godwa, A.M., Osman, I.M., Mikhali, E.G. (2009). Comparative studies on six pomegranate cultivars under Beni-Suef governorate conditions. *J. Agric. Sci. mansoura Univ.*, 34(11), 10527-10541.
- Haddioui, A. (2012). La culture du grenadier (*Punica granatum* L.) au Maroc. Proceedings of II International symposium on the pomegranate, 79-81.
- Hassan, N.A., El-Halwagi, A.A., Sayed H.A. (2012). Phytochemicals, antioxidant and chemical properties of 32 pomegranate accessions growing in Egypt. *World applied Sciences Journal*, 16(8), 1065-1073.
- Karatas, S., Ozdogan, N. (2013). Efficiency of pomegranate seed oil. *International journal of electronics, mechanical and engineering*, 3(3), 591-597.
- Koppel, K., Chambers IV, E. (2010). Lexicon to describe appearance and flavor of pomegranate juice. *J. Sens. Stud.* 25, 819-837.
- Kotwal, G.J. (2007). Genetic diversity-independent neutralization of pandemic viruses (e.g.HIV), potentially pandemic (e.g. H5N1 strain of influenza) and carcinogenic

- (e.g. HBV and HCV) viruses and possible agents of bioterrorism (variola) by enveloped virus neutralizing compounds (EVNCs). *Vaccine*, 26, 3055-3058.
- Legua, P., Melgarejo, A. Haddioui, J.J., Martínez, R., Martínez, I., Hmid, H. Hanine, F., Hernández (2012). Characterization of six varieties of Moroccan pomegranate. Proceedings of II International symposium on the pomegranate, 83-86.
- Mahdavi, R., Nikniaz, Z., Rafrat, M., Jouyban, A. (2010). Determination and comparison of total polyphenol and vitamin C contents of natural fresh and commercial fruit juices. *Pakistan journal of nutrition*, 9(10), 968-972.
- Makkar, H.P.S., Bluemmel, M., Borowy, N.K., Becker, K. (1993). Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods, *J. Sci. Food Agric.* 61 (1993) 161-165.
- Mars, M., (1998). Pomegranate plant material: genetic resources and breeding (review). I Symposium Internacional sobre el granado. MV-0. Orihuela (Alicante).
- Martínez, J.J., Melgarejo, P., Legua, P., Martínez, R., Hernández, F. (2012). Diversity of pomegranate (*Punica granatum* L.) germplasm in Spain. Proceedings of II International symposium on the pomegranate, 53-56.
- Melgarejo, P., Martínez, J.J., Hernández, F., Legua, P., Melgarejo-Sánchez, P., Martínez Font, R. (2012). The pomegranate tree in the world: Its problems and uses. Proceedings of II International symposium on the pomegranate, 11-26.
- Ozgen, M., Durgac, C., Serce, S., Kaya, C. (2008). Chemical and antioxidant properties of pomegranate cultivars grown in the region of Turkey. *Food Chemistry*, 111, 703-706.
- Pekmezci, M., Erkan, M. (2004). Pomegranate. available on the Internet (<http://www.ba.ars.usda.gov/hb66/113pomegranate.pdf>).
- Radunić, M., Goreta Ban, S., Gadze, J., Jukić Špira, M., Tomasović, P., Maclean D. (2011). Pomological and chemical characteristics of pomegranate cultivars (*Punica granatum* L.) in the valley of the river Neretva. Proceedings. 46th Croatian and 6th International Symposium on Agriculture, Opatija, 1048-1051.
- Radunić, M., Jukić Špika, M., Goreta Ban, S., Gadže, J., Mac Lean, D. (2012). Chemical composition of pomegranate (*Punica granatum* L.) cultivars grown in Croatia. Proceedings of II International symposium on the pomegranate, 87-89.
- Reddy, M.K., Gupta, S.K., Jacob, M.R., Khan, S.I., Ferreira, D. (2007). Antioxidant, antimalarial and antimicrobial activities of tannin-rich fractions, elagitannins and phenolic acids from *Punica granatum* L.. *Planta Medica*, 73, 461-467.
- Saeedi, A.M., Mohammad, G., Samadi, G.R., Abdiani, S., Giordani, E. (2012). The Pomegranate national collection of Afghanistan. Proceedings of II International Symposium on the pomegranate, 57-60.
- Schwartz, E., Glazer, I., Bar-Ya'akov, I., Matityaha, I., Bar-Ilan, I., Holland, D., Amir, R. (2009). Changes in chemical constituents during the maturation and ripening of two commercially important pomegranate accessions. *Food Chemistry*, 115, 965-973.
- Tehraniifar, A., Zarei, M., Nemati, Z., Esfandiari, B., Vazifeshenas, M.R. (2010). Investigation of physico-chemical properties and antioxidant activity of twenty Iranian pomegranate (*Punica granatum* L.) cultivars. *Scientia Horticulturae*, 126, 180-185.
- Tezcan, F., Gultekin-Ozguven, M., Diken, T., Ozcelik, B., Bedia Erim, F. (2009). Antioxidant activity and total phenolic, organic acid and sugar content in commercial pomegranate juices. *Food Chemistry*, 115, 873-877.
- Westwood, N.H. (1982). *Fruticultura de Zonas Templadas*. Ediciones Mundi-Prensa. Madrid. 461.
- Zarei, M., Azizi, M., Bachiri-Sadr, Z. (2010). Studies of physico-chemical properties and bioactive compounds of six pomegranate cultivars grown in Iran. *Journal of food technology* 8 (3): 112-117.
- Zarei, M., Azizi, M., Bashir-Sadr Z. (2011). Evaluation of physicochemical characteristics of pomegranate (*Punica granatum* L.) fruit during ripening. *Fruits*, 6, 121-129.

ФИЗИЧКИ И ХЕМИСКИ КАРАКТЕРИСТИКИ НА КАЛИНКАТА (*PUNICA GRANATUM* L.), СОРТА КАРАМУСТАФА

Виолета Димовска¹, Фиданка Илиева¹, Сања Костадиновиќ¹, Љупчо Михајлов¹

¹Земјоделски факултет, Универзитет „Гоце Делчев“ - Штип, „Крсте Мисирков“, 10-А, 2000 Штип,
Република Македонија

violeta.dimovska@ugd.edu.mk; fidanka.ilieva@ugd.edu.mk; sanja.kostadinovik@ugd.edu.mk;
ljupco.mihajlov@ugd.edu.mk

Резиме

Калинките се добро адаптирани овошни видови во суви и полупустински услови и високоценети култури кои се одгледуваат во земјите на Медитеранот и во Азија. Најважните сорти калинка во Македонија се *бејнарија*, *лифанка* и *карамустафа*. Целта на оваа студија е да се евалуираат физичките и хемиските параметри на калинката сорта *карамустафа*, собрани од две различни локации во Република Македонија. Физичките параметри вклучуваат тежина на свежо овошје, процент на зрно (%), тежина од 100 бобинки (г), процент на сок (%). Сортата *карамустафа* (239.57- 304.79 г) припаѓа на група големи плодови. Сокот од родот се движи од 47,83% (Валандово) до 50,63% (Јосифово). Врз основа на содржината на вкупните киселини (ТС), сортата *карамустафа* ѝ припаѓа на групата со слатка сорта (ТА <1).

Врз основа на резултатите пронајдовме значителни разлики во тежината на овошјето, содржината на вкупните киселини и содржината на вкупните полифеноли во сокот.

Клучни зборови: *Punica granatum* L, овошје, сок, вкупни киселини, вкупно полифеноли