

*Martin Lukarevski and Dan Stefan Marinescu*

*A refinement of the Kooi's inequality, Mittenpunkt  
and applications*

# JOURNAL OF MATHEMATICAL INEQUALITIES

## AIMS AND SCOPE

Journal of Mathematical Inequalities (**JMI**, J. Math. Inequal.) presents carefully selected original research articles from all areas of pure and applied mathematics, provided they are concerned with mathematical inequalities. **JMI** will also periodically publish invited survey articles with interesting results treating the theory of inequalities, as well as relevant book reviews.

**JMI** is published quarterly, in March, June, September, and December.

## SUBMISSION OF MANUSCRIPTS

Authors are requested to submit their articles electronically as  $\text{T}_{\text{E}}\text{X}$  or  $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$  files, with enclosed Adobe Acrobat PDF format through the provided web interface on journal web page.

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the **JMI** regarding the article, makes any revisions, and reviews and releases the author proofs. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

## COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to the **JMI**. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal while under review for the **JMI**, and that its submission for publication has been approved by all of the authors.

## OPEN ACCESS

Journal of Mathematical Inequalities is published as open access journal. All interested readers are allowed to view, download, print, and redistribute any article without paying for subscription. Offprints of each article may be ordered from **JMI** prior to publication.

## PREPARATION OF MANUSCRIPTS

Manuscripts should be written in English, using prepared journal LaTeX style.

The first page should contain the article title, authors' names (complete first names and family names), complete affiliations (name of institution, address, city, state, and zip code), e-mail addresses, proposed running head (not more than 40 characters), a short abstract (not more than 150 words), a list of key words and phrases, and the AMS 2010 Mathematics Subject Classification primary (and secondary) codes. Avoid abbreviations, mathematical symbols and formulas, diagrams, and reference to the text of the article.

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

Bibliographic references should be listed alphabetically at the end of the article, each numbered by an Arabic number between square brackets. The following information should be provided for references to journals: names of authors, full title of the article, abbreviated name of the journal, volume, year of publication, and range of page numbers. For standard abbreviations of journal names, the authors should consult the latest Abbreviations of Names and Serials reviewed in Mathematical Reviews.

## SOURCE FILE, PROOFS

Upon acceptance of the article, the authors will be asked to send the related LaTeX source file to the Editorial Office, in accordance to the journal style. In order to accelerate the publication process, the AMS LaTeX package is strongly preferred. PDF proofs will be sent by e-mail to the corresponding author.

## FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

**JMI** is published by Publishing House **ELEMENT**, Zagreb, Croatia.

All correspondence and subscription orders should be addressed to the Editorial Office:

[www.ele-math.com](http://www.ele-math.com)  
e-mail: [jmi@ele-math.com](mailto:jmi@ele-math.com)

Journal of Mathematical Inequalities  
Editorial Office  
Menceticeva 2, 10000 Zagreb, Croatia  
Fax: +385 1 6008799

## A REFINEMENT OF THE KOOI'S INEQUALITY, MITTENPUNKT AND APPLICATIONS

MARTIN LUKAREVSKI AND DAN STEFAN MARINESCU

(Communicated by T. Burić)

*Abstract.* Kooi's inequality is important in the theory of triangle inequalities, giving bound on the semiperimeter in terms of circumradius and inradius. We give one refinement, then relate the inequality to the Mittenpunkt, and finally apply it to improve several known inequalities.

### 1. The inequality

Although not as widely known in the theory of triangle inequalities as the celebrated Gerretsen's inequality [7], the inequality of O. Kooi [1, 5.7] which has the same flavor - it gives bound from above on the square of the semiperimeter  $s$  of a triangle in terms of its circumradius  $R$  and inradius  $r$  - is also significant. The inequality in question is

$$s^2 \leq \frac{R(4R+r)^2}{2(2R-r)}. \quad (1)$$

It is stronger than the RHS of the Gerretsen's inequality

$$16Rr - 5r^2 \leq s^2 \leq 4R^2 + 4Rr + 3r^2, \quad (2)$$

since by Euler's inequality  $R \geq 2r$  follows that  $\frac{R(4R+r)^2}{2(2R-r)} \leq 4R^2 + 4Rr + 3r^2$ .

### 2. Sharpened Kooi's inequality

The reference [6] where Kooi's inequality appeared for the first time is not easily accessible and apart from it we were not able to find a proof in the literature. In this section we give a sharpened version of the inequality and a proof thereof, based on the RHS of an inequality that has been labeled "fundamental inequality of a triangle" [1, 5.10]

$$\begin{aligned} & 2R^2 + 10Rr - r^2 - 2(R-2r)\sqrt{R(R-2r)} \leq s^2 \\ & \leq 2R^2 + 10Rr - r^2 + 2(R-2r)\sqrt{R(R-2r)}. \end{aligned} \quad (3)$$

*Mathematics subject classification* (2010): 51M04, 51M16.

*Keywords and phrases:* Kooi's inequality, Mittenpunkt, barycentric coordinates, Hadwiger-Finsler inequality.

It is sometimes called Blundon’s inequality, despite its appearance in the literature being much older. The inequality is recorded in 1851 as a problem posed by C. Ramus [12], with a solution by Emile Rouché (1832-1910), who is best known for his theorem in complex analysis.

REMARK 1. Gerretsen’s inequality can also be derived from (3) or by calculating distances between triangle centers and showing that  $9GI^2 = s^2 - 16Rr + 5r^2$ ,  $HI^2 = 4R^2 + 4Rr + 3r^2 - s^2$ , where  $G$ ,  $I$  and  $H$  are the centroid, incenter and orthocenter of the triangle respectively. There is also a simple proof with an algebraic twist [7]. Sharp version of (3) is given in [14].

The next result is a refinement of the Kooi’s inequality (1).

THEOREM 1. *For the seiperimeter  $s$ , circumradius  $R$  and inradius  $r$  of a triangle it holds*

$$s^2 \leq \frac{R(4R+r)^2}{2(2R-r)} - \frac{r^2(R-2r)}{4R}.$$

*Proof.* We put  $t := r/R$ ,  $0 < t \leq 1/2$ ,  $K(R, r) := \frac{R(4R+r)^2}{2(2R-r)} - \frac{r^2(R-2r)}{4R}$  and  $f(R, r) := 2R^2 + 10Rr - r^2 + 2(R-2r)\sqrt{R(R-2r)}$ . Then  $f(R, r) \leq K(R, r)$  is equivalent to

$$2 + 10t - t^2 + 2(1-2t)\sqrt{1-2t} \leq \frac{(4+t)^2}{4-2t} - \frac{t^2(1-2t)}{4},$$

which can be rewritten as

$$2(1-2t)\sqrt{1-2t} \leq \frac{2(1-2t)(t^3 - 24t + 16)}{4(4-2t)}.$$

Squaring and expanding of this last inequality gives

$$t^3(t^3 - 48t + 160) \geq 0,$$

which is obviously true for  $0 < t \leq 1/2$ . Hence by the fundamental inequality  $K(R, r) \geq f(R, r) \geq s^2$  and the improvement of the Kooi’s inequality (1) is proven.  $\square$

To understand thoroughly Kooi’s inequality, it is desirable to have a geometric proof in which triangle centers are involved, a proof similar in spirit to the above mentioned proof of Gerretsen’s inequalities. Such a proof we give in the next section.

### 3. Mittenpunkt and barycentric coordinates

For a point  $P = (x : y : z)$  with normalized barycentric coordinates, its squared distance  $OP^2$  from the circumcenter  $O$  of the triangle  $ABC$  with sides  $a$ ,  $b$ ,  $c$  and circumradius  $R$  is given by [13]

$$OP^2 = R^2 - a^2yz - b^2zx - c^2xy. \tag{4}$$

In the geometric proof of Gerretsen's inequalities all the classical triangle centers are involved. We consider now one lesser-known point, the Mittenpunkt  $M$ , which is still a household center for all the triangle geometers. That is the point which is the intersection of the lines passing through the centers of the excircles and the corresponding midpoints of the sides of the triangle (hence its name, given by the German mathematician Nagel in 1836). For more properties of  $M$  see [5], where the point is denoted by  $X_9$ .

**THEOREM 2.** *The distance from the circumcenter  $O$  to the Mittenpunkt  $M$  of the triangle is*

$$OM^2 = R^2 - \frac{2Rs^2(2R - r)}{(4R + r)^2}.$$

*Proof.* The Mittenpunkt  $M$  has normalized barycentric coordinates [16],  $M = \left(\frac{a(s-a)}{D} : \frac{b(s-b)}{D} : \frac{c(s-c)}{D}\right)$ , where  $D = \sum a(s-a) = 2r(4R+r)$ , by the well-known relation  $a^2 + b^2 + c^2 = 2(s^2 - 4Rr - r^2)$ , [11, p.52]. We use  $\sum a(s-b)(s-c) = 2rs(2R-r)$ , which follows from  $ab + bc + ca = s^2 + 4Rr + r^2$  and  $abc = 4Rrs$ . Hence by (4), we get

$$\begin{aligned} OM^2 &= R^2 - \frac{abc}{D^2} \sum a(s-b)(s-c) \\ &= R^2 - \frac{2Rs^2(2R - r)}{(4R + r)^2}. \quad \square \end{aligned}$$

**COROLLARY 1.** *Kooi's inequality (1), which is equivalent to  $OM^2 \geq 0$ .*

### 4. Equivalent Forms of Kooi's inequality

In this section we will give few interesting equivalent forms of the Kooi's inequality. By the well-known identities

$$\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \frac{r}{4R}$$

and

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} = \frac{(4R+r)^2}{s^2} - 2,$$

Kooi's inequality (1) rewritten as  $\frac{(4R+r)^2}{s^2} - 2 \geq 2 - \frac{2r}{R}$ , can be trigonometrised, giving the Garfunkel-Bankoff inequality [4]

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} \geq 2 - 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}.$$

Another equivalent form of the inequality is

$$4R + r \geq s \sqrt{3 + \frac{R - 2r}{R}}, \tag{5}$$

which will be used in the last section. Using the identity

$$\tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} = \frac{4R+r}{s},$$

we obtain the sharpening

$$\tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} \geq \sqrt{3 + \frac{R-2r}{R}}$$

of the well-known inequality  $\tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} \geq \sqrt{3}$ .

### 5. Improved Hadwiger-Finsler, Gordon's, Curry's and Leunberger's Inequality

As an application, we show that the celebrated Hadwiger-Finsler inequality [3]

$$4\sqrt{3}A + Q \leq a^2 + b^2 + c^2 \leq 4\sqrt{3}A + 3Q, \quad (6)$$

where  $Q := (a-b)^2 + (b-c)^2 + (c-a)^2$  and  $A$  is the area of the triangle, can be improved using the Kooi's inequality to [8]

$$a^2 + b^2 + c^2 \geq 4\sqrt{3 + \frac{R-2r}{R}}A + Q. \quad (7)$$

Indeed, using the well-known identity  $ab + bc + ca = s^2 + 4Rr + r^2$  and (5), we get

$$\begin{aligned} a^2 + b^2 + c^2 - [(a-b)^2 + (b-c)^2 + (c-a)^2] &= 4(ab + bc + ca) - (a+b+c)^2 \\ &= 4r(4R+r) \geq 4rs\sqrt{3 + \frac{R-2r}{R}}, \end{aligned}$$

which proves (7).

REMARK 2. For a proof of Hadwiger-Finsler inequality (6) based on Gerretsen's inequality (2) see [7]. The LHS of (6) can also be derived [9] from the seemingly weaker Weitzenbock inequality  $a^2 + b^2 + c^2 \geq 4\sqrt{3}A$ .

REMARK 3. From (7) directly follows

$$ab + bc + ca \geq 4\sqrt{3 + \frac{R-2r}{R}}A,$$

which is an improvement of the Gordon's inequality [1, 4.5]  $ab + bc + ca \geq 4\sqrt{3}A$ .

Kooi's inequality (5) together with Euler's inequality  $R \geq 2r$  gives

$$9R \geq 2(4R+r) \geq 2s\sqrt{3 + \frac{R-2r}{R}}.$$

Now  $\frac{9R}{2s} \geq \sqrt{3 + \frac{R-2r}{R}}$  is equivalent to

$$\frac{9abc}{a+b+c} \geq 4A\sqrt{3 + \frac{R-2r}{R}},$$

which improves Curry's inequality [2]  $\frac{9abc}{a+b+c} \geq 4A\sqrt{3}$ .  
 Leuenberger's inequality [1, 5.22] is

$$\frac{\sqrt{3}}{R} \leq \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \leq \frac{\sqrt{3}}{2r}.$$

Its LHS is equivalent to Gordon's inequality, so it can be improved to

$$\frac{\sqrt{3 + \frac{R-2r}{R}}}{R} \leq \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

As a next application, we give a simpler proof of a result from [15] which seemingly generalizes (7) for  $p = 1$  but is actually equivalent to it.

**THEOREM 3.** *Let  $p \geq 1$ . Then*

$$a^{2p} + b^{2p} + c^{2p} \geq 3^{1-p} \left[ (4A)^p \left( 3 + \frac{R-2r}{R} \right)^{p/2} + (a-b)^{2p} + (b-c)^{2p} + (c-a)^{2p} \right].$$

*Proof.* By Jensen's inequality applied to the convex function  $f(x) = x^p$  we have  $3^{p-1}(a^{2p} + b^{2p} + c^{2p}) \geq (a^2 + b^2 + c^2)^p$ . The superadditivity of  $f(x) = x^p$  and (7) yield

$$\begin{aligned} (a^2 + b^2 + c^2)^p &\geq (a-b)^{2p} + (b-c)^{2p} + (c-a)^{2p} + (a^2 + b^2 + c^2 - Q)^p \\ &\geq (a-b)^{2p} + (b-c)^{2p} + (c-a)^{2p} + (4A)^p \left( 3 + \frac{R-2r}{R} \right)^{p/2}. \end{aligned}$$

Putting together the two inequalities finishes the proof.  $\square$

We conclude this note by asking for a geometric proof of the sharpening of Kooi's inequality.

REFERENCES

- [1] O. BOTTEMA, R. Z. DJORDJEVIC, R. R. JANIC, D. S. MITRINOVIC, P. M. VASIC, *Geometric inequalities*, Groningen, Wolters-Noordhoff, 1969.
- [2] T. R. CURRY, *The American Mathematical Monthly*, **73** (1966), no. 2, Problem E 1861.
- [3] P. FINSLER, H. HADWIGER, *Einige Relationen im Dreieck*, *Commentarii Mathematici Helvetici*, **10** (1937), no.1, 316–326.
- [4] Problem 825 (proposed by J. Garfunkel, solution by L. Bankoff), *Crux Math.* **9** (1983), 79 and **10** (1984), 168.
- [5] C. KIMBERLING, *Central Points and Central Lines in the Plane of a Triangle*, *Math. Mag.* **67** (1994), no. 3, 163–187.

- [6] O. KOOI, *Inequalities for the triangle*, Simon Stevin, **32** (1958), 97–101.
- [7] M. LUKAREVSKI, *An alternate proof of Gerretsen's inequalities*, *Elem. Math.* **72**, No. 1 (2017), 2–8.
- [8] M. LUKAREVSKI, *The American Mathematical Monthly*, **123**, (2016), no. 9, Problem 11938.
- [9] M. LUKAREVSKI, *The excentral triangle and a curious applications to inequalities*, *The Math. Gazette* **102**, No. 555, (November 2018), 531–533.
- [10] D. S. MARINESCU, M. MONEA, M. OPINCARIU, M. STROE, *Note on Hadwiger-Finsler's inequalities*, *J. Math. Inequal.* **6**, No. 1, 57–64 (2012).
- [11] D. S. MITRINOVIĆ, J. PEČARIĆ, V. VOLENEC, *Recent Advances in Geometric Inequalities*, Kluwer Academic Publishers, Dordrecht (1989).
- [12] RAMUS-ROUCHÉ, *Question 233*, *Nouv. Ann. Math.* 10 (1851), 353.
- [13] V. VOLENEC, *Metrical relations in barycentric coordinates*, *Math. Commun.* 8, No. 1, 55–68 (2003).
- [14] S. WU, *A sharpened version of the fundamental triangle inequality*, *Math. Inequal. Appl.* 11 (2008), no. 3, 477–482.
- [15] S. WU, L. DEBNATH, *Parametrized Garfunkel-Bankoff inequality and improved Finsler-Hadwiger inequality*, *Appl. Math. Lett.* **25**, No. 3, 331–336 (2010).
- [16] P. YIU, *The uses of homogeneous barycentric coordinates in plane Euclidean geometry*, *Internat. J. Math. Ed. Sci. Tech.* 31 (2000), no. 4, 569–578.

(Received March 14, 2017)

*Martin Lukarevski*  
*Department of Mathematics and Statistics*  
*University "Goce Delcev"*  
*Stip, Macedonia*  
*e-mail: martin.lukarevski@ugd.edu.mk*

*Dan Stefan Marinescu*  
*Colegiul National "Iancu De Hunedoara"*  
*Hunedoara, Romania*  
*e-mail: marinescuds@gmail.com*