

Representation of diabetes in children and adolescents in Eastern Macedonia

Panova G., Ivanoska B., Panov N., Velichkova N., Kamcev N., Kamceva G., Nikolova L.

Faculty of medical science, University Goce Delcev – Stip, R.Macedonia

Background

Two more children under five years Europe will be infected by type 1 diabetes from 2005 to 2020 *. In diabetes type 1 beta stations gradual collapse of the thyroid gland that produce insulin to complete loss of insulin.

- This indicates research published by the on - line UK shop "D Lancet".

Methods

Type 1 diabetes often appears in contrast to type 2 and a rate of seven to 10 percent in all cases of diabetes. In children, they scale conflict in a larger number drzhavi.Nauchnitsite found that the total number of sufferers grows by 3.9 percent per year. The study does not explain the trend of growth, but scientists believe that too an increasing number of sufferers can be prescribed to genetic factors. They focus on the role of obesity and more caesarean section.Number of sick children from type 1 diabetes in this age postijano and grew rapidly primarily as a result of genetic predisposition. Treatment with insulin should regulate parents.

Results

In the period of 2009. In my municipality were registered suffering from diabetes and this: From 3-6 years. 35 children infected of whom 15 women and 20 children 6-11, the male children..Od. Infected 47 children of whom 35 women and 12 children 12-18, the male children.Od. 50 children infected of whom 27 women and 23 male children. Period of 2010. In my municipality were registered suffering from diabetes and this: From 3-6 years. Infected 42 children of whom 20 women and 22 children guys. 6-11 god.Ill patients 50 children, of which 35 female and 15 male children 12-18 .Sufferers 58 children, 38 women children and 20 boys.

Conclusion

The display may be noted that the number of sick children and adolescents with diabetes each year have experienced steady growth.
