

GOCE DELCEV UNIVERSITY - STIP
FACULTY OF AGRICULTURE



JOURNAL OF AGRICULTURE AND PLANT SCIENCES

YEAR 2019

VOLUME 17, Number 2

EDITOR IN CHIEF

Editor in Chief

Liljana Koleva Gudeva, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, liljana.gudeva@ugd.edu.mk

Editors

Ljupco Mihajlov, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, ljupco.mihajlov@ugd.edu.mk

Fidanka Trajkova, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, fidanka.trajkova@ugd.edu.mk

Administrator

Biljana Atanasova, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, biljana.atanasova@ugd.edu.mk

Technical Editing

Slave Dimitrov, Publishing Department, Goce Delcev University, Stip,
Republic of North Macedonia, slave.dimitrov@ugd.edu.mk

Language Editors

Biljana Ivanova, MA, Senior lecturer, Faculty of Philology, Goce Delcev University, Stip,
Republic of North Macedonia Macedonia, biljana.petkovska@ugd.edu.mk – English language Editor

Danica Gavrilovska Atanasovska, Goce Delcev University, Stip, Republic of North Macedonia
Macedonia, danica.gavrilovska@ugd.edu.mk – Macedonian language Editor

Editorial Office

Faculty of Agriculture, Goce Delcev University, Stip, Krste Misirkov Str., No.10-A P.O. Box 201, 2000
Stip, Republic of North Macedonia
japs@ugd.edu.mk
<http://js.ugd.edu.mk/index.php/YFA>

**GOCE DELCEV UNIVERSITY - STIP, REPUBLIC OF NORTH MACEDONIA
FACULTY OF AGRICULTURE**

UDC 63(058)
ISSN 2545-4447 print
ISSN 2545-4455 on line



Journal of Agriculture and Plant Sciences, JAPS, Vol 17, No. 2

Successor of the Yearbook of Faculty of Agriculture of GDU, Vol 14

YEAR 2019

VOLUME XVII, Number 2

EDITORIAL BOARD

Aco Kuzelov,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, aco.kuzelov@ugd.edu.mk

Danijela Raičević,

Biotechnical Faculty, University of Montenegro, Mihaila Lalica bb., Podgorica,
Montenegro, nelar@mail.com

Dragan Skorić,

Serbian Academy of Sciences and Arts, Knez Mihajlova 35, 11000 Belgrade,
Serbia, draganskoric@sbb.rs

Dragomir Vlcev,

Institute of Agriculture – Karnobat, Bulgaria, vulchevd@abv.bg

Hatice Gülen,

Istanbul Bilgi University, Faculty of Engineering and Natural Sciences,
Department of Genetics and Bioengineering, Istanbul, Turkey,
hatice.gulen@bilgi.edu.tr

Jovica Vasin,

Institute of Field and Vegetable Crops, Novi Sad, Serbia, jovica.vasin@ifvcns.ns.ac.rs

Kiril Bahcevandziev,

Coimbra Agricultural School, 3045-601 Coimbra, Portugal, kiril@esac.pt

Klemen Lisjak,

Agricultural Institute of Slovenia, Hacquetova ulica 17, Ljubljana, Slovenia, Klemen.Lisjak@kis.si

Marijan Bubola,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, marijan@iptpo.hr

Maryna Mardar,

Odessa National Academy of Food Technologies, Odessa, 65039, Kanatnaya Str., Ukraine,
marinamardar2003@gmail.com

Rubin Gulaboski,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, rubin.gulaboski@ugd.edu.mk

Sanja Radeka,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, sanja@iptpo.hr

Sasa Mitrev,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, sasa.mitrev@ugd.edu.mk

Shuhe Wei,

Institute of Applied Ecology, Chinese Academy of Sciences, China, shuhewei@iae.ac.cn
Velichka Nikolova Rodeva,
Maritsa Vegetable Crops Research Institute, 32 Brezovsko shosse St., Plovdiv 4003, Bulgaria,
velirod@yahoo.com

Violeta Dimovska,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, violeta.dimovska@ugd.edu.mk

Wolfram Schnäckel,

Anhalt University of Applied Sciences, Bernburger Straße 55, 06366 Köthen, Germany,
Wolfram.Schnaeckel@hs-anhalt.de

CONTENT

Darko Andronikov, Aco Kuzelov, Julija Sazdova, Kiro Mojsov, Aco Janevski, Sonja Jordeva, Saška Golomeova Longurova QUANTITATIVE CHARACTERISTICS OF RABBIT HYBRIDS	9
Katerina Bojkovska, Nikolche Jankulovski, Goran Mihajlovski, Gordana Dimitrovska, Elena Joshevska CONSUMER PREFERENCES AS DETERMINANT OF THE SUCCESSFUL QUALITY MANAGEMENT OF DAIRY PRODUCTS	15
Daniela Dimovska, Ljupco Mihajlov, Fidanka Trajkova, Liljana Koleva Gudeva, Igor Iljovski THE INFLUENCE OF ORGANIC FERTILIZERS ON THE GROWTH AND YIELD OF BROCCOLI (<i>Brassica oleracea</i> L. var. <i>italica</i>)	25
Elena Joshevska, Aco Kuzelov, Gordana Dimitrovska, Katerina Bojkovska IMPROVEMENT OF THE PRODUCTION TECHNOLOGY OF MEAT PRODUCTS WITH THE ADDITION OF FUNCTIONAL COMPONENTS	31
Karolina Kocoska, Ilija Risteski THE SIZE AND NUMBER OF MIDDLE BELT LEAVES IN SOME VARIETIES OF WILD TOBACCO	39
Liljana Koleva Gudeva, Valentina Butleska Gjoroska, Fidanka Trajkova, Ljupco Mihajlov ACTIVITY OF ENZYME CATALASE IN ALFALFA (<i>Medicago sativa</i> L.) AS AN INDICATOR FOR ABIOTIC STRESS	45
Daniela Nikolovska Nedelkoska, Tatjana Kalevska, Natalija Atanasova-Pancevska, Mitko Karadelev, Zora Uzunoska, Dzoko Kungulovski EVALUATION OF BACTERICIDAL ACTIVITY OF SELECTED WILD MACROFUNGI EXTRACTS AGAINST <i>Escherichia coli</i>	53
Silvana Pashovska, Trajko Miceski PERSPECTIVES AND POSSIBILITIES FOR DEVELOPMENT OF TOBACCO PRODUCTION IN THE REPUBLIC OF NORTH MACEDONIA	59

INTRODUCTION

In the past twelve years the educational, research and applicative activities of the Faculty of Agriculture – Stip, Goce Delcev University – Stip, contributed to the development of agriculture sector in the country and broader region.

The Faculty of Agriculture organized the 2nd International Meeting Agriscience & Practice (ASP 2019), giving an opportunity to the participants for presentation and discussion of original scientific and practical results in different fields of agriculture.

The 2nd International Meeting Agriscience & Practice (ASP 2019), held on 12 April 2019 at the Faculty of Agriculture - Stip, was organized with intention to bring together all agricultural stakeholders for sharing their knowledge, experience and obstacles. One of the main aims was to link research and field work in agricultural sector in the country and broader, giving it an international dimension. All oral presentations as well as poster presentations at ASP 2019 were organized in several scientific sessions:

- Agricultural economics,
- Plant biotechnology,
- Plant production,
- Plant protection,
- Quality control and food safety,
- Soil science and hydrology,
- Viticulture, enology and fruit production.

The main goal of the Meeting was linking and promoting scientific achievements and practical knowledge, presented in different thematic areas, which were achieved in the Republic of North Macedonia and wider in the region.

Journal of Agriculture and Plant Sciences Vol. 17, No. 2 mostly contains the presented papers from the 2nd International Meeting Agriscience & Practice (ASP 2019).

Editorial Board,

December, 2019

Editor in chief,

Prof. Liljana Koleva Gudeva, PhD



QUANTITATIVE CHARACTERISTICS OF RABBIT HYBRIDS

**Darko Andronikov^{1*}, Aco Kuzelov², Julijana Sazdova³, Kiro Mojsov¹, Aco Janevski¹,
Sonja Jordeva¹, Saška Golomeova Longurova¹**

¹*Goce Delcev University, Faculty of Technology, Stip, Krste Misirkov 10-A, 2000 Stip,
Republic of North Macedonia*

²*Goce Delcev University, Faculty of Agriculture, Stip, Krste Misirkov 10-A, 2000 Stip,*

³*SOU Kole Nehtenin, Stip, Republic of North Macedonia*

*Corresponding e-mail: darko.andronikov@ugd.edu.mk

Abstract

Flemish giant rabbit and New Zealand white rabbits (group ON), were used as a material for the study. Rabbits were fed ad libitum with commercial pelleted feed, with the addition of small amounts of green feed and hay. The experiment included 6 rabbits, 3 males and 3 females. Previously defined live weight of 1800 to 2500 g was achieved within 77 days and after that they were slaughtered. Study results presented in this paper show fattening and slaughter properties of Flemish giant rabbit and New Zealand white rabbit. The achieved average weight of male rabbits was 2792 g and meat-to-bone ratio was 47.92%, 2597 g and 48.61% respectively in female rabbits. Average participation of the back part of the carcass in male/female rabbits was 28.02% and 27.34%, dorsal part was 20.93% and 22.65%, front of the carcass was 25.26% and 25.39%.

Key words: rabbit, hybrids, live weight

INTRODUCTION

The food of animal origin still constitutes one of the most important components of human nutrition, because it is a source of numerous essential nutrients that a person needs for growth, development and normal functioning.

Rabbit meat products can be evaluated according to carcass quality and to meat quality as for other livestock. Carcass quality has to satisfy economic objectives, such as saleable meat yield and attractiveness to consumer (Gjurič, 1985; Dalle Zotte, 2002). The dressing percentage is a very important economic variable in the rabbit market. Some combinations of measurements such as retail cut weights or length measurements are necessary to predict lean percentage in the carcass (Yalçın et al., 2006).

The meat of rabbit belongs to the group of meat rich with protein, significantly low in fat and cholesterol and it contains essential nutrients: essential amino and fatty acids, calcium, copper, iron, vitamins from the B complex, low energy value, and less connective

tissue (Koch & Pavčič, 2000; Wood et al. 2003; Dal Bosko et al. 2001; Pogány Simonová, et al. 2010; Chrenek, et al. 2012; Para et al. 2015). The meat has a mild taste and is easily digestible therefore it is known as a dietary food. It is recommended for children in growth, for a diet of patients with arteriosclerosis and enlarged lipids, as well as for patients with stomach illness and elderly people (Grün, 2002; Lah, 2006).

World rabbit meat production amounted to 1.56 million tonnes in 2014. The leading world producer of rabbit meat is China with 762,627 t year, while, in Europe, the main producer is Italy (268,980 t), followed by Spain (63,790 t), France (53,292 t), Czech Republic (38,602 t) and Germany (34,253 t year). Unfortunately, for most Balkan countries we have not managed to find data, except for Greece (6,799 t), Bulgaria (6,629 t) and Romania (143 t year) (FAOSTAT, 2014; Belichovska et al. 2017).

The purpose of cultivating the hybrids is to obtain the meat, that is, they are kept solely for slaughter. When creating animals - hybrids, two (or more) selected lines are crossed along by

either the father or the mother. During crossing two lines hybrids are created, which in turn will give the best features from both lines. The mother line brings fertile properties (increased number of offspring), while the father gives racial qualities (ability to grow rapidly, good food utilization, good slaughter quality and meatiness) (Grün, 2002).

The influences in the nutrition over the parts of the trunk, the fattening and slaughtering properties of rabbits have been studied by many authors (Panić & Petrović, 1989; Ozimba & Lukefahr 1991; Dalle Zotte et al. 1996; Skandro

et al. 2004; Metzger et al. 2006; Omojola 2007; Baiomy & Hassanien 2011; Kostovska 2012). Chemical composition of rabbit meat has been especially investigated (Dalle Zotte et al. 1996; Hernández et al. 1998; Gondret et al. 1998; Nizza & Moniello 2000; Metzger et al. 2003; Wood et al. 2003; Pascual et al. 2004; Polak et al. 2006; Ali, 2007; Nistor et al. 2013).

The purpose of this paper is to present the fattening and slaughtering properties of the meat of crossed hybrids based upon our results and data.

MATERIAL AND METHODS

Six rabbit's hybrids (3 male and 3 female), that is, crossed units from the so-called Belgium oriash and New Zealand's white rabbits (ON) fed ad libitum with industrially pelleted food (containing alfalfa, barley, corn, wheat, soy, granules sunflower premixes, salt, vitamins and minerals) were the research material for our study.

The rabbits reached the defined weight from 1800 to 2500 g in 77 days kept in separate wire cages. 24 hours before slaughter, their feeding was stopped. Slaughtering and primary processing of rabbits was performed in the usual way. The slaughter was done after the

veterinary examination and looting. 24 post mortem cool carcasses were cut in basic peaces and measured on an electronic scale, there in each organ individually, and then cut to the front, the middle and the back part (Bivolarski et al., 2011).

Meat-to-bone ratio (slaughter weight expressed in percentage) was calculated in compliance with the recommended processing procedure (Gjurič, 1985; Caklovica et al. 1986; Omrčen, 1995; Skandro et al. 2004; Ali, 2007).

Statistical analyses were made by using the statistical software SPSS ver. 21 (SPSS Inc, Chicago, IL, 2012).

RESULTS AND DISCUSSION

The middle values of live weight of male/female rabbits are shown following the cycle of 77 days of their age. The table shows that the live weight of male rabbits on average was 2792 grams while at female rabbits it was 2597 grams. The results that we found in relation to the live weight of male vaccines are in line with the results obtained by Kostovska (2012), while in relation to female units the results obtained by Kostovska (2012) compared to our results are higher for 191 grams.

The values we found in relation to the live mass compared to the values of live mass found by Ali (2007) and Skandro et al. (2008) are higher. Within our study the male rabbits achieved a higher body mass compared to females (for 213 grams), while in the studies by Skandro et al. (2008) the female rabbits achieved a higher body mass than the males for 10 grams.

Table 1. Meat-to-bone ratio (%) - slaughter and live weight (g) of male and female rabbits.

	ON Male	ON Female
Live weight (g)	2792±140	2597±90
Mass slaughter (g)	1338±90	1280±50
Meat-to-bone ratio (%)	47.92	49.29

\bar{x} – mean, Sd – standard deviation; number of pieces = 3

Meat-to-bone ratio (%) of male rabbits is 47.92% while at female rabbits it is 49.29%. The results that we obtained in relation to the meat-to-bone ratio (%) are in line with the allegations of other researchers (Kovačević and Rašeta,

1983; Urošević et al. 1986; Skandro et al. 2008). Tafro et al. (1989); Skandro et al. (2004), Ali (2007) and Kostovska (2012) referred to meat-to-bone ratio (%) values for rabbits between 40 and 53%.




Figure 1. Meat-to-bone ratio (%) of male and female rabbits.

The average level of blood in the total body mass is 3.34%; at male rabbits it is 3.22% and at female rabbits it is 3.46%. A bit lower results in relation to the results obtained by us in terms of the prevalence of blood in the live weight of rabbits is stated by Kostovska (2012). The same author found blood levels in relation to the live weight of male rabbits of 2.95% and at female rabbits of 2.97%.

Skandro et al. (2008) found blood levels in relation to the live weight of rabbits, which is 7.20%, that is 7.70% at male rabbits and 6.70% at female rabbits. The results they received were greater than the results obtained by us probably because they also calculated the crack of cutting.




Figure 2. Participation of major meat categories in carcass composition of male and female rabbits.

The head with the total body weight at male rabbits is 4.83% and at female rabbits it is 4.74%. The results that we obtained are lower compared to the results obtained by Skandro et al. (2008); Kostovska (2012). Compared to our results, Kostovska (2012) found a slightly higher participation of the head in the body weight of the fuselage of 5.71% at male rabbits and 5.34% at female blacks, probably as a result of researching a

larger number of rabbits (n = 18; 9 male / 9 female).

Skandro et al. (2008) determined the participation of the head in the final body mass of the body 8.68% (at male rabbits 8.71%, at female rabbits 8.65%). Our values were lower than the results obtained by Skandro et al. (2008) and Kostovska (2012) probably due to the larger number of male and female individuals (n = 24 rabbits; 12 male / 12 female).




Figure 3. Participation of fat and major meat categories in carcass composition of male and female rabbits

The average participation of internal organs in the total body mass of male and female rabbits is 25.79%. Our results were lower than the results obtained by Ali (2007) who found a share of internal organs of 30.72%, and those were greater

than the values obtained by Skandro et al. (2008) and Kostovska (2012), who established a share of internal organs of 23.84% at male rabbits and 23.48% at female rabbits, that is, 24.39% at male rabbits and 24.83% at female rabbits.

CONCLUSIONS

Based on results of investigation of quantitative characteristics of meat of Flemish giant rabbit and New Zealand white rabbit, it may be concluded as follows:

- Average carcass dressing percentage, without head, was higher in female rabbits,

- Participation of hind and back parts, considered as the qualitative parts of meat in dressed and cooled carcass, was higher in male rabbits.

REFERENCES

- Ali T. (2007): Utjecaj ishrane i načina uzgoja na kvalitativno – kvantitativna svojstva novozelandskog bijelog kuniča, Magistarski rad, Sarajevo.
- Baiomy, A.A., & Hassanien H.H.M. (2011). Effect of breed and sex on carcass characteristics and meat chemical composition of New Zealand white and Californian rabbits under upper Egyptian environment. *Egypt Poultry Science*, 31, 275–284.
- Bivolarski, B.L., Vachkova E.G., & Ribarski S.S. (2011). Effect of weaning age upon the slaughter and physicochemical traits of rabbit meat. *Veterinarski Arhiv*, 81, 499–511.
- Belichovska, D., Belichovska, K., Pejkovski, Z., & Uzunoska, Z. (2017). Effect of genotype on certain physico-chemical characteristics of rabbit meat. *Scientific Journal "Meat Technology"*, 58(1), 10-15.
- Caklovica, F., Milanović, A., & Tafro, N. (1986). Prilog istraživanjima klaoničnih vrijednosti kuniča, X Savetovanje o dijagnostici, profilaksi i terapiji u savremenoj stočarskoj proizvodnji. *Veterinaria, Primošten, Croatia. Proceedings*: 35(1), 91 – 95.
- Chrenek, P., Makarevich, A., Kozelová, D., & Rafay J. (2012). Meat quality of transgenic rabbit. *Slovak Journal of Animal Science*, 45, 60–62.
- Dal Bosko A., Castellini C., & Bernardini M. (2001). Nutrition-al quality of rabbit meat as affected by cooking procedure and dietary vitamin E. *Journal of Food Science*, 66, 1047–1051.
- Dalle Zotte, A. (2002): Perception of rabbit meat quality and major factors influencing the rabbit carcass and meat quality. *Livestock Production Science*, 75, 11-32.
- Dalle Zotte, A., Ouhayoun, J., Parigi Bini, R., & Xiccato G. (1996). Effect of age, diet and sex on muscle energy metabolism and on related physicochemical traits in the rabbit. *Meat Science*, 43, 15-24.

- FAOSTAT (2014). Food and Agriculture Organization statistical database. Retrieved from: <http://www.fao.org/faostat/en/#data/QL>. Access: Production / Livestock Primary, 2014.
- Gjurič, A. (1985): Kuničarstvo. Popularno – stručna biblioteka, Nakladni zavod znanja, Zagreb, Croatia.
- Gondret, F., Juin, H., Mourot, J., & Bonneau M. (1998). Effect of age at slaughter on chemical traits and sensory quality of longissimus lumborum muscle in the rabbit. *Meat Science* 48, 181-187.
- Grün, P. (2002, April). Reja kuncev. Ljubljana, Kmečki glas, pp. 106-126.
- Hernández, P., Pla, M., & Blasco A. (1998). Carcass characteristics and meat quality of rabbit lines selected for different objectives: II. Relationships between meat characteristics. *Livestock Production Science*. 54, 125-131.
- Koch, V., & Pavčič M. (2000). Kriteriji za ocenjevanje mesa in mesnin z vidika varovalnega živila. V: Meso in mesnine za kakovostno prehrano. 2. Posvet o vlogi in pomenu mesa v normalni-zdravi in dietni prehrani, Portorož, Slovenija. Proceedings: 10-14.
- Kostovska A. (2012): Кланични карактеристики на зајците од калифорниската раса. Магистерски труд, Битола.
- Kovačević, M., & Rašeta, D. (1983). Praktično kuničarstvo. Beograd, Republic of Serbia.
- Lah L. (2006): Vsebnost maščob in maščobnokislinska sestava mesa kuncev različnih pasemskih linij, spolov in starosti. Diplomsko delo, Ljubljana.
- Metzger, Sz., Kustos, K., Szendro, Zs., Szabo, A., Eiben, Cs., & Nagy I. (2003). The effect of housing system on carcass traits and meat quality of rabbit. *World Rabbit Science*, 11, 1–11.
- Metzger, Sz., Odermatt M., Szendro Zs., Mohaupt M., Rom-vári R., Makai A., Biró-Németh E., Radnai I., & Sipos L. (2006). Comparison of carcass traits and meat quality of Hyplus hybrid, purebred Pannon White rabbits and their crossbreds. *Archiv Tierzucht*, 49, 389–399.
- Nistor, E., Bampidis V.A., Păcală N., Pentea M., Tozer J., & Prundeanu H. (2013). Nutrient content of rabbit meat as compared to chicken, beef and pork meat. *Journal of Animal Production Advances*, 3, 172–176.
- Nizza, A. & Moniello G. (2000). Meat quality and caecal content characteristics of rabbit according to dietary content and botanical origin of starch. *World Rabbit Science*, 8, 3–9.
- Omojola A. B. 2007. Effect of delayed bleeding on carcass and eating qualities of rabbit meat. *Pakistan Journal of Nutrition*, 6, 438–442.
- Omrčen, S. (1995). Kuničarstvo, Nakladni zavod Globus, Zagreb, Croatia.
- Ozimba, C.E., & Lukefahr S. D. (1991). Comparison of rabbit breed types for postweaning litter growth, feed efficiency, and survival performance traits. *Journal of Animal Science*, 69, 3494–3500.
- Panić, M., & Petrović M. (1989). Fenotipska povezanost osnovnih delova trupa i tkiva kunića brojlera. IX Jugoslovensko savjetovanje, kvalitet i standardizacija mesa stoke za klanje, peradi, divljači i riba. Donji Milanovac, Serbia. Proceedings: 427-432.
- Para, A.P., Ganguly, S., Wakchaure, R., Sharma, R., Mahajan T., & Praveen K. P. (2015). Rabbit meat has the potential of being a possible alternative to other meats as a protein source: A Brief Review. *International Journal of Pharmacy & Biomedical Research*, 2, 17–19.
- Pascual, M., Aliaga, S., & Pla M. (2004). Effect of selection for growth rate on carcasses and meat composition in rabbits. 8th World Rabbit Congress, Puebla City, Mexico. Proceedings: pp. 1435-1440.
- Pla, M., Guerrero, L., Guardia, D., Oliver, M.A., & Blasco A. (1998). Carcass characteristics and meat quality of rabbit lines selected for different objectives: I. Between lines comparison. *Livestock Production Science*, 54, 115–123.
- Pogány Simonová, M., Chrastinová, L., Mojto, J., Lauková, A., Szabóová, R., & Rafay J. (2010). Quality of rabbit meat and phyto-additives. *Czech Journal of Food Sciences*, 28, 161–167.
- Polak, T., Gašperlin, L., Rajar, A., & Žlender B. (2006). Influence of genotype Lines, Age at Slaughter and Sexes on the Composition of Rabbit Meat. *Food*

- Technology and Biotechnology, 44,1, 65-73.
- Ristovski, T., Micevski, C., & Janevski, B. (2004). Одгледување и болести кај домашните зајаци. Priracnik, Skopje, Republic of Macedonia.
- Skandro, M., Čaklović, F., Gagić, A., Kobilic-Zubčević V., Smajlović, M., Kustura A., & Goletić T. (2004). Slaughter yield of rabbit carcasses produced in besieged Sarajevo during war years 1992-1995. 27th World veterinary congress, Tunis. Proceedings: 52(3), 273-281.
- Skandro, M., Tariq, A., Alić, B., Goletić, T., & Kostura A. (2008). Kvalitativno - kvantitativna svojstva mesa novozelandskog bijelog kunića. First Croatian Journal on Meat - Meso, 10(2): 122-126.
- Skandro, M. (1998). Mali kuhar za jela od mesa kunića. Izdavač – Grafo Art, Sarajevo, Bosnia and Hercegovina.
- Tafro, N., Urošević, M., Panić, M., Bursać, N., & Dizdarević D. (1989). Tovne vrijednosti kunića hibrida "Emiliano". Savez veterinarara i veterinarskih tehničara Jugoslavije XIII Jugoslovensko savetovanje veterinarara. Zbornik kratkih sadržaja, Primošten, Croatia Proceedings: 21-25.
- Urošević, M., Panić, M., Petrović, & Tafro, N. (1986). Hranljiva vrednost i klanična svojstva mesa kunića., VII seminar inovacije u stočarstvu, Univerzitet u Beogradu – Zemun, Serbia. Proceedings: 16-24.
- Wood, J.D., Richardson R.I., Nute G.R., Fischer A.V., Campo M.M., Kasapidou E., Sheard P.R., & Enser M. (2003). Effects of fatty acids on meat quality: A review. Meat Science 66, 21-32.
- Yalçın, S., Onbaşilar, E., & Onbaşilar, I. (2006). Effect of Sex on Carcass and Meat Characteristics of New Zealand White Rabbits Aged 11 Weeks. Asian-Australian. Journal of Animal Science 19(8):1212-1216

КВАНТИТАТИВНИ КАРАКТЕРИСТИКИ НА ХИБРИДИТЕ ЗАЈАЦИ

**Дарко Андроников¹, Ацо Кузелов², Јулија Саздова³, Киро Мојсов¹, Ацо Јаневски¹,
Соња Јордева¹, Сашка Голомеова-Лонгурова¹**

¹Универзитет „Гоце Делчев“, Технолошко-технички факултет - Штип,
ул. „Крсте Мисирков“ 10-А, 2000 Штип, Република Северна Македонија

²Универзитет „Гоце Делчев“, Земјоделски факултет - Штип,
Крсте Мисирков 10-А, 2000 Штип, Република Северна Македонија

³СОУ „Коле Нехтенин“ - Штип, Република Северна Македонија

*Контакт автор: darko.andronikov@ugd.edu.mk

Резиме

Како материјал за ова истражување ни послужија хибриди на вкрстени единки од т.н. белгиски оријаши и новозеландски бели зајаци (ОН), кои беа хранети ад либитум со индустриски пелетирана храна. Во истражувањето беа опфатени шест зајаци и тоа три машки и 3 три женски. Дефинираната жива маса од 1800 до 2500 г зајациите ја постигнаа за 77 дена, по што се изврши колењето. Во трудот се изнесени резултатите од истражувањето на товните и клавните својства, како и утврдување на составот на месото од зајаци. Достигната е просечна жива маса на машките зајаци од 2792 г, и рандман од 47,92%, додека кај женските зајаци просечната жива маса беше 2695 г и рандман од 48,61%. Просечен удел на задниот дел од трупот кај машките/женските зајаци 28,02% и 27,34%, на грбниот дел 20,93% и 22,65%, додека на предниот дел на трупот 25,26% и 25,39%.

Клучни зборови: зајаци, хибриди, жива маса.