

# HPLC Analysis of Hydroxycinnamic Acid Derivatives in Smederevka and Chardonnay Wines

Violeta Ivanova<sup>1,2</sup>, Emmanuelle Meudec<sup>3</sup>, Jean-Marc Souquet<sup>3</sup>, Borimir Vojnoski<sup>2</sup>, Veronique Cheynier<sup>3</sup>, Marina Stefova<sup>1</sup>

<sup>1</sup>Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Sts. Cyril and Methodius University, Skopje, Republic of Macedonia

<sup>2</sup>Department for Enology, Institute of Agriculture, Sts. Cyril and Methodius University, Skopje, Republic of Macedonia

<sup>3</sup>UMR 1083 Sciences Pour l'Oenologie, INRA, Montpellier, France

e-mail:violeta.ivanova@zeminst.edu.mk



## INTRODUCTION

Phenolic compounds are considered as main factors responsible for the quality of grapes, and thus for their corresponding wines, and also, they are specific for different cultivars. Wine phenolics belong to two main groups: nonflavonoids and flavonoids. The major nonflavonoid phenolic compounds of white wines are hydroxycinnamic acid derivatives, such as caffeoyltartaric (caftaric) acid, *p*-coumaroyltartaric (*p*-coutaric) acid and feruloyltartaric (fertaric) acid (Fig. 1). For white wine production, maceration is kept to a minimum and seldom lasts more than few hours. White wines are usually made at low temperatures (14–18 °C). The juice runs freely from the crushed grapes, which are protected with SO<sub>2</sub> to prevent the enzymatic oxidation.

Hydroxycinnamic acid derivatives	R <sub>1</sub>	M <sub>r</sub>
Coutaric acid	H	296
Caftaric acid	OH	312
Fertaric acid	OCH <sub>3</sub>	326




Fig. 1. Structures of hydroxycinnamic acids derivatives

The presented study represents the first attempt to analyze Macedonian white wines for their polyphenolic acid content with HPLC: **Smederevka**, as typical for the Balkan region and the most wide spread variety at Macedonian vineyards, and **Chardonnay**, as well known grape variety, in order to compare them and correlate the contents with the winemaking protocols.

## MATERIALS AND METHODS

Macedonian white wines, Smederevka and Chardonnay (*Vitis vinifera* L.) were subject of investigation. Winemaking procedures for both varieties included addition of two doses of SO<sub>2</sub> (50 and 100 mg/L) and two yeasts for fermentation (Vinalco and Levuline). A reversed phase liquid chromatographic method was used for identification and quantification of hydroxycinnamic acid derivatives in the wines. Separation of the components, by direct injection of the wines into HPLC (Waters 2690 system), was performed using reversed-phase Atlantis dC18 column, monitored at 320 nm. The mobile phase consisted of water/formic acid (99:1; solvent A), and acetonitrile/water/formic acid (80:19:1; solvent B). In addition, HPLC-MS (Waters 2690 system equipped with ThermoFinnigan LCQ Advantage ion trap mass spectrometer) analysis was carried out to confirm the identity of the separated compounds, recording the spectra in negative ion mode.

## RESULTS AND DISCUSSION

### MS and UV-Vis identification




Fig. 2. MS and UV-Vis spectra of Caftaric and Coutaric acids identified in Smederevka and Chardonnay wines

Hydroxycinnamic acid derivatives, caffeoyltartaric (caftaric) acid at *m/z* 311 (fragment ions: *m/z* 179, 149) and maximum absorbance at 327.9 nm) and *p*-coumaroyltartaric (coutaric) acid at *m/z* 295 (fragment ion at *m/z* 163, 149 and maximum absorbance at 313.6 nm) have been detected in the wines. These compounds produce same fragment ion [M-H]-132] which corresponds to loss of tartaric acid residue.

Thus, molecular ion [M-H]<sup>-</sup> at *m/z* 311 after fragmentation produces two fragments, [M-H]<sup>-</sup> at *m/z* 179, corresponding to caffeic acid and [M-H]<sup>-</sup> at *m/z* 149, obtained after elimination of tartaric acid. This compound was identified as *trans*-caffeoyltartaric acid or *trans*-caftaric acid.

Molecular ion [M-H]<sup>-</sup> at *m/z* 295 giving two fragment ions, [M-H]<sup>-</sup> at *m/z* 162.9 corresponding to the *p*-coumaric residue and [M-H]<sup>-</sup> at *m/z* 149, corresponding to the tartaric acid residue, was identified as *cis*-*p*-coumaroyltartaric acid or *cis*-coutaric acid.

### Quantification

Table 1. Content of hydroxycinnamic acid derivatives in Chardonnay wines

Compounds	Ch-Mac-50	Ch-Mac-100	Ch-Fr-50	Ch-Fr-100
<i>trans</i> -Caftaric acid	57.14	132.18	81.00	142.18
<i>cis</i> -Coutaric acid	25.71	31.37	33.10	32.35
<i>trans</i> -Coutaric acid	30.90	47.03	28.00	81.35
Total	113.75	210.58	142.10	225.88

Table 2. Content of hydroxycinnamic acid derivatives in Smederevka wines

Compounds	Sm-Mac-50	Sm-Mac-100	Sm-Fr-50	Sm-Fr-100
<i>trans</i> -Caftaric acid	3.62	7.17	3.33	11.78
Coutaric acid ( <i>trans</i> + <i>cis</i> )	11.43	26.87	13.77	27.81
Total	15.05	34.04	17.1	39.59

Labels: Ch-Chardonnay, Sm-Smederevka, Mac-Macedonia yeast, Vinalco, Fr-French yeast, Levuline, 50– 50 mg/L SO<sub>2</sub>, 70– 70 mg/L SO<sub>2</sub>

Chardonnay wines were richer with phenolic acid derivatives compared to Smederevka wines. The dominant component in Chardonnay wines was *trans*-caftaric acid, while, *trans*-coutaric acid dominated in Smederevka wines. Regarding the influence of SO<sub>2</sub>, wines with higher dose of SO<sub>2</sub> contained higher levels of acids, since SO<sub>2</sub> suppress the activity of oxidases, preventing oxidation of these readily oxidizable phenols and wine browning. The influence of the yeast on the content of hydroxycinnamic acid derivatives was not significantly different. Principal component analysis (Fig. 3) was performed in order to check if the studied wines can be distinguished, observing separation of the samples according to the variety and SO<sub>2</sub> doses.

### Statistical analysis




Fig.3 Principal Component score plot (A) and correlation scatterplot (B) of the variables with PC1 and PC2 based on hydroxycinnamic acids for the analyzed Smederevka and Chardonnay wines

**Acknowledgements:** The work was supported by the FP7 project CHROMLAB-ANTIOXIDANT.