

THE EFFECT OF ALVOGYL IN THE POSTEXTRACTED PERIOD

Dimova Cena PhD, Evrosimovska Biljana Ms. Sci., Popovska Lidija Ph D, Muretzani Arben

AIM

The aim of this study was to show the analgesic effect of ALVOGYL in post extracted pain conditions.



ALVOGYL

Jodoform	15,8g
Paraminobenzoat butyl	25,7g
Eugenol	13,7g
Penghavar	3,5g
Xscipient k.s.p.	100g

PATIENTS AND METHODS


The study included 30 patients, divided in three groups of ten subjects, as follows:

- First group with typical extractions (with clinical diagnosis Paradontitis chronic in the posterior teeth),
- Second group with atypical extractions (with the same diagnosis as the first group and separation of the roots)and,
- Third group with operative extractions of semiimpacted third molar.


RESULTS

The patients were evaluated during the period of two weeks. Post extraction pain was established in eighteen patients (60%) who were treated with ALVOGYL in production of Septodont, France. The reduction of pain with ALVOGYL was shown 15-20 min after applying the medicament.

N = 30	First application of alvogyl		Second application of alvogyl
	I & II day	III day	
First group (n=10) (typical extractions)	3 (33%)	1 (10%)	0
Second group (n=10) (atypical extractions)	4 (44%)	2 (22%)	1 (10%)
Third group (n=10) (operative extractions)	5 (55%)	3 (33%)	2 (22%)
Total N' = 18	12 (66%)	6 (33%)	3 (16%)



Extraction wound after 48 hours



Applied alvogyl in extraction wound



Extraction wound after 5 days

FIRST GROUP



SECOND GROUP



THIRD GROUP

CONCLUSION

A traumatic work during the tooth extraction should be imperative in preventing complications like after extraction pain, and timely controlling the healing of the extraction wounds.