

Српско геолошко друштво

Зборник радова XVI Конгреса геолога Србије



Proceedings of the XVI Serbian Geological Congress

ОПТИМАЛНО ИСТРАЖИВАЊЕ И ОДРЖИВО КОРИШЋЕЊЕ ГЕОЛОШКИХ РЕСУРСА

OPTIMAL RESEARCH AND SUSTAINABLE USAGE OF THE GEOLOGICAL RESOURCES



XVI Конгрес геолога Србије: Зборник радова (Национални конгрес с међународним учешћем)

XVI Serbian Geological Congress: Proceedings (National Congress with International Participation)

Donji Milanovac, 22-25.05.2014.

За издавача / For the Publisher Zoran Stevanović, Председник Српског геолошког друштва / President of the Serbian Geological Society

> Главни уредник / Editor-in-Chief Vladica Cvetković

Уређивачки одбор / Editorial Board

Biljana Abolmasov, Katarina Bogićević, Meri Ganić, Rade Jelenković, Aleksandra Maran Stevanović, Vesna Matović, Vesna Ristić Vakanjac, Aleksandar Ristović, Ljupko Rundić

Техничка припрема / Technical Preparation Veljko Marinović, Ognjen Jevtić, Milan Vukićević, Nemanja Krstekanić, Jelena Krstajić, Nikola Vojvodić, Bojan Kostić, Kristina Šarić

Издавач / Publisher

Српско геолошко друштво / Serbian Geological Society Kamenička 6, P. Box 227, 11001 Belgrade, Serbia http//:www.sgd.rs; e-mail: office@sgd.rs Тираж: 300 примерака / Circulation: 300 copies

Штампа / Printing Šprint, Alekse Nenadovića 28, Beograd, Serbia

ISBN 978-86-86053-14-5

Напомена: *Аутори су одговорни за садржај и квалитет својих саопштења* Note: The authors are responsible for the content and quality of their contributions



ХVІ КОНГРЕС ГЕОЛОГА СРБИЈЕ

Чланови одбора XVI SERBIAN GEOLOGIAL CONGRES Committee Members

Почасни одбор / Honorary Committee

Zoran Maksimović, Stevan Karamata, Vidojko Jović, Aleksandar Grubić, Jelena Obradović, Ivan Antonijević, Rajka Radojčić, Zora Sučić-Protić, Branislav Ćirić, Radule Popović, Jovanka Mitrović, Nadežda Krstić, Mihajlo Kalenić, Ivan Filipović

Почасни одбор (институције) / Representatives of the Institutions

Adam Dangić (Društvo geoloških inžinjera i tehničara Srbije), Ivan Obradović (Univerzitet u Beogradu – Rudarsko-geološki fakultet), Dragoman Rabrenović (Geološki zavod Srbije), Milutin Ignjatović (CIP, Beograd), Srđan Stefanović (NP Đerdap), Vlastimir Trujić (Institut za rudarstvo i metalurgiju, Bor), Milan Dimkić (Institut za vodoprivredu Jaroslav Černi), Dragan Mitrović (Geološki zavod Republike Srpske), Vashkevich Aleksey Alexandrovich (Gazprom Neft), Alexander Rodionov (NIS)

Организациони одбор / Organizing Committee

Zoran Stevanović (predsednik), Nenad Banjac (potpredsednik), Ljupko Rundić (potpredsednik), Tivadar Gaudenyi (tehnički sekretar), Boško Jevtović (tehnički sekretar), Milovan Rakijaš (blagajnik), Petar Stejić, Nenad Grubin, Mile Bugarin, Dejan Koželj, Aleksandra Maran Stevanović, Vesna Lesić, Marija Kukavica

Научни одбор / Scientific Committee

Vladica Cvetković (predsednik), Milan Sudar, Dragan Milovanović, Radmila Pavlović, Vladan Radulović, Vesna Poharc-Logar, Aleksandar Đorđević, Rade Jelenković, Veselin Dragišić, Nebojša Vasić, Slobodan Knežević, Divna Jovanović, Dragutin Jevremović, Slobodan Vujasinović, Aleksandar Kostić, Spomenko Mihajlović, Ljiljana Karanović, Ivan Dulić, Dejan Prelević, Milenko Burazer

Студенти волонтери / Students Congress volunteers

Jelena Krstajić, Marina Prodanović, Veljko Marinović, Milan Vukićević, Ognjen Jevtić, Nemanja Krstekanić, Bojan Kostić, Marina Zečević

САДРЖАЈ / CONTENT

Aleksandar Orubie	
ONEGEOLOGY – GEOSCIENCE DATA AT THE TOUCH OF A BUTTON Marko Komac, Francois Robida, Tim Duffy	
STANJE I POTENCIJALI MINERALNIH RESURSA ISTOČNE SRBIJE STATUS AND POTENTIAL OF MINERAL RESOURCES OF EASTERN SERBIA Rade Jelenković	
O KENOZOJSKOM MAGMATIZMU SRBIJE – REČIMA KOJE RAZUME SVAKI GEOLOG ABOUT CENOZOIC MAGMATISM IN SERBIA – BY WORDS UNDERSTANDABLE TO EVERY GEOLO Vladica Cvetković	GIST
СТРАТИГРАФИЈА, ПАЛЕОНТОЛОГИЈА И ТЕКТОНИКА STRATIGRAPHY, PALAEONTOLOGY AND TECTONICS	
RHODOPE EVOLUTION IN THE HEART OF BALKAN GEOLOGY Ivan S. Zagorchev	
CIKLIČKA SEDIMENTACIJA NA RUBOVIMA MARINSKOG ŠELFA ZA VRIJEME SREDNJEG I GORI PERMA U CENTRALNOM DIJELU VELEBITA (HRVATSKA) CYCLIC SEDIMENTATION IN MARGINAL MARINE SHELF ENVIRONMENT AT THE MIDDLE/UPPER PERMIAN BOUNDARY IN CENTRAL PART OF THE VELEBIT MT. (CROATI Ana Šimičević, Jasenka Sremac	NJEG (A)
THE LAKE PANNON – SERBIAN SIDE OF THE STORY Dejan Radivojević, Imre Magyar, Marten ter Borgh, Ljupko Rundić	
STABLE ADRIA AND THE EXTERNAL DINARIDES: TECTONIC MODELS IN THE CONTEXT JURASSIC-CRETACEOUS PALEOMAGNETIC RESULTS Emő Márton, Vlasta Ćosović, Alan Moro	T OF
АНИЗИЈСКИ КАРБОНАТИ ЦРМНИЦЕ И ОКОЛИНЕ (У ЦРНОЈ ГОРИ) ANISIAN CARBONATES OF CRMNICA AND SURROUNDINGS (IN MONTENEGRO) Damjan Čađenović, Jelena Milutin, Martin Đaković, Novo Radulović	
MAJOR PHASES OF NEOGENE-QUATERNARY TECTONIC EVOLUTION OF THE SOUTH-EASTERN I OF PANNONIAN BASIN Ivan Dulic, Snežana Marjanovic, Vladislav Gajic, Goran Bogicevic	PART
СТРАТИГРАФСКИ МОДЕЛ КВАРТАРА ЗА РЕЧНЕ ПОЛИЦИКЛИЧНЕ СЕДИМЕНТЕ НА ПРОСТОРУ СРЕ STRATIGRAPHICAL MODEL OF QUATERNARY FOR FLUVIAL POLYCYCLIC SEDIMENTS IN TERRITORY OF SERBIA	БИЈЕ THE
Draženko Nenadić, Tivadar Gaudenyi ПЛЕИСТОЦЕНСКЕ ФЛУВИЈАЛНЕ НАСЛАГЕ ЈУГОИСТОЧНОГ СРЕМА И ДОЛИНА ДУНАВА И МОРАН PLEISTOCENE FLUVIAL DEPOSITS FROM THE SOUTHEASTERN SREM AND THE VALLEYS OF DANUBE AND MORAVA Draženko Nenadić Katarina Rogićević	BE THE
MIDDLE JURASSIC RADIOLARIANS FROM OPHIOLITIC MÉLANGE OF DINARIDE OPHIOLITE 2 (DOZ) - AGE AND GEODYNAMIC SIGNIFICANCE (PAPRATNICA RIVER, NORTHERN BOSNIA HERZEGOVINA)	ZONE AND
Branimir Šegvić, Alan Vranjković, Duje Kukoč, Vlatko Brčić, Špela Goričan, Elvir Babajić, Hazim Hrvatović Dragičević, Majda Pađen	, Ivan
СЕЛИМЕНТОЛОГИЈА И БИОСТРАТИГРАФИЈА САНТОНСКО-КАМПАНСКИХ КРЕЧЊАКА ОКОЈ	ЛИНЕ

ANALIZA I OBRADA KAROTAŽNIH MERENJA ZA POTREBE IZRADE SEIZMIČKE INVERZIJE I PROGNOZE PETROFIZIČKIH PARAMETARA NA OSNOVU 3D SEIZMIČKIH PODATAKA WELL LOG PROCESSING AND ANALYSIS FOR SEISMIC INVERSION AND PREDICTING OF PETROPHYSICAL PARAMETERS DISTRIBUTION Stevan Doroškov, Aleksandar Ristović	699
СТАНДАРДИЗАЦИЈА ПОПРАВКИ ПРИ РАЧУНАЊУ ГРАВИМЕТРИЈСКИХ АНОМАЛИЈА У СРБИЈИ STANDARDIZATION IN REDUCTION FOR CALCULATION OF GRAVITY ANOMALIES IN SERBIA Ivana Vasiljević, Snežana Ignjatović, Branislav Sretenović, Oleg Odalović	708
ГЕОМАГНЕТСКИ ПРЕМЕР НА ТАЧКАМА ДРУГОГ РЕДА РЕПУБЛИКЕ СРБИЈЕ GEOMAGNETIC SECOND ORDER SURVEYS AT SERBIA D.Andrejić, D. Popeskov, M. Mihajlčić	718
PRAĆENJE PROMENA GEOMAGNETSKOG POLJA, PARAMETARA ATMOSFERE I ZEMLJOTRESA U PROJEKTU FP7 BLACKSEAHAZNET MONITORING OF EARTHQUAKES, GEOMAGNETIC FIELD AND ATMOSPHERE PARAMETER IN FP7 BLACKSEAHAZNET PROJECT Milena Cukavac, Strachimir Cht. Mavrodiev, Lazo Pekevski, Spomenko J. Mihajlovic	728
PRIMENA SEIZMIČKE INVERZIJE I MULTIATRIBUTIVNE ANALIZE PRI INTERPRETACIJI 3D SEIZMIČKIH PODATAKA KIKINDE 2013 THE APPLICATION OF SEISMIC INVERSION AND MULTIATTRIBUTE ANALYSIS FOR INTERPRETATION OF 3D SEISMIC DATA KIKINDA 2013 Janko Sovilj	733
ЕКОНОМСКА ГЕОЛОГИЈА И РУДНА ЛЕЖИШТА ECONOMIC GEOLOGY AND ORE DEPOSIT	737
GEOLOGY AND MINERALIZATION OF THE NEW COOPER AND GOLD DISCOVERY SOUTH OF BOR – TIMOK MAGMATIC COMPLEX Miodrag Banješević, Duncan Large	720
ОДРЖИВО ПЛАНИРАЊЕ АГРЕГАТА У ЈУГОИСТОЧНОЈ ЕВРОПИ - ЗАШТО И КАКО SUSTAINABLE AGGREGATES PLANNING IN SOUTHEAST EUROPE – WHY AND HOW?	739
GEOLOŠKE KARTE KAO PODLOGE ZA ODRŽIVO PLANIRANJE SNABDEVANJA AGREGATIMA (NA PRIMERU REPUBLIKE HRVATSKE) GEOLOGICAL MAPS AS BACKGROUND FOR SUSTAINABLE AGGREGATE SUPPLY PLANNING (EXAMPLE OF THE REPUBLIC OF CROATIA) Vladimir Šimić, Slobodan Miko, Boris Kruk, Željko Dedić, Dragana Životić, Nevena Andrić, Zoran	741
PROCEDURE FOR MARBLE CLASSIFICATION FROM BOREHOLES WITH PARTICULAR REFERENCES TO THE SIVEC MINE Mice Trkaleski, Blazo Boey, Ilias Rigoupolous	743
QUALITATIVE-QUANTITATIVE CHARACTERISTICS OF THE MARBLES FROM TUMBA AREA (REPUBLIC OF MACEDONIA) AND OPPORTUNITIES FOR THEIR EXPLOITATION	745
GENERAL REVIEW OF MINERAL ECONOMY IN MODERN CONDITIONS OF SUSTAINABLE DEVELOPMENT	751
Radule Tošović	756
ZNACAJ PRIMENE MODERNOG I EFIKASNOG MENADŽMENTA LJUDSKIH RESURSA U MINERALNOM SEKTORU THE IMPORTANCE OF MODERN AND EFFECTIVE HUMAN RESOURCE MANAGEMENT IN THE MINERAL SECTOR Radule Tošović	745
QUALITATIVE-QUANTITATIVE FEATURES OF CLAY FROM THE SITE BAZERNIK (WESTERN MACEDONIA) Orce Spasovski, Daniel Spasovski	765
CHARACTERIZATION OF THE ORIGIN OF THE MTE ANOMALIES IN THE "CAP DE GARDE", EDOUGH MASSIF, ANNABA (NE ALGERIA) Soraya Hadj Zobir, Menana Daif, Saadane Djorfi	776

КВАРЦНИ ПЕСАК И ПЕСКОВИТА ГЛИНА ЛЕЖИШТА БОШЊАНЕ КАО СИРОВИНА ЗА ЦЕМЕНТНУ ИНДУСТРИЈУ QUARTZ SAND AND SANDY CLAY FROM THE BOŠNJANE DEPOSIT AS RAW MATERIAL FOR CEMENT INDUSTRY Miroslava Maksimović, Milenko Jovanović, Slađana Krstić, Goran Pačkovski, Marinko Pavlović	777
GEOCHEMICAL CHARACTERISTICS OF POLYMETALLIC DEPOSIT KAZANDOL (R. MACEDONIA) Jordanka Nikolova, Orce Spasovski, Daniel Spasovski	785
ТЕХНОГЕНО ЛЕЖИШТЕ НА ПРОСТОРУ СТАРОГ ФЛОТАЦИЈСКОГ ЈАЛОВИШТА У БОРУ (ПОЉЕ 1 И ПОЉЕ 2) ТЕСНЛОGENIC DEPOSIT IN THE AREA OF THE OLD FLOTATION TAILING DUMP IN BOR (FIELD 1 AND FIELD 2) Miroslava Maksimovic, Mile Bugarin, Vladan Marinković, Zoran Stevanović	789
DISPERZNA ORGANSKA MATERIJA U LEŽIŠTU BORATA PISKANJA (SRBIJA) POREKLO I USLOVI DEPOZICIJE DISPERSED ORGANIC MATTER IN THE PISKANJA BORATE DEPOSIT (SERBIA): SOURCE AND	
DEPOSITIONAL ENVIRONMENT Nevena Andrić, Dragana Životić, Vladimir Simić	799
3D MODEL LEŽIŠTA "RAVNA REKA"- BLOK IV 3D MODEL OF THE DEPOSIT "RAVNA REKA"- BLOCK IV Nikola Pašajlić	802
ИСТРАЖИВАЊА УГЉА У ДЕСПОТОВАЧКОМ БАСЕНУ EXPLORATION OF COAL IN THE DESPOTOVAC BASIN Svetlana Simić	810
ПОДРУЧЈЕ ХОМОЉА ГЕОЛОШКО-ЕКОНОМСКИ ЦЕНТАР КАМЕНА (АГК И ТГК) СА АСПЕКТА ПЕРСПЕКТИВНОСТИ, ВРСТЕ, КВАЛИТЕТА, КОЛИЧИНА И УПОТРЕБЉИВОСТИ ТНЕ НОМОLJE AREA, ТНЕ GEOLOGICAL AND ECONOMIC CENTRE OF STONE,	
WELL AS THE USABILITY OF THE DECORATIVE STONE IN SERBIA Cvetko Živković	816
ΓΕΟΕДУΚΑЦИЈА, ЗАШТИТА ГЕОНАСЛЕЂА И ГЕОЕКОЛОГИЈА GEOEDUCATION, GEOHERITAGE AND GEOECOLOGY	825
GEOZAŠTITA U SRBIJI: PROŠLOST, SADAŠNJOST, BUDUĆNOST GEOCONSERVATION IN SERBIA: PAST, PRESENT, FUTURE Aleksandra Maran Stevanović	827
УЛОГА И ЗНАЧАЈ ГЕОЕДУКАЦИЈЕ У САВРЕМЕНОМ ДРУШТВУ ROLE AND IMPORTANCE OF GEOEDUCATION IN THE MODERN SOCIETY Ljupko M. Rundić, Marina M. Ilić	832
METOДОЛОШКЕ ОСНОВЕ ЗА ПРОЦЕНУ ОБЈЕКАТА ГЕОНАСЛЕЂА METHODOLOGICAL GUIDELINES FOR ASSESSMENT OF GEOHERITAGE SITES Aleksandra Maran Stevanović	842
"UPOZNAJ ZEMLJU - ZAVIRI U MIKROSKOP!" - PROMICANJE GEOLOGIJE I GEOLOŠKE BAŠTINE "MEET THE EARTH-PEEK UNDER THE MICROSCOPE!" - PROMOTION OF GEOLOGY AND GEOHERITAGE PROTECTION	846
Lidija Galović, Marija Bošnjak заштита и презентација геолошких објеката у урбаним условима	0+0
THE PROTECTION AND PRESENTATION OF GEOLOGICAL SITES IN THE URBAN ENVIRONMENT Erna Šehovac, Milan Popović, Srećko Stefanović, Branka Lazović, Ljupko Rundić Slobodan Knežević	850
POTENTIALLY TOXIC METALS WITHIN THE ZATON-BULBA KARST SYSTEM (MEHEDINIT PLATEAU, ROMANIA) Cristian-Mihai Munteanu, Andrei Giurginca, Marius Vlaicu	857
POTENCIJALNO TOKSIČNI MIKROELEMENTI U MRKIM UGLJEVIMA SRBIJE POTENTIALLY TOXIC TRACE ELEMENTS IN SERBIAN BROWN COALS Dragana Životić, Vladimir Simić, Ivan Gržetić	. 860
GEOCHEMICAL PROPERTIES OF SOIL AND ATTIC DUST AROUND THE BOR OPET PIT AND Cu SMELTER (SERBIA) Robert Šajn, Trajče Stafilov, Suzana Erić	864
ОСНОВНЕ ВРЕДНОСТИ ПОДРУЧЈА ЂЕРДАПА, ПОТЕНЦИЈАЛНОГ ГЕОПАРКА MAIN VALUES OF THE DJERDAP AREA, POTENTIAL GEOPARK Dragoman Rabrenović, Srdjan Belij, Ivana Mojsić, Mališa Mladenović	. 866

QUALITATIVE QUANTITATIVE FEATURES OF CLAY FROM THE SITE BAZERNIK (WESTERN MACEDONIA)

Orce Spasovski and Daniel Spasovski

Faculty of Natural and Technical Sciences, "Goce Delčev" University, Goce Delčev 89, MK-200, Štip, Republic of Macedonia orce.spasovski@ugd.edu.mk

Key words: deposit, Bazernik, clay, schist, qvarcporfire, qualitative and quantitative characteristics, mineralogical compound, chemical compound.

Abstract: Clay schist and clay from the vicinity of Bazernik - Demir Hisar in the future may represent an important economic entity in the production of clays in Macedonia, requires monitoring of economic parameters in function of current trends in assessment of the justification of the use of clay and clay schist and the possibility of valuation and associated components too.

Following the release mode, multi-layered system of cracks and cracks degree of compactness and purity fragmented clay enabled the determination of the qualitative and quantitative characteristics of the clay. The clay schist and the clay from the deposit Bazernik, based on their qualitative characteristics, can be found in their natural state, and might be applied in the production of acid-resistant ceramic products with great quality. They can be found in their enriched state, so they might be applied in the paper industry, cosmetic industry etc.

INTRODUCTION

The site Bazernik is located near the village Bazernik, an area that territorially is a part of the municipality of Demir Hisar. It is located at a distance of 50km from Bitola and 10 km from the railway station Sopotnica (Fig. 1). The site Bazernik is explored with excavation on the surface, and within the scope of 1020 using pit exploration. Regarding this exploration it is determined that the site is comprised of white clay, clay shist, as well as green metamorphic quartz porphyry. The clay and the clay shist represent a mineral mass and the metamorphic quartz porphyry is a slag.



Figure 1. Map of the R. Macedonia with position the Bazernik lokality

The clays and the clayed schist from the surroundings of Bazernik – Demir Hisar may represent an important economical subject for the clay production in Macedonia in the future. The monitoring of the economical parameters for the contemporary trends allows to evaluate the justification of the clays and clay schist usage and the possibility to evaluate their accompanied components.

Within the hitherto explorations, there is a particular attention on the radio – structural explorations, which have determined the existence of the main mineral stages during the creation of the clay and clay schist layers.

The site Bazernik has been discovered in 1969 when the basic geological data has been collected (Zatkovski, 1969). In 1970 a program for detailed exploration of the site has been developed, and within the same year a detailed explorations regarding tachometric recording, geological mapping of the site, chemical examinations, diggings, technological examinations, chemical analysis, radio and petrographic examinations have been conducted (Zatkovski, 1971). A mine geodesic recording, mapping of the mine exploration, undermining and excavations have also been conducted. The newest data about the qualitative and quantitative characteristics of the site Bazernik can be found in Jovanov's papers (2010)

GEOLOGICAL CHARACTERISTICS

The geological composition of the mineral body is determined based on the geological mapping and the surface and pit explorations.

The mineral body "Bazernik" has a shape of an elongated body comprised of white clay, dark – green metamorphic quartz porphyry, clay schist and white rough metamorphic quartz porphyry. Given that the diameters do not intersect the mineral body, it wasn't possible to establish if the top part is comprised solely of metamorphic quartz porphyry, as it was predicted on the transversal geological profiles.

The clay and the clay schist represent a mineral matter, whereas the white and the dark – green metamorphic quartz porphyry represent non – mineral matter (Fig. 2).

The clay (Fig. 3), seen macroscopically, has white color and plastic properties and it's comprised of rough and fine - granulated fractions. The first fraction is represented with 44 - 66 % with grains bigger than 0.06mm, and the second fraction with microscopical grains. Using radioscopic thermodiferential examinations it is determined that the mineral mass consists of kaolin and illit. There are quartz grains and fragments of a sound rock mass in the clay matter. It's found on the base part of the mineral body and it's shaped like an irregular wire 3-8 meters thick. It declines astride northeast under variable declining angle from 15 to 65 degrees. It is also found with the same structure within the native rock, having well preserved elements of schist with clearly expressed forms of the clay feldspars.

The clay schist (Fig. 4), macroscopically speaking, has greenish white colour and rather small firmness and hardness. When pressed insignificantly with the fingers it decomposes in a fine decomposition comprised of leaf – like fragments and clay matter. It's comprised of white clay and schist matter that consist of basic matter built from Si matter, sericite, chlorite, epidote, zoisit, a little allanite, in which there are Phenoklasts and rarely K – feldspars. It declines astride northeast under variable declining angle.

The dark – green metamorphic quartz porphyry is situated in the central part of the mineral body, between the white clay and the clay schist. It has a remarkable firmness and hardness. It is comprised of a basic, significantly siliphied matter with sericite and fine – dispersed opaque matter. Within the basic matter there are phenoplasts comprised of quartz and certain amounts of K – feldspare.





1.purple and other schist, 2. white rough metamorphic quartz porphyry, 3. clay schist, 4. white clay, 5. dark – green metamorphic quartz porphyry, 6. gray crystalline limestone, 7. route the cross section, 8. tag to undermine and shaft.

The whitish quartz porphyry macroscopically has a greenish white colour. The schist surfaces are streaked with manganic dendrites. Under compression it decomposes in parallelopipedic decomposition. It's mineralogically comprised of partially syliphied and carbonized basic matter weaved with fine dispersed opaque matter, where quartz aggregates and sericit relics can be seen. The phenoclasts are comprised of quartz and partially carbonized schist that declines astride northeast under viable declining angle.



Figure 3. White clay at the deposit of Bazernik



Figure 4. Clay schist at the deposit of Bazernik

QUALITATIVE CHARACTERISTICS

In order to distinguish the qualitative characteristics of the clay and the clay schist from the site Bazernik, representative samples are used for distinguishing the main components. In the Table 1 are shown the given results from the completed chemical explorations of the clay. From the given chart we can see that the clay consist of high contents of SiO₂ with medium composition of 71.46 %, followed by Al₂O₃ with medium composition of 12.12 %, and the Fe, Fe₂O₃, CaO, K₂O are least included. In the Table 2 are shown the given results from the completed chemical exploration on the clay schist. According to the chart the SiO₂ and Al₂O₃, like in the clay, are the most included, and the Fe, Fe₂O₃, CaO, K₂O are less included. According to the charts we can conclude that the clay and the clay schist have almost equable chemical composition.

Compo-		Average										
nents	1'	2'	3'	4	9'	12'	11'	14'	17	21 [']	n	composition
G z	3.86	4,46	4,32	3,86	4,63	3,71	5,97	7,01	4,75	4,83	10	4.65
SiO ₂	74.91	72,37	72,12	71,82	70,99	73,97	70,04	68,36	69,46	70,56	10	71.46
Fe ₂ O ₃	1.50	1,80	1,90	2,10	2,80	2,00	3,80	3,00	2,60	3,29	10	2.47
Fe	1.04	1,25	1,32	1,45	1,94	1,39	2,54	2,09	1,80	2,25	10	1.70
FeO	1.20	1,36	1,30	1,52	1,68	1,10	0,58	0,20	2,05	0,40	10	1.14
Al ₂ O ₃	11.99	12,42	13,18	12,86	12,35	12,16	11,96	9,88	12,01	12,44	10	12.12
TiO ₂	0.14	0,18	0,20	0,14	0,20	0,20	0,18	0,12	0,17	0,14	10	0.16
CaO	3.12	3,90	3,86	3,44	2,86	2,86	3,14	5,72	3,12	4,05	10	3.61
MqO	1.33	1,52	1,42	1,53	1,47	1,50	1,68	,66	2,28	1,68	10	1.71
CO ₂	0.20	0,18	0,30	0,20	0,32	0,20	0,36	5,04	0,75	0,41	10	0.80
Na ₂ O	1.20	1,30	1,30	2,30	1,30	1,00	2,30	1,30	1,65	0,95	10	1.46
K ₂ O	1.95	2,05	1,75	1,95	2,40	2,60	1,85	1,95	1,95	2,10	10	2.05

Table 1. Chemical composition of the clay (%)

 Table 2. Chemical composition of the clay schist (%)

Compo-	Number of sample												Average	
nents	6	7'	8'	10 [°]	13'	15 [°]	18'	19'	20'	22'	23'	24'	n	composition
G z	2.84	4.38	3.16	2.66	4.12	4.11	3.44	4.62	4.38	3.76	4.24	4.30	12	3.84
SiO ₂	72.45	71.90	72.31	73.28	71.78	71.54	72.83	70.97	72.55	71.40	71.93	72.44	12	72.12
Fe ₂ O ₃	2.00	1.60	2.20	4.60	3.24	3.40	20.70	3.40	2.20	2.14	1.60	2.20	12	2.61
Fe	1.39	1.50	1.50	3.20	2.25	2.36	1.86	2.36	1.94	1.46	1.10	1.53	12	1.94
FeO	0.96	1.20	1.36	0.45	0.31	0.30	1.44	1.30	0.90	1.08	0.92	1.32	12	0.96
Al ₂ O ₃	12.87	12.99	12.87	9.36	11.42	11.40	12.22	12.18	11.96	12.38	11.64	12.56	12	12.04
TiO ₂	0.18	0.12	0.19	0.14	0.16	0.22	0.21	0.14	0.17	0.21	0.18	0.14	12	0.17
CaO	3.38	3.90	3.51	3.69	3.82	3.44	3.24	3.43	3.28	3.76	3.16	3.35	12	3.49
MqO	1.33	1,90	1.61	1.52	1.86	2.14	1.96	1.86	1.94	2.10	1.75	1.68	12	1.81
CO ₂	6.40	0.38	0.35	0.42	0.38	0.20	0.25	0.20	0.35	0.34	0.38	0.40	12	0.34
Na ₂ O	3.20	0.70	2.30	2.30	1.65	1.65	1.30	1.30	0.95	2.30	1.20	1.55	12	1.80
K ₂ O	1.75	2.45	1.85	2.10	1.95	2.10	2.10	2.10	1.95	1.95	1.30	1.95	12	1.96

QUANTITATIVE CHARACTERISTICS

Four mineral masses are distinguished with the determination of the clay's and clay schist's quantitative characteristics, which have been explored on a different level.

According to the level of exploration the mineral mass 2 is categorized within the reserves of the A category, the mineral masses 1 and 3 belonged to the B category, while the mineral mass 4 in the C_1 and C_2 category (Fig. 5).



Figure 5. Schematic block diagarm the deposit Bazernik 1.clay, 2. clay schist, 3. surface of clay (m^2) , 4. surface of schist (m^2) , 5. tag category. 6. length (m).

According to the estimation of the mineral reserves, using the Profile method (Jankovic, 1957), (Blecic and Milovanovic, 1999) 7200 tons of the A category and 35. 774 tons of B category are estimated to be clay reserves.

Within these estimations 16,800 tons of clay schist reserves belong to the A category, 52,790 tons to the B category and 836,550 tons to the C_1 category.

73,610 tons of clay reserves and 910,140 tons of clay schist of A, B, C_1 and C_2 category are determined.

CONCLUSION

According to the afore mentioned in the graduate thesis we can conclude the following: The site Bazernik is located at a distance of 50 km from Bitola and 10 km from the railway station Sopotnica. The wider surrounding of the site Bazernik is comprised from Paleozoic and Triassic rocks represented with different schists, partially crystal dark-grey lime stones and metamorphic quartz porphyry, purple schists, double – basis series and strong sediment of crystal multi colored limestone. The site Bazernik is explored with excavation on the surface, and within the scope of 1020 using pit exploration. Regarding this exploration it is determined that the site is comprised of white clay, clay schist, as well as green metamorphic quartz porphyry. The clay and the clay schist represent mineral matter and the metamorphic quartz porphyry is a slag. Within the composition of the mineral body four masses are distinguished divided into A, B, C₁ and C₂ category. There are 73,610 tons of clay reserves, out of which 7200 tons belong to A, 35, 774ons belong to B and 30, 636 tons to C₁ and C₂ category. There are 908, 140 tons reserves of clay schist, out of which 16,800 tons belong to A category, 52, 790 belong to B category and 836, 550 are from C₁ and C₂ category. Using detailed explorations it is determined that the clay and the clay schist can be used as acidulous resistant products in their natural condition. When supplemented they can be used in the food industry, the paper industry, the cosmetics industry and the paints and varnishes industry.

REFERENCES

- Blecic, N., and Milovanovic, D., (1999): Metode proracuna rudnih rezervi. *Rudarsko geoloski fakultet* Univerziteta u Beogradu. pp. 176.
- Jankovic, S., (1957): Examining and estimation of the mineral raw materials reserves. *Faculty of Mining and Geology, Belgrade*. pp. 379.
- Javanov, D., 2010: Kvalitativno kvantitativni karakteristiki na glinite od naogalisteto Bazernik Dimir Hisar. Univerzitet Goce Delcev, Stip, Fakultet za prirodmi i tehnicki nauki. Diplomska rabota. Str. 45.
- Zatkovski, P., (1969): Izvestaj za izvrseni prethodni geoloski istrazuvanja na naogalisteto na glina Bazernik. Str. Fodn na Biro za geoloski i rudarski istrazuvanja – Bitola. pp.25.
- Zatkovski, P., 1971: Elaborate on the geological composition of the mineral reserves on the clay and clay schist site Bazernik. *Department for geological and mining exploration*. pp. 50.