

Българско геологическо дружество
Bulgarian Geological Society



НАЦИОНАЛНА КОНФЕРЕНЦИЯ

с международно участие

NATIONAL CONFERENCE

with international participation

**ГЕОНАУКИ
GEOSCIENCES
2011**

PROCEEDINGS

**СБОРНИК
РАЗШИРЕНИ РЕЗЮМЕТА**

Българско геологическо дружество
Bulgarian Geological Society
Sofia • 2011

Small foraminifers from the Paleogene basins in the Republic of Macedonia

Малки фораминифери от палеогенските басейни в Република Македонија

Violeta Stojanova, Goše Petrov, Violeta Stefanova
Виолета Стоянова, Гоше Петров, Виолета Стефанова

Faculty of Natural and Technical Sciences, University “Goce Delcev”, Štip, R. Macedonia;
E-mail: violeta.stojanova@ugd.edu.mk; gose.petrov@ugd.edu.mk; violeta.stefanova@ugd.edu.mk

Key words: Paleogene, benthic and planktonic foraminifers, basins.

Introduction

Paleogene sediments are broadly spread on the territory of Republic of Macedonia, especially in the central and the eastern part, named Vardar zone and the Serbo-Macedonian massif. According to its distribution, the Paleogene in Republic of Macedonia can be divided into 4 major (larger) basins and few isolated masses, mostly located along the shells and covers with an orientation NW–SE. They are: Tikveš, Ovče Pole, Skopje-Kumanovo and Delčevo basins, as

main ones, and Deve Bair, Valandovo-Gevgelija and Ilovica-Shtuka within the Strumica valley, as isolated blocks (Fig.1).

As a result of the lithostratigraphic studies of the Paleogene sediments taken from the above mentioned basins, 5 superpositionally distributed lithostatigraphic units (lithozones) have been divided: basal, lower flysch, lithozone of yellow sandstones, upper flysch lithozone and carbonate – sandy lithozone.

The geological age of the sediments in all basins has been determined as Upper Eocene to Lower



Fig. 1. Distribution of Paleogene sediments in Republic of Macedonia: ZMZ, Western Macedonian zone; VZ, Vardar zone; SMM, Serbian-Macedonian massif

Basins: 1, Tikveš; 2, Ovče Pole; 3, Skopje-Kumanovo; 4, Delčevo; 5, Deve Bair; 6, Valandovo-Gevgelija; 7, Strumica

Oligocene on the base of many fossil remainings coming from macrofossil groups. The total thickness of the Paleogene sediments (Upper Eocene–Oligocene) is estimated as 3000–3500 m (seemingly, according to some other data the thickness reaches up to 4000 m).

Materials and methods

Sampling of foraminiferal fauna was carried out on open Paleogene cross sections in basins of Macedonia. Technical work was done with classical methods of micropaleontological analysis (break up, washing, drying, selecting and determination). Taxonomic determinations were carried out with a microscope binocular Zeiss, 50 to 80 magnifications. Microphotographs were taken with an electronic microscope JMS-5510-JEOL on selected foraminiferal tests.

Results

As a result of the micropaleontological research of the Paleogene sediments from Macedonian basins, rich and diverse association of benthic and planktonic foraminiferal fauna has been found. Research collection consists of 68 samples from borehole sections located in Ovče Pole and Tikveš basins and 180 samples of the open Paleogene cross sections in Tikveš, Ovče Pole, Skopje-Kumanovo, Delčevo, Valandovo-Gevgelija and Strumica basins. 76 species belonging to 41 genera and 25 families were established: Spiroplectamminidae Cushman, 1927 – 2 species, Eggerellidae Cushman, 1937 – 1 species, Textulariidae Ehrenberg, 1838 – 2 species, Spiroloculinidae Wiesner, 1920 – 1 species, Hauerinidae Schwager, 1896 – 4 species, Nodosariidae Ehrenberg, 1838 – 5 species, Lagenidae Reuss, 1862 – 4 species, Polymorphinidae d'Orbigny, 1839 – 2 species, Glandulinidae Reuss, 1860 – 2 species, Globigerinidae Carp., Park. and Jones – 12 species, Catapsydracidae Loeblich and Tappan, 1957 – 1 species, Boliviniidae Glaessner, 1937 – 7 species, Buliminidae Jones, 1875 – 3 species, Fursenkoinidae Loeblich and Tappan, 1961 – 1 species, Caucasinidae Bykova, 1959 – 2 species, Stilostomellidae Finlay, 1947 – 2 species, Baggenidae – 2 species, Eponidae

– 2 species, Parrelloidae – 1 species, Cibicidae – 5 species, Nonionidae Schultze, 1854 – 4 species, Chilostomellidae Brandy, 1881 – 1 species, Heterolepidae Gonzáles-Donoso, 1969 – 4 species, Gavelinellidae Hofker, 1956 – 3 species, Rotaliidae Ehrenberg, 1839 – 2 species.

Benthic foraminifers are more taxonomically diverse and represented with a huge number of specimens and sorts that are vastly spread in the basins. Planktonic foraminifers are less in amount, and they show limited spread in Ovče Pole, Tikveš and Skopje-Kumanovo basins.

Benthic and planktonic associations found in Ovče Pole and Tikveš basins, has enabled the process of identifying one biostratigraphic zone *Catapsydrax dissimilis* – *Globigerinatheka tropicalis* (Toumarkine, Luterbacher, 1985) and one biostratigraphic subzone *Bolivina antegressa* from the zone *Planulina costata* (Burgova, 1988), which are significant to the geological age of those basins.

Systematic classification of the foraminiferal fauna was done after Loeblich and Tappan (1988).

Conclusion

Micropaleontological analysis of the Paleogene basins in R. of Macedonia, revealed a wide spectrum of foraminiferal species spread into whole Eocene section. The vast number of the determined foraminiferal taxa is mutual forms for all basins. The mutual foraminiferal fauna in the Paleogene basins in Macedonia refers to similar bionomic conditions and the existence of relation between those reservoirs during the sediment's depositing.

The presence and stratigraphic distribution of many benthic and plankton foraminiferal species from the open Paleogene cross sections in Tikveš, Ovče Pole, Skopje-Kumanovo, Delčevo, Valandovo-Gevgelija and Strumica basins enable us to complete stratigraphy of these basins with new data.

Acknowledgements: The authors are thankful to Assoc. Prof. Dr. Sava Juranov from Sofia University for the consultations for the paper.

References

- Loeblich, A., H. Tappan. 1988. *Foraminiferal Genera and Their Classification*. New York, Van Nostrand Reinhold, 900 p.
- Toumarkine, M., H. Luterbacher. 1985. Paleocene and Eocene planktic foraminifera. – In: Bolli, H., J. B. Saunders, K. Perch-Nielsen (Eds.). *Plankton Stratigraphy*. Cambridge Univ. Press, 87–154.
- Burgova, E. M. 1988. Zonal subdivision of the south USSR on benthic foraminifers. – *Acad. Sci. of the USSR*, 300, 1, 169–171 (in Russian).