


**УНИВЕРСИТЕТ ПО ХРАНИТЕЛНИ ТЕХНОЛОГИИ -
ПЛОВДИВ**

**UNIVERSITY OF FOOD TECHNOLOGIES -
PLOVDIV**



SCIENTIFIC WORKS

Volume LXI

part I

Plovdiv, October 24-25, 2014

НАУЧНА КОНФЕРЕНЦИЯ С МЕЖДУНАРОДНО УЧАСТИЕ

“ХРАНИТЕЛНА НАУКА, ТЕХНИКА И ТЕХНОЛОГИИ 2014”

‘FOOD SCIENCE, ENGINEERING AND TECHNOLOGIES 2014’

НАУЧНИ ТРУДОВЕ

**Том LXI
part I**

Пловдив, 24- 25 октомври 2014



- OPTIMIZATION OF SALT COMPOSITION OF NUTRIENT MEDIUM FOR LIPASE PRODUCTION BY RHIZOPUS ARRHIZUS, Hristina Strinska, Georgi Dobrev, Valentina Dobрева, Boriana Zhekova, Petia Ivanova 491
102. ОПТИМИЗАЦИЯ ПРОЦЕССА КУЛЬТИВИРОВАНИЯ ЧИСТОЙ КУЛЬТУРЫ ДРОЖЖЕЙ, Е.М. Моргунова, Ю.С. Назарова
OPTIMIZATION OF PROCESS OF CULTIVATION OF PURE CULTURE OF YEAST, E.M. Morgynova, Y.S. Nazarova 497
103. КОРЕЛАЦИЯ МЕЖДУ НЯКОИ МОРФОЛОГИЧНИ И БИОХИМИЧНИ СВОЙСТВА НА ОРИЗ (*Oryza sativa* L.), Верица Илиева, Илия Каров, Наталия Маркова Руждик
CORRELATION BETWEEN SOME MORPHOLOGICAL AND BIOCHEMICAL TRAITS OF RICE (*ORYZA SATIVA* L.), Verica Ilieva, Ilija Karov, Natalija Markova Ruzdik 501
104. АЛТЕРНАТИВНИ НА ДЕСТИЛАЦИЯТА МЕТОДИ ЗА ПОЛУЧАВАНЕ НА АЛКОХОЛ, М. Лазарова, Е. Кирова
ALTERNATIVES TO DISTILLATION FOR ALCOHOL RECOVERY, M. Lazarova, E. Kirova 506
105. ИЗСЛЕДВАНЕ ВЛИЯНИЕТО НА ЕСТЕСТВЕНАТА ИЗМЕНЧИВОСТ НА *BACILLUS SUBTILIS* TS 01 ВЪРХУ НЕГОВАТА АНТИБИОТИЧНА АКТИВНОСТ, Севдалина Тодорова, Любка Кожухарова
INVESTIGATION ON THE INFLUENCE OF THE NATURAL VARIABILITY OF *BACILLUS SUBTILIS* TS 01 ON ITS ANTIBIOTIC ACTIVITY, Sevdalina Todorova, Lubka Kozhuharova 510
106. МИКРОБНА ПРОДУКЦИЯ НА ИНУЛИНАЗИ ОТ БАКТЕРИИ ОТ РОД *BACILLUS* И ВЛИЯНИЕ НА ИЗТОЧНИЦИТЕ НА ВЪГЛЕРОД И АЗОТ, Симеон Гавраилов, Камий Перашон, Карол Трюше, Вяра Иванова
MICROBIAL PRODUCTION OF INULINASES FROM BACTERIA OF GENUS *BACILLUS* AND EFFECTS OF NITROGEN AND CARBON SOURCES, Simeon Gavrilov, Camille Perrachon, Carole Truchet, Viara Ivanova 517
107. ИНТЕГРИРАНЕ НА ФЕРМЕНТАЦИЯТА И ПЕРВАПОРАЦИЯТА, М. Лазарова, Е. Кирова
COMBINING FERMENTATION WITH PERVAPORATION, M. Lazarova, E. Kirova 523
108. ПРЕЧИСТВАНЕ НА ОТПАДНИ ВОДИ ОТ 1,2 – ДИБРОМЕТАН, Evgenia Vasileva, Venko Beschkov
WASTEWATER TREATMENT FROM 1,2 – DIBROMOETHANE, Evgenia Vasileva, Venko Beschkov 527
109. ИЗСЛЕДВАНЕ НА СИСТЕМАТА NI-SN ПО МЕТОДА НА ДИФУЗИОННИТЕ ДВОЙКИ, Вая Гандова, Димитър Димитров
DIFFUSION COUPLES STUDIES IN THE NI-SN SYSTEM, Vanya Gandova, Dimitar Dimitrov 533
110. ЕЛЕКТРОХИМИЧНА РЕДУКЦИЯ НА НИТРИТНИ ЙОНИ, Цветомила Първанова-Манчева
ELECTROCHEMICAL REDUCTION OF NITRITE IONS, Tsvetomila Parvanova-Mancheva, Venko Beschkov 538
111. ПРИЛОЖЕНИЕ НА АНТОЦИАНИ ЗА ОЦЕНКА НА МИКРОБИОЛОГИЧНАТА СТАБИЛНОСТ И БЕЗОПАСНОСТ НА КОНДУКТИВНО ЗАГРЯВАНИ ХРАНИ, Цветко Прокопов
APPLICATION OF ANTHOCYANINS FOR ASSESSMENT OF MICROBIOLOGICAL STABILITY AND SAFETY OF FOOD HEATED BY CONDUCTION, Tsvetko Prokopov 542
112. BIOSORPTION OF METHYL ORANGE BY WASTE BIOMASS OF *STREPTOMYCES FRADIAE*, Z. Velkova, G. Kirova, V. Gochev, V. Kafadarova, M. Stoytcheva 546
113. ОБ УСЛОВИЯХ СИНТЕЗА ДИГИДРОФОСФАТА ЦИНКА ТЕТРАГИДРАТА, Антрапцева Н.М., Била Г.Н., Леонова Б.И.
ABOUT CONDITIONS OF SYNTHESIS OF ZINC DIHYDROGENPHOSPHATE TETRAHYDRATE, Antrapitseva N.M., Bila G.N., Leonova B.I. 552
114. О СОСТАВЕ ПРОДУКТОВ ТЕРМООБРАБОТКИ ЦИНК-МАРГАНЕЦ(II) ДИФОСФАТОВ ПЕНТАГИДРАТОВ, Антрапцева Н.М., Солод Н.В., Била Г.Н., Гаврилюк О.В.,



КОРЕЛАЦИЯ МЕЖДУ НЯКОИ МОРФОЛОГИЧНИ И БИОХИМИЧНИ СВОЙСТВА НА ОРИЗ (*Oryza sativa* L.)

Верица Илиева * Илия Каров * Наталија Маркова Руждик *

*Универзитет „Гоце Делчев“, Земјоделски факултет, „Гоце Делчев“ No: 89, 2000 Штип,
Република Македонија, verica.ilieva@ugd.edu.mk, ilija.karov@ugd.edu.mk,
natalija.markova@ugd.edu.mk

CORRELATION BETWEEN SOME MORPHOLOGICAL AND BIOCHEMICAL TRAITS OF RICE (ORYZA SATIVA L.)

Verica Ilieva*, Ilija Karov*, Natalija Markova Ruzdik*

*, „Goce Delcev” University, Faculty of Agriculture, „Goce Delchev” No: 89, 2000 Stip, Republic of
Macedonia, verica.ilieva@ugd.edu.mk, ilija.karov@ugd.edu.mk, natalija.markova@ugd.edu.mk

Abstract

The relationship between some morphological and biochemical traits in ten domestic rice varieties (Kocanski, Osogovka, Biser-2, Nada-115, Ranka, Prima riska, Montessa, N° 51, N° 69 and B 30-303), and three lines of special interest (79/22-2, 78/12-3-4 and 78/12-3-5) were analyzed. The results showed that shape of grain has the highest significantly positive correlation with length of grain ($r=0.897$). Number of grain per panicle is significantly correlated with mass of grains per panicle ($r=0.877$) and fertility with leaf area ($r=0.689$). Proteins content has significantly negative correlation with iron in paddy ($r=-0.862$) and with carbohydrates content in white rice ($r=-0.686$), while iron content with width of grain ($r=-0.719$ in paddy). The derived results can be used as selection index for future rice improvement.

Keywords: rice, morphological traits, biochemical traits, correlation.

Introduction

Rice is a one of the major agricultural crop in the world which is used in nutrition of the population. In the Republic of Macedonia the production of rice is an important industry because fully satisfies the needs of domestic rice and also is an export oriented. The rice production takes a place mainly in the Eastern part of the country, along the river Bregalnica. About 90% of total production is produced in Kocani region and the remaining 10% in Stip, Vinica and Veles region.

Rice is an important source of starch and protein. However, rice scientists have long recognized its micronutrient deficiencies, which are the basis of numerous human health problems worldwide (Welch et al., 2004). Rice is a poor source of essential micronutrients such as Fe and Zn (Bouis, and Welch, 2010, according Anuradha et al., 2012). It only has limited contents of Fe and Zn, and moreover the loss of minerals, particularly of Fe, during rice milling is high (Doesthale et al., 1979, according Tamanna et al., 2013). There is few available

information on the relationship between 8 mineral elements, plant morphological and grain quality traits in Yunnan rice. Therefore, advance in understanding this relationship can provide reference for rice quality breeding and special rice industrialization (Ya-wen et al., 2005).

For selection in rice, information on correlation coefficient always has been helpful as a basis for selection in a breeding programme (Akhtar et al., 2011). The aim of this research was to analyze the relationship between some morphological and biochemical traits in different rice genotypes from Macedonia.

Materials and Methods

Ten rice varieties from Macedonia (No 51, No 69, B-30-303, kocanski, nada-115, ranka, osogovka, biser-2, prima riska and montesa), and three lines of special interest (79/22-2, 78/12-3-4 and 78/12-3-5), were grown on the experimental areas for rice in Kocani. Polish experiments were performed by using



randomized block system in three repetitions. The standard growing measures were applied during the vegetation. The morphological traits (culm height, leaf length, leaf width, panicle length, number of grains in panicle, mass of grains per panicle, 100-grains weight, grain length and grain width) are determined according to the International descriptors for rice (Bioversity International, IRRI and WARDA, 2007; IBPGR and IRRI, 1980). Based on the number of full and empty grains per panicle is calculated the percentage of fertility and from length and width of the leaf is calculated the leaf area. The length and width of the grain, are measured with caliper, and from the ratio of the measured values, was determine the shape of grain. From the biochemical

traits were examined the content of soluble carbohydrates, protein content and content of microelements iron – Fe, zinc – Zn and copper – Cu in categories paddy, cargo and white rice. For getting cargo category, paddy is peeled by hand machine and for getting white rice, bleaching is performed with rice electrical peeling machine. The content of soluble carbohydrates is determined by the method of Dubois et al., 1956, the total nitrogen content is determined by the method of micro Kjeldahl, and the content of microelements is determined by the method of wet combustion and atomic absorption spectroscopy type – PERNIN ELMER 200.

The relationship between examined traits is calculated using software SPSS.

Results and Discussion

The coefficient of correlation between analyzed morphological traits is given in table 1. Significantly positive correlation was obtained between shape and length of grain ($r=0.897$), mass and number of grains per panicle ($r=0.877$) and between fertility and the leaf area ($r=0.689$). The leaf area is positively correlated with the length of leaf ($r=0.611$) but only at significant level $p=0.05$. Positive correlation at significant level, $p=0.05$ exists between grain length and 100 grains weight ($r=0.577$) and between fertility and panicle length ($r=0.569$). The width and length of

leaf showed negative relationship and correlation coefficient ($r=-0.608$) significant only for $p=0.05$.

The results from inter correlation of different morphological traits in rice are presented in researches of Naseem et al., 2014, Maji and Shaibu, 2012, Ashfaq et al., 2012, Zahid et al., 2006. Generally, the panicle length is positively correlated with culm height, the number of grains per panicle with panicle length and leaf area. Direct effect on the yield per plant has the number of grains per panicle and 1000 grains weight.

Table 1. Correlation between some morphological traits of the rice genotypes analyzed

| Traits | Leaf length | Leaf width | Leaf area | Panicle length | Number of grains per panicle | Fertility | Mass of grains per panicle | 100-grains weight | Grain length | Grain width | L/B ratio |
|------------------------------|-------------|------------|-----------|----------------|------------------------------|-----------|----------------------------|-------------------|--------------|-------------|-----------|
| Culm height | 0.447 | -0.238 | 0.295 | 0.366 | 0.014 | 0.094 | -0.041 | -0.118 | 0.328 | 0.419 | 0.105 |
| Leaf length | | -0.608* | 0.611* | 0.408 | -0.014 | 0.227 | -0.217 | -0.472 | 0.114 | 0.074 | 0.077 |
| Leaf width | | | 0.245 | 0.081 | -0.008 | 0.410 | 0.040 | 0.163 | 0.036 | -0.129 | 0.097 |
| Leaf area | | | | 0.519 | -0.078 | 0.689** | -0.276 | -0.432 | 0.149 | 0.419 | 0.177 |
| Panicle length | | | | | 0.521 | 0.569* | 0.356 | -0.180 | 0.050 | 0.339 | -0.115 |
| Number of grains per panicle | | | | | | 0.107 | 0.877** | 0.084 | 0.192 | -0.069 | 0.175 |
| Fertility | | | | | | | 0.102 | -0.007 | 0.073 | 0.194 | -0.020 |
| Mass of grains per panicle | | | | | | | | 0.545 | 0.433 | 0.118 | 0.304 |
| 100-grains weight | | | | | | | | | 0.577* | 0.326 | 0.354 |
| Grain length | | | | | | | | | | -0.030 | 0.897** |
| Grain width | | | | | | | | | | | -0.466 |

Level of significant $p=0.01$ **

Level of significant $p=0.05$ *



The relationship between the analyzed biochemical traits at paddy, cargo and white rice from examined genotypes is given in table 2, 3 and 4 respectively. In paddy (table 2) negative mutual dependence, with high significant was obtained for the content of iron and protein content ($r=-0.862$). Highly significant negative correlation of protein

content was found in the content of carbohydrates in white rice ($r=-0.686$). At other analyzed traits in paddy and white rice was not found significant correlation. In cargo category was not found any significant correlation between examined biochemical traits (table 3).

Table 2. Correlation between some biochemical traits at paddy rice

| Traits | Proteins | Fe | Zn | Cu |
|-----------------------|----------|----------|--------|--------|
| Carbohydrates content | 0.320 | -0.255 | 0.184 | 0.127 |
| Protein content | | -0.862** | 0.103 | -0.095 |
| Fe | | | -0.170 | 0.042 |
| Zn | | | | -0.260 |

Level of significant $p=0.01^{**}$

Table 3. Correlation between some biochemical traits at cargo rice

| Traits | Proteins | Fe | Zn | Cu |
|-----------------------|----------|--------|--------|--------|
| Carbohydrates content | 0.224 | -0.015 | 0.138 | 0.153 |
| Protein content | | -0.317 | 0.080 | -0.096 |
| Fe | | | -0.187 | 0.057 |
| Zn | | | | 0.259 |

Table 4. Correlation between some biochemical traits at white rice

| Traits | Proteins | Fe | Zn | Cu |
|-----------------------|----------|--------|--------|--------|
| Carbohydrates content | -0.686** | -0.132 | -0.098 | 0.041 |
| Protein content | | -0.265 | -0.063 | -0.161 |
| Fe | | | 0.277 | 0.477 |
| Zn | | | | -0.157 |

Level of significant $p=0.01^{**}$

In table 5, 6 and 7 is given the correlation between examined biochemical traits in paddy, cargo and white rice, with examined morphological traits. The content of iron at paddy, cargo and white rice showed

a negative correlation with the grain width. But the correlation is significant for both levels of significance only in paddy.

Table 5. Correlation between some morphological and biochemical traits at paddy rice

| Traits | Carbohydrates content | Protein content | Fe | Zn | Cu |
|------------------------------|-----------------------|-----------------|----------|--------|--------|
| Culm height | 0.491 | 0.501 | -0.549 | 0.488 | -0.001 |
| Leaf length | -0.333 | 0.238 | -0.281 | 0.340 | 0.002 |
| Leaf width | 0.536 | 0.121 | 0.003 | -0.081 | -0.066 |
| Leaf area | 0.119 | 0.383 | -0.346 | 0.371 | 0.007 |
| Panicle length | 0.342 | 0.284 | -0.279 | 0.228 | -0.394 |
| Number of grains per panicle | 0.359 | -0.290 | 0.304 | 0.110 | -0.178 |
| Fertility | 0.272 | 0.569* | -0.562* | 0.009 | 0.025 |
| Mass of grains per panicle | 0.354 | -0.174 | 0.096 | 0.095 | -0.318 |
| 100-grains weight | 0.159 | 0.093 | -0.271 | -0.015 | -0.344 |
| Grain length | 0.226 | 0.208 | -0.286 | 0.401 | -0.422 |
| Grain width | 0.274 | 0.508 | -0.719** | 0.189 | -0.086 |
| L/B ratio | 0.0746 | -0.032 | 0.056 | 0.286 | -0.325 |

Level of significant $p=0.01^{**}$, Level of significant $p=0.05^{*}$



At cargo category and white rice obtained values are significant only at $p=0.05$. In paddy, the content of iron is negatively correlated with fertility ($r=-0.562$) but only at the level of significance at $p=0.05$. Fertility showed significant correlation with the protein content in paddy ($r=0.569$) and in the category cargo ($r=0.636$) but only at level of significance $p=0.05$. The content of zinc in white rice is negative significant correlation with the panicle length ($r=-0.749$) for both levels of significance and with trait

fertility ($r=-0.590$) only at level of significance $p=0.05$.

According to Ekka et al., 2011, Sellappan et al., 2009, the grain size determines the quantity of important microelements, such as iron and zinc in grain. In researches of Ekka et al., 2012, grain dimensions are not in significant correlation with content of iron and zinc. Negative correlation between the iron content and grain dimensions and zinc content and dimensions of grains show the researches of Yawen et al., 2005

Table 6. Correlation between some morphological and biochemical traits at cargo rice

| Traits | Carbohydrates content | Protein content | Fe | Zn | Cu |
|------------------------------|-----------------------|-----------------|---------|--------|--------|
| Culm height | 0.473 | 0.470 | -0.461 | 0.217 | 0.029 |
| Leaf length | -0.325 | 0.319 | -0.526 | 0.204 | -0.058 |
| Leaf width | 0.402 | 0.084 | 0.371 | 0.144 | 0.2701 |
| Leaf area | 0.016 | 0.447 | -0.297 | 0.412 | 0.242 |
| Panicle length | 0.016 | 0.630* | -0.112 | 0.123 | -0.126 |
| Number of grains per panicle | -0.008 | 0.085 | 0.440 | 0.455 | -0.167 |
| Fertility | 0.176 | 0.636* | -0.332 | 0.151 | 0.162 |
| Mass of grains per panicle | 0.154 | -0.020 | 0.274 | 0.237 | -0.398 |
| 100-grains weight | 0.415 | -0.217 | -0.133 | -0.248 | -0.491 |
| Grain length | 0.328 | -0.068 | -0.178 | 0.330 | -0.507 |
| Grain width | 0.198 | 0.314 | -0.668* | -0.340 | -0.108 |
| L/B ratio | 0.193 | -0.212 | 0.124 | 0.448 | -0.386 |

Level of significant $p=0.05^*$

Table 7. Correlation between some morphological and biochemical traits at white rice

| Traits | Carbohydrates content | Protein content | Fe | Zn | Cu |
|------------------------------|-----------------------|-----------------|---------|----------|--------|
| Culm height | -0.003 | 0.129 | -0.548 | -0.068 | -0.014 |
| Leaf length | -0.175 | 0.105 | -0.302 | -0.031 | 0.223 |
| Leaf width | 0.069 | -0.107 | 0.050 | -0.335 | -0.041 |
| Leaf area | -0.102 | -0.005 | -0.351 | -0.364 | 0.211 |
| Panicle length | -0.155 | 0.032 | -0.109 | -0.749** | 0.352 |
| Number of grains per panicle | -0.389 | -0.004 | 0.539 | -0.109 | 0.490 |
| Fertility | 0.077 | -0.086 | -0.453 | -0.590* | 0.140 |
| Mass of grains per panicle | -0.187 | -0.094 | 0.404 | -0.011 | 0.149 |
| 100-grains weight | 0.364 | -0.252 | -0.022 | 0.188 | -0.478 |
| Grain length | -0.060 | -0.136 | -0.091 | 0.360 | -0.396 |
| Grain width | 0.182 | 0.308 | -0.580* | -0.414 | -0.247 |
| L/B ratio | -0.140 | -0.239 | 0.161 | 0.503 | -0.253 |

Level of significant $p=0.01^{**}$

Level of significant $p=0.05^*$



Conclusion

The analysis of correlation between some morphological and biochemical traits in thirteen genotypes of rice from Republic of Macedonia showed that the shape of the grain is significantly positive correlation with the length of grain and mass of grains per panicle with the number of grain per

panicle. Fertility is in positively correlated with the leaf area. A negative correlation was found between the protein content and content of iron in paddy, and content of carbohydrates in white rice. The content of iron in paddy, cargo and white rice is negatively correlated with the grain width.

References

- [1] Agahi, K., Fotokian, M., Farshadfar, E., (2007). Correlation and path coefficient analysis for some Yield – Related Traits in Rice Genotypes (*Oryza sativa* L.). Asian Journal of Plant Sciences 6 (3): 513-517.
- [2] Akhtar, N., Nazir, M., Rabnawaz, A., Mahmood, T., Safdar, M., Asif, M., Rehman, A., (2011). Estimation of heritability, correlation coefficient analysis in fine grain rice (*Oryza sativa* L.). The Journal of Animal & Plant Sciences, 21(4), 660-664.
- [3] Anuradha, K., Agarwal, S., Batchu, A., Babu, A., Swamy, B., Longvah, T., Sarla, N., (2012). Evaluating rice germplasm for iron and zinc concentration in brown rice and seed dimensions. Journal of Phytology, 4(1): 19-25.
- [4] Ashfaq, M., Khan, S., Ahmad, R., (2012). Association of various morphological traits with yield and genetic divergence in rice (*Oryza sativa* L.). Int. J. Agric. Biol., 14: 55-62.
- [5] Bioversity International, IRRI and WARDA. (2007). Descriptors for wild and cultivated rice (*Oryza* spp.). Bioversity International, Rome, Italy; International Rice Research Institute, Los Banos, Philippines; WARDA, Africa Rice Center, Cotonou, Benin.
- [6] Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A., & Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 33, 1574-1579.
- [7] Ekka R., Sarawgi A., Kanwar R. (2011). Correlation and Path Analysis in Traditional Rice Accessions of Chhattisgarh. Journal of Rice Research, Vol. 4 No 1 & 2, 11-18.
- [8] International Board for Plant Genetic Resources and International Rice Research Institute (1980). Descriptors for rice *Oryza sativa* L. By IBPGR - IRRI Rice Advisory Committee, Manila, Philippines.
- [9] Maji, A., Shaibu, A., (2012). Application of principal component analysis for rice germplasm characterization and evaluation. Journal of Plant Breeding and Crop Science, Vol. 4(6), pp. 87-93.
- [10] Naseem, I., Khan, A., Akhter, M., (2014). Correlation and path coefficient studies of some yield related traits in rice (*Oryza sativa* L.). International Journal of Scientific and Research Publications. Volume 4, Issue 4, 1-5.
- [11] Tamanna, S., Parvin, S., Kumar, S., Dutta, A., Ferdoushi, A., Siddiquee, M., Biswas, S., Howlader, M., (2013). Content of some minerals and their bioavailability in selected popular rice varieties from Bangladesh. International Journal of Current Microbiology and Applied Sciences, Vol. 2, No 7, 35-43.
- [12] Welch, R. M., Graham, R. D., (2004). Breeding for micronutrients in staple food crops from a human nutrition perspective. *J. Exp. Bot.*, 55, 353–364.
- [13] Ya-wen, Z., Shi-quan, S., Lu-xiang, W., Jia-fu, L., Xiao-yuing, P., Juan, D., Min, Q., (2005). Correlation of Plant Morphological and Grain Quality Traits with Mineral Element Contents in Yunnan Rice. Rice Science, 12(2): 101-106.
- [14] Zahid, M., Akhter, M., Sabar, M., Manzoor, S., Awan, T., (2006). Correlation and Path Analysis Studies of Yield and Economic Traits in Basmati Rice (*Oryza sativa* L.). Asian Journal of Plant Sciences 5 (4): 643-645.