

8th CMAPSEEC

Conference on Medicinal and Aromatic Plants of Southeast European Countries

organized by

- Association for Medicinal and Aromatic Plants of Southeast European Countries (AMAPSEEC)
- Albanian Academy of Science



PROCEEDINGS

May 19-22, 2014
Durrës, Albania

Title: **Proceedings of the Eighth Conference on Medicinal and Aromatic Plants of Southeast European Countries**
(Proceedings of the 8th CMAPSEEC)

Editors: Prof. Dr. Zora Dajić Stevanović
Assoc. Prof. Dr. Alban Ibraliu

All papers within the Proceedings of the 8th CMAPSEEC were reviewed by the members of the Scientific Committee and kind assistance of some members of other Conference bodies. Reviewed papers were categorized as review papers, original scientific papers and additional papers.

Reviewers: Prof. Dr. Zora Dajić Stevanović
Prof. Dr. Dragoja Radanovic
Prof. Dr. Milen Georgiev
Prof. Dr. Zlatko Šatovic
PhD. Asmund Asdal
Prof. Dr. Dea Baričević
Prof. Dr. Danijela Stešević
Prof. Dr. Efigjeni Kongjika
Prof. Dr. Ilkay Orhan
Prof. Dr. Nazim Sekeroglu
Prof. Dr. Biljana Bauer Petrovska
Assoc. Prof. Dr. Diamanto
Prof. Dr. Ivan Salamon
Prof. Dr. Jelena Cvejic
Prof. Dr. Monica Hancianu
AsSoc. Prof. Dr. Sonia Melino
Assoc. Prof. Dr. Vilma Papajani
Assoc. Prof. Dr. Gjosh Stefkov

Published by: Academy of Science of Albania, Agricultural University of Tirana and Association for Medicinal and Aromatic Plants of Southeast European Countries (AMAPSEEC)

Technical Support: Alban Ruci

ISBN: **978-99956-10-66-1**

**8th Conference on Medicinal and Aromatic Plants of
Southeast European Countries
(8th CMAPSEEC)**

May 19-22, 2014, Adriatik Hotel, Durrës, Albania

Organized by:

**Association for Medicinal and Aromatic Plants of Southeast European Countries
(AMAPSEEC)**

Albanian Academy of Science

Agricultural University of Tirana

CONFERENCE COMMITTEES

HONORARY BOARD

Prof. Dr. Fatos Harizaj
(Agricultural University of Tirana, Rector)

Prof. Dr. Salvatore Bushati
(Scientific Secretary of Natural &
Technical Section, Academy of Sciences,
Albania)

Prof. Dr. Edmond Panariti
(Minister of Agriculture, Rural
Development and Water Management)

Prof. Dr. Nazim Gruda
(Agricultural University of Tirana and
University of Bonn, Germany)

Prof. Dr. Uran Asllani
(Representative of Albanian MAP
Industry)

Prof. Dr. Zef Sima
(Faculty of Pharmacy, Medicine
University, Albania)

CHAIRPERSON

Assoc. Prof. Dr. Alban Ibraliu
(President of 8th CMAPSEEC)

HONORARY CHAIRPERSON

Prof. Dr. Zora Dajic-Stevanovic
(President of AMAPSEEC)

ORGANIZING COMMITTEE MEMBERS

Assoc. Prof. Dr. Alban Ibraliu

Dr. Dejan Pljevljakusic

Assoc. Prof. Dr. Julian Shehu

Assoc. Prof. Dr. Fetah Elezi

Prof. Dr. Sokol Abazi

Assoc. Prof. Dr. Ndoc Faslia

SCIENTIFIC COMMITTEE

Assoc. Prof. Dr. Alban Ibraliu - Albania

Prof. Dr. Alfred Mullaj - Albania

Prof. Dr. Andrea Pieroni - Italy

Prof. Dr. Ardian Maci - Albania

PhD. Asmund Asdal - Norway
Prof. Dr. Biljana Bauer Petrovska - FYR
Macedonia
Prof. Dr. Danijela Stešević - Montenegro
Prof. Dr. Dea Baricevic - Slovenia
Assoc. Prof. Dr. Diamanto Lazari - Greece
Prof. Dr. Dragoja Radanovic - Serbia
Acad. Ass. Prof. Dr. Biology Efigjeni
Kongjika - Albania
Assoc. Prof. Dr. Gjoshe Stefkov - FYR
Macedonia
Prof. Dr. Ilkay Orhan - Turkey
Prof. Dr. Ivan Salamon - Slovakia
Prof. Dr. Jelena Cvejic - Serbia

Prof. Dr. Joachim Müller - Germany
Prof. Dr. Johannes Novak - Austria
Dr. Kiril Tkachenko - Russia
Prof. Dr. Milen Georgiev - Bulgaria
Prof. Dr. Monica Hancianu - Romania
Prof. Dr. Muhammad Iqbal - India
Prof. Dr. Nazim Sekeroglu - Turkey
Asoc. Prof. Dr. Sonia Melino - Italy
Prof. Dr. Stella Kokkini - Greece
Assoc. Prof. Dr. Vilma Papajani - Albania
Prof. Dr. Zlatko Šatovic - Croatia
Prof. Dr. Zora Dajic Stevanovic - Serbia

CONFERENCE'S SECRETARY

Najada Kadiasi

Rea Tako

Forward

Medicinal and aromatic plants (MAP) have been essential resources for human health from ancient times to the present day. The majority of the world's population depends on traditional medicine for primary health care needs. More than 35.000 plant species are used in herbal medicine and as spices, out of the most are of a local importance due to traditional use. Because of their increasing appliance in pharmaceutical, food, cosmetic and beverage industry, as well as use in folk and official medicine, veterinary and plant protection, herbal industry has been recognized as an important element of global economy. Together with growth in global demand for medicinal plants and in local demand for plant based traditional medicines, the pressure on the existing populations of medicinal plants has increased tremendously during the last few decades. The extinction or scarcity of these plants is not only a problem for conservation – it also results in serious problems for people's health and livelihoods. Cultivation may reduce harvesting pressure on some wild species, particularly rare and threatened species, and thus can also be an important production strategy that supports conservation. South East Europe is particularly appreciated for richness in indigenous MAP resources and long tradition in use of MAP and their products. The region is known as one of the main suppliers of MAP raw material into EU and US. In addition, medicinal plants were being the subject of a great scientific interest in the SEE region, where significant contribution to understanding of various research aspects in number of MAP species was achieved by Conferences on Medicinal and Aromatic Plants of Southeast European Countries, under organization and support of the Association for Medicinal and Aromatic Plants of Southeast European Countries (AMAPSEEC) established in 2000 by the Institute for Medicinal Plant Research "Dr Josif Pančić", Belgrade, R. of Serbia. We are very proud that, the 8th Conference on MAP of SEE countries took place in Durres, Albania, after the second was held in Greece, the third in Slovakia, followed by Romania, Czech Republic, Turkey and Serbia. The 8th CMAPSEEC gathers together not only the researchers from SEE region, but from all over the world. At the Conference participated over 320 researchers, experts, company's representatives and guests interested in MAP diversity, biology, conservation, ecology, phytochemistry, pharmacology, breeding, cultivation and biotechnology. Near 283 summaries of research contributions were presented, out of about 100 in the form of full papers. All contributions covered very different research areas, and were classified into the three distinct groups: "MAP diversity at all levels and tools for its evaluation", "Pharmacology and biological effects of active MAP compounds" and "Map cultivation, breeding and biotechnology".

The full papers, categorized as review papers and original scientific papers, within this proceedings were reviewed by our Scientific Committee. In addition, a small group of papers was issued either without referring on reviewer's comments. However, we decided to include them as well, considering the subjects and research approaches interesting. We strongly believe that presented results will contribute to a general knowledge on MAP, and will encourage both young researchers and processing companies to deal with many species whose composition and biological effects were appointed as promising. Moreover, we hope that pleasant and collegial atmosphere additionally contributed to establishing of the new professional and personal contacts and to strengthening of the ones already established. Finally, strong network on scientists and professionals interested in MAP under AMAPSEEC umbrella might bring new project ideas and new value in the near future.

Editors, Prof. Dr. Zora Dajić Stevanović and Assoc. Prof. Dr. Alban Ibraliu

Foreword from the Minister of Education and Sports
Mrs. Lindita Nikolla

Ladies and Gentleman

Please Allow me to initially thank the organizers of this special event, who made possible this important conference regarding scientific research in Albania.

I also wish to express my congratulations to all the participants in the conference and to the authors of the research work to be presented today, especially our foreign guests.

Today's conference is important, because scientific research is not only an academic achievement, but also in that research and innovation determine economic growth, create new job opportunities and establish the progress of a country.

The government of the Republic of Albania, mandated in September 2013, has immediately initiated concrete measures to be taken in the field of research, while considering scientific research as a compass which can direct towards the most successful economic sectors. With an edict from the prime minister Mr. Edi Rama, an independent commission was established for the reformation of higher education and scientific research in Albania.

After several months of intensive work the commission has designed the first draft of the report on the reform, which is being discussed these days with all the interest groups and actors involved. After being finalized this report will serve as the base document for the new higher education law.

The reform will provide concrete steps toward the quality improvement of scientific research in Albania. An essential element for this improvement is the development of technical capacities and project managing, but foremost is the international cooperation at the academic level. For this reason, the ministry of education and sports has undertaken the steps for the association of Albania in the EU scientific research program, Horizon 2020. This program offers good opportunities for the development of scientific research at the regional and global level.

Our government by being a part of the Horizon 2020 program and also Erasmus Plus, has created the opportunities that favor the internationalization of higher education and scientific research, promotion of student exchange, young researchers and academic staff. In these programs it has been clearly stated that maximal support will be offered for the cooperation between native and foreign researchers.

Also the ministry of education and sport will support scientific research programs of quality and innovation, with structural and program financing.

Scientific research transforms every challenge into an opportunity, therefore your contribution is needed in protecting our biodiversity and promoting the application of biotechnology in creating new varieties of cultivated crops.

Honored participants, I believe that the endeavours and collaborations in the field of science, research and innovation are one of the better ways in strengthening our regional partnership and advancing our national and regional development

Thank you for your contribution, and I wish you further success in the proceeding of the conference



8th Conference on Medicinal and Aromatic Plants of Southeast European Countries

May 19-22, 2014, Adriatik Hotel, Durres, Albania



SUPPORTING INSTITUTIONS




Ministry of Education and Sports of the Republic of Albania

SILVER SPONSOR



DONORS



SALVIANORD

TABLE OF CONTENTS

Section I - "MAP diversity at all levels and tools for its evaluation"	12
ARE THERE STILL NEGLECTED MEDICINAL PLANTS BEYOND OFFICIAL AND TRADITIONAL CONSIDERATION?	
<i>Dajic Stevanovic Zora¹, Pedja Janackovic², Milan Stankovic³</i>	13
GENETIC RESOURCES OF MEDICINAL AND AROMATIC PLANTS OF ALBANIA – CURRENT STATUS OF THE NATIONAL COLLECTION OF MAPS	
<i>Ibraliu Alban¹, Mullaj Alfred², Elezi Fetah³, Shehu Julian¹ and Gixhari Belu³</i>	23
MEDICINAL AND AROMATIC PLANTS (MAP) IN VASCULAR FLORA OF THE DRENICA MOUNTAIN-REPUBLIC OF KOSOVO	
<i>Krasniqi Elez & Krasniqi Hazbije</i>	32
STATUS OF WILD-GROWING MEDICINAL PLANTS IN THE ANINEI MOUNTAINS (WESTERN ROMANIA): RECENT DATA	
<i>Antal S. Diana, Ardelean Florina</i>	41
GEOGRAPHIC DISTRIBUTION AND DIVERSITY ASSESSMENT IN EX SITU COLLECTION OF ALBANIAN MEDICINAL PLANTS	
<i>Gixhari Belu¹, Hobdari Valbona¹, Kadiasi Najada², Faslja Ndoc², Ibraliu Alban²</i>	51
GEOGRAPHIC DISTRIBUTION AND SPATIAL GAPS ASSESSMENT IN EX SITU COLLECTION OF <i>ORIGANUM VULGARE</i> L. STORED IN ALBANIAN GENEBA	
<i>Gixhari Belu¹, Hobdari Valbona¹, Kadiasi Najada², Faslja Ndoc²</i>	59
MEDICINAL MUSHROOMS AND THERAPY: TRANSLATING A TRADITIONAL PRACTICE INTO THE WESTERN MEDICINE	
<i>¹Bauer Biljana, ²Karadelev Mitko</i>	67
USE OF ST JOHN'S WORT THROUGH THE AGES	
<i>Bauer Biljana¹, Kostic Vesna²</i>	76
MICROMORPHOLOGICAL RESEARCH REGARDING THE GLANDULAR HAIRS OF <i>THYMUS PRAECOX</i> OPIZ SSP. <i>POLYTRICHUS</i> (A. KERN. EX BORBAS) JALAS	
<i>Boz Irina¹, Toma Constantin², Zamfirache Maria Magdalena², Gille Elvira^{3*}</i>	83
DISTRIBUTION OF <i>SIDERITIS RAESERI</i> BOISS. ET HELDR. IN ALBANIA – STATE OF ITS POPULATIONS AND RECOMMENDATIONS FOR CONSERVATION	
<i>Aneva Y. Ina¹, Evstatieva N. Luba¹, Zhelev Peter², Papajani - Toska Vilma³, Ibraliu Alban⁴</i>	89
FLORA AND VEGETATION OF BERATI CASTLE IN ALBANIA	
<i>Shehu Julian¹, Imeri Alma¹, Mullaj Alfred²</i>	101
ON THE REPRODUCTIVE BIOLOGY OF <i>SIDERITIS SYRIACA</i> L. (LAMIACEAE)	
<i>Yankova-Tsvetkova Petrova Elina, Aneva Ina</i>	112

ETHNOBOTANICAL STUDY OF MEDICINAL PLANTS TRADITIONALLY USED IN FIERI DISTRICT, ALBANIA.	
<i>Papajani Vilma¹, Ibraliu Alban² Miraçi Mirela¹, Rustemi Anjeza¹</i>	123
RESEARCH AND DEVELOPMENT OF THE PLANT MEDICINE BLOSSOMS IN SLOVAKIA – NEW VARIETIES	
<i>Salamon Ivan</i>	134
TURKISH ANISE (PIMPINELLA ANISUM L.)	
<i>AVCI Ayse Betül¹, GIACHINO AKCALI R. Refika²</i>	140
Section II - "Pharmacology and biological effects of active MAP compounds"	146
ANTI-NOCICEPTIVE, ANTI-INFLAMMATORY AND ANTIPYRETIC EFFECTS OF MORINGA OLEIFERA PLANT	
<i>Ghazal Nabil, M. Atef, EL Banna, H. A.</i>	147
ANTIOXIDANT CAPACITY AND PHENOLIC CONTENT OF SALVIA L.	159
<i>¹ Muráriková Andrea*, ¹ Neugebauerová Jarmila, ¹ Kaffková Katarína, ² Raab Simona</i>	159
COULD MEDICINAL PLANTS OFFER ALTERNATIVES TO IMPROVE THE HUMAN IRON POOL?	
<i>Antal S. Diana¹, Ardelean Florina, Dehelean A. Cristina</i>	164
FATTY ACID COMPONENTS OF SOME ENDEMIC SIDERITIS GÜMÜŞÇÜ, Ahmet¹ AKBULUT, Mehmet² GÜMÜŞÇÜ, Gönül³	171
THE CONTENT OF SATURATED, MONOUNSATURATED AND POLYUNSATURATED FATTY ACIDS IN THE SEEDS OF DIFFERENT CANOLA VARIETIES	
<i>Bauer Biljana¹ Kostić Vesna²</i>	176
VOLATILE COMPOUNDS AND ANTIOXIDANT ACTIVITY OF PEEL ESSENTIAL OILS AND METHANOLIC EXTRACTS OF FOUR GREEK CITRUS SPECIES: CITRUS BERGAMIA, CITRUS MEDICA, CITRUS AURANTIUM AND FORTUNELLA JAPONICA	
<i>Sarrou Eirini¹, Chatzopoulou Paschalina², Dimasi-Theriou Kortessa¹, Therios Ioannis¹</i>	182
EVALUATION OF THE ANTIFUNGAL ACTIVITIES OF MACEDONIAN WILD MUSHROOM EXTRACTS AGAINST SELECTED FUNGAL STRAINS	
<i>Ivanova Emilija¹, Atanasova-Pancevska Natalija¹, Karadelev Mitko¹, Bogdanov Jane², Kungulovski Dzoko¹</i>	193
CHEMICAL COMPOSITION AND ANTIFUNGAL ACTIVITY OF ESSENTIAL OILS FROM ALBANIAN MEDICINAL PLANTS.	
<i>Mamoci Erjon, Hodaj Entela, Hasalliu Rozeta</i>	201
PROPOLIS PROFILING OF SAMPLES FROM SOUTHWESTERN ROMANIA (LUGOJ REGION): NEW DATA	
<i>Ardelean Florina¹, Dragos Dan¹, Szabadai Zoltan¹, Antal S. Diana¹</i>	208

DETERMINATION OF FUMONISINS AND BEAUVERICIN IN ANIS SEED BY LIQUID CHROMATOGRAPHY/TANDEM MASS SPECTROMETRY

Vukovic LJ. Gorica¹, Aleksić A. Goran², Kuzmanović T. Slobodan², Starović S. Mira², Vljaković M. Jelena¹, Vojislava P. Bursić³

214

HOW DO LYCOPENE AND ANTIOXIDATIVE ACTIVITY VARY IN TWO TOMATO GENOTYPES UNDER DEFICIT IRRIGATION TREATMENTS

Pećinar, M. Ilinka¹, Rančić, V. Dragana¹, Pekić Quarrie, V. Sofija¹, Milosavić, B. Nenad², Dajić Stevanović, P. Zora¹, Bertin, Nadia³ and Stikić, I. Radmila¹

221

QUANTITATIVE ANALYSIS OF GLYCYRRHIZIC ACID OF GLYCYRRHIZA L. TAXA FROM THE CZECH REPUBLIC AND KYRGYZSTAN

¹NEUGEBAUEROVÁ JARMILA, ²SAZYKULOVA J. GULBAIRA

227

DETERMINATION OF MACRO NUTRIENT CONTENT IN SOME HERBAL DRUGS FROM THE BLACK SEA PROVINCES IN TURKEY

KORKMAZ Kürşat¹, KARA Şevket Metin², AKGÜN Mehmet¹, BATI Ebru³

235

ESSENTIAL OIL OF COMMON JUNIPER (*JUNIPERUS COMMUNIS* L.) IN ALBANIA

Salamon Ivan¹, Ibraliu Alban², Fejer Jozef¹

239

ESSENTIAL MACRO NUTRIENT PROFILES OF SELECTED MEDICINAL AND AROMATIC PLANTS FROM THE FAMILY OF LAMIACEAE

Kara Şevket Metin¹, ÖzkutluFaruk², Açıkgoz M. Akif¹, Batı Ebru³

245

CORRELATION BETWEEN ANTIOXIDATIVE POTENTIAL OF PURE CAPSAICIN AND CAPSICUM OLEORESINS

Maksimova Viktorija¹, Koleva G. Liljana², Ruskovska Tatjana¹, Cvetanovska Ana³, Gulaboski Rubin¹

251

TOXICOLOGICAL EVALUATION OF *JUNIPERUS* SPECIES FROM FLORA OF THE R. MACEDONIA

Hamidi R. Mentor¹, Jovanova Blagica¹, Karapandzova Marija², Stefkov Gjoshë², Cvetkovikj Ivana², Kulevanova Svetlana², Kadifkova Panovska Tatjana¹

257

SEED PROGENY OF PORTUGUESE FENNEL WILD POPULATIONS: MORPHOLOGICAL AND ESSENTIAL OILS VARIABILITY

Lopes VR¹, Barata AM¹, Rocha F¹, Bettencourt E², Mota AS³, Silva L.³, Figueiredo AC⁴

265

ANTIOXIDANT ACTIVITY, TOTAL POLYPHENOLS CONTENT AND FLAVONOIDS CONTENT OF AQUEOUS EXTRACTS OF SOME ALBANIAN MEDICINAL PLANTS

Aruci (Neza) Edlira¹, Neza Jonida²

276

PHYTOCHEMISTRY OF THE ESSENTIAL OIL OF *MELLISA OFFICINALIS* L. GROWING WILD IN MOROCCO: PREVENTIVE APPROACH AGAINST NOSOCOMIAL INFECTION

JALAL Zineb, ERRAI Siham, LYOUSSI Badiia and ABDELLAOUI Abdelfattah

283

Section III - "MAP Cultivation, Breeding and Biotechnology"

288

MOLECULAR AND BIOCHEMICAL CHARACTERIZATION, AND <i>IN VITRO</i> CONSERVATION OF SOME ALBANIAN POPULATIONS OF SAGE (<i>SALVIA OFFICINALIS</i> L.)	
<i>Kongjika Efigjen¹, Bacu Ariola², Babani Fatbardha², Sota Valbona², Ricciardi Luigi³</i>	289
EFFECT OF NITROGEN FERTILIZATION UPON MOUNTAIN SAVORY YIELD AND ESSENTIAL OIL AND ITS STABILITY ESTIMATION	
<i>Crnobarac¹ Jovan, Adamović² Dušan, Danojević² Dario, Jaćimović¹ Goran</i>	300
VALUE CHAIN ANALYSIS OF MEDICINAL PLANTS FROM BERATI REGION IN CENTRAL ALBANIA	
<i>Qose Anisa¹, Tabaku Vath², Toromani Elvin², Mine Luljeta³</i>	306
EVALUATION OF HOW BEESWAX OF ALBANIAN ORIGIN AFFECTS THE SPF OF SUN CREAM	
<i>MYFTARI Brunilda¹, JUCA Besnik¹, MALAJ Ledjan¹, TOSKA Vilma¹, MYFTARI Elton²</i>	315
PHYTOPLASMA DISEASE OF MEDICINAL PLANTS IN SERBIA	
<i>Pavlović, D. Snežana¹, Stojanović, D. Saša², Jošić, Lj. Dragana³, Starović, S. Mira²</i>	321
FUNGI ASSOCIATED WITH CARAWAY FRUIT IN SERBIA	
<i>Stojanović D. Saša¹, Pavlović DJ. Snežana², Aćimović G. Milica⁴, Aleksić A. Goran¹, Kuzmanović T. Slobodan¹, Jošić LJ. Dragana³</i>	330
NAPHTODIANTHRONE PRODUCTION IN <i>HYPERICUM PERFORATUM</i> L. TRANSGENIC SHOOTS	
<i>Tusevski Oliver¹, Trajkovska Ljubica¹, Ivanova Lozenka¹, Shijakova Kristiana¹, Petreska Stanoeva Jasmina², Stefova Marina², Gadzovska Simic Sonja¹</i>	335
<i>IN VITRO</i> CULTURE OF MATURE ZYGOTIC POMEGRANATE EMBRYOS (<i>PUNICA GRANATUM</i> L.)	
<i>Sota Valbona¹, Kongjika Efigjen²</i>	345
MID-TERM <i>IN VITRO</i> CONSERVATION OF MYRTLE (<i>MYRTUS COMMUNIS</i> L.) – A VALUABLE MEDICINAL PLANT	
<i>Sota Valbona¹, Kongjika Efigjen²</i>	353
ESTABLISHMENT OF <i>GLORIOSA SUPERBA</i> CELL SUSPENSION CULTURES	
<i>Zarev Yancho¹, Ionkova Iliana¹</i>	362
Additional Papers	369
MEDICINAL AND AROMATIC PLANTS FROM THE STUDENICA REGION- REPUBLIC OF KOSOVO	
<i>Ukaj Shkëlzim^{1*}, Millaku Fadil², Shala Albana¹, Sallaku Fatbardh³</i>	370
TESTING OF SOME ECOTYPES OF SAGE FOR PRODUCTIVITY AND ACTIVE PRINCIPLES QUALITY.	
<i>Mato Arqilea¹, Mero Gjergji², Hajkola Kostandin³</i>	376
SAFFRON (<i>CROCUS SATIVUS</i> L.) - A NEW AROMATIC AND MEDICINAL PLANT AND ITS CULTIVATION	

<i>KUTROLLI Florenc¹</i>	383
<hr/>	
INCREASED EXPORT POSSIBILITIES, A NEW TREND FOR ORGANIC CULTIVATION OF MEDICINAL AND AROMATIC PLANTS	
<i>LAMA Enilda¹; KUTROLLI Florenc²; MIHO Liri³, IBRALIU Engjellushe⁴</i>	391
<hr/>	
CHEMICAL ANALYSIS OF HYDROLATES OF <i>SATUREJA MONTANA</i> PRODUCED FROM STEAM DISTILLATION INDUSTRY IN ALBANIA	
<i>BUÇI Aurora¹, CIKO Lorena¹, CELA Dorisa¹, ÇELIBASHI Lule¹, ABAZI Sokol¹</i>	400
<hr/>	
ANTIOXIDANT ACTIVITY OF THE METHANOL AND ETHANOL EXTRACTS OF ENDEMIC <i>THYMUS MALYI</i> RONNINGER	
<i>Marin A. Marija¹, Novaković M. Miroslav²</i>	405
<hr/>	
<i>HALACSYA SENDTNERI</i> (BOISS.) DÖRFL. - ANTIOXIDANT ACTIVITY OF THE METHANOL EXTRACT	
<i>Branković R. Snežana¹, Marin A. Marija²</i>	408
<hr/>	
EFFECT OF DIRECT SELECTION ON PRODUCTIVE TRAITS OF MARSHMALLOW (<i>ALTHAEA OFFICINALIS</i> L.)*	
<i>Slobodan B. Dražić</i>	411
<hr/>	

THE CONTENT OF SATURATED, MONOUNSATURATED AND POLYUNSATURATED FATTY ACIDS IN THE SEEDS OF DIFFERENT CANOLA VARIETIES

Bauer Biljana¹ Kostic Vesna²

¹Institute of Pharmacognosy, Faculty of Pharmacy, SS. Cyril and Methodius University,
Majka Tereza No 47, 1000 Skopje, Republic of Macedonia

²Institute of Public Health of Republic of Macedonia, 50 Divizija 6,
1000 Skopje, Republic of Macedonia

ABSTRACT

Canola is a name applied to edible oilseed rape. This plant belongs to the *Brassicaceae* (mustard) family along with 3,000 other species. Close relatives of this crop have been cultivated for food since the earliest recordings of man. The name "canola" was registered in 1979 by the Western Canadian Oilseed Crushers Association to describe "double-low" varieties. Double low indicates that the processed oil contains less than 2% erucic acid (C_{22:1}). Like soybean, canola contains both high oil content as well as high protein content. It contains about 40% oil and 23% protein compared to 20 and 40%, respectively, for soybean consumption.

Commercial varieties of canola were developed from three species: *Brassica napus* L. (Argentine type), *Brassica campestris* L. (Polish type) and *Brassica juncea* L. (canola quality brown mustard). There are considerable differences in agronomic characteristics, yield, and fatty acid (FA) composition of seed oil between species and between varieties.

The main objective of this work was identification and determination the FA composition of the seed oil of the two canola varieties grown in the Republic of Macedonia, during 2012. For that purpose, a total of hundred samples of the seeds of the two types of canola varieties were analyzed for the presence of total saturated fatty acid (SFA), total monounsaturated fatty acids (MUFA), and total polyunsaturated fatty acids (PUFA).

The results of the study, showed different FA content among the two canola varieties. The canola variety type 2, was found to be high linolenic with the average content of linolenic acid (C_{18:3}) 44.0% ± 2.02. The canola variety type 1, was found to be high oleic with the average content of oleic acid (C_{18:1}) 59.5% ± 1.91. The average content of erucic acid (C_{22:1}) was below 0.2% in the both varieties.

The canola variety type 1 contained lower mean value of the total SFA (9.6% ± 0.56) in comparison with canola variety type 2, which had higher mean value of the total SFA (17.4% ± 0.67). The canola variety type 1, had higher content of the total MUFA unlike the canola variety type 2. The differences in the FA composition, as well as, the total SFA, MUFA, and

the PUFA content, the both canola varieties had similar values of polyunsaturated/saturated indexes (P/S), which were found to be 3.2, and 3.4, respectively. This means that the both varieties of the oil had the same nutritional value.

Keywords: *Canola, fatty acid, gas chromatography, polyunsaturated / saturated index*

INTRODUCTION

Canola was developed in the early 1970s using traditional plant breeding techniques by Canadian plant breeders to remove the anti-nutritional components (erucic acid and glucosinolates) from rapeseed to assure its safety for human and animal consumption. The canola plant also produced seeds with a very low level of saturated fat, seven percent or below [1]. There is an internationally regulated definition of canola that differentiates it from rapeseed, based upon its having less than two percent erucic acid (C_{22:1}) and less than 30 µmol/g glucosinolates [2]. Oilseed products that do not meet this standard cannot use the trademarked term "Canola." Like soybean, canola contains both high oil content as well as high protein content. It contains about 40% oil and 23% protein compared to 20 and 40%, respectively, for soybean consumption [3].

Commercial varieties of canola were developed from three species: *Brassica napus* L. (Argentine type), *Brassica campestris* L. (Polish type) and *Brassica juncea* L. (canola quality brown mustard). There are considerable differences in agronomic characteristics, yield, and fatty acid (FA) composition of seed oil between species and between varieties. In order to develop herbicide tolerance of the canola plant, and to improve quality of the canola seed, some different innovations have been established. *Roundup Ready* and *Liberty Link* canola varieties were developed using a traditional plant breeding technique called mutagenesis. Another innovation is the development of hybrid canola varieties. Hybrids can increase yields and are increasing in acreage [4].

In the Republic of Macedonia, two different hybrids of canola varieties have been developed in the past decade. The main objective of our work was identification and determination the FA composition of the seed oil of the two canola varieties, which were grown in the Republic of Macedonia, during 2012. For that purpose, a total of hundred samples of the seeds of the two types of canola varieties were analyzed for the presence of total saturated fatty acid (SFA), total monounsaturated fatty acids (MUFA), and total polyunsaturated fatty acids (PUFA). The values of polyunsaturated / saturated indexes (P/S) were calculated for the both canola varieties.

MATERIAL AND METHODS

Materials

A total of hundred seed samples of two different hybrids of canola varieties were collected from the local producers during the 2012 (canola variety type 1, n=45; canola variety type 2, n=55).

Methods

Sample preparation

The total of 5 g of grinded samples was mixed with 5 g of anhydrous sodium sulphate (Na_2SO_4). An aliquot of 2 g was placed in Soxlet extractor system, and extraction was performed with petroleum – ether (40-69) within 24 hours. After cooling at room temperature, the solvent was evaporated to dryness in the stream of nitrogen.

Preparation of fatty acid methyl esters (FAMES)

Fatty acid (FA) composition of the oils and fats was determined as their corresponding methyl esters. Preparation of FAMES was carried out according to the modified ISO method (5). 0.1 - 0.2 g of oil was dissolved in 10 mL 0.2 mol/L H_2SO_4 prepared in anhydrous methanol. Esterification was performed by refluxing for 30 minutes at 100 °C in tightly sealed Pyrex tubes. After cooling at room temperature, 10 mL of petroleum ether (40 - 60) was added followed by 10 mL of deionized water, mixed gently and allowed to settle until the upper petroleum ether layer becomes clear. The distinct upper layer of methyl esters in petroleum ether was separated carefully in a capped vial and used for analysis. 2 μL of the petroleum ether aliquots were injected into the chromatographic column and peaks were recorded for their respective retention times and areas by the data processor unit of the GC. Identification of each individual fatty acid methyl ester was achieved by comparison with authentic reference standards. All solvents and standards were of analytical grade (Merck, Fluka).

Chromatography

HP model 5890 series II (plus) gas chromatograph equipped with an HP automatic liquid sampler and a flame-ionization detector (FID) was used either with a nonpolar fused silica capillary column (30 m x 0.32 mm id. x 1 μm film thickness) coated with 100% poly (dimethylsiloxane), commercially available as SPB^{TM-1} obtained from Supelco (USA). The carrier gas (nitrogen) flow rate was 1.5 mL/min and the split ratio was 1:10. The injection port was maintained at 250 °C and the FID at 280 °C. Oven temperature was set at 200 °C (1 minute) increasing for 5 °C/min. The final oven temperature was maintained at 250 °C (20 minutes). For confirmation of identified and determined FAMES in oils and fats, a polyethylene glycol TPA modified polar column commercially available as HP-FFAP (25 m x 0.32 mm id x 0.52 μm) was used with the same HP model 5890 series II (plus) gas chromatograph. The carrier gas (nitrogen) flow rate was 1.5 mL/ min and the split ratio was 1:10. The injection port was maintained at 230 °C and the FID at 260 °C. Oven temperature was set at 180 °C increasing for 2 °C/ min. The final oven temperature was maintained at 230 °C (4 minutes).

RESULTS AND DISCUSSION

A total of 100 seed samples of two different hybrids of canola varieties were analyzed on the composition of fatty acids using gas chromatographic method. The content of following saturated and unsaturated fatty acids was tested in the samples: caproic acid ($\text{C}_{6:0}$), caprylic acid ($\text{C}_{8:0}$), capric acid ($\text{C}_{10:0}$), lauric acid ($\text{C}_{12:0}$), myristic acid ($\text{C}_{14:0}$), palmitic acid ($\text{C}_{16:0}$), stearic acid ($\text{C}_{18:0}$), arachidic acid ($\text{C}_{20:0}$), behenic acid ($\text{C}_{22:0}$), lignoceric acid ($\text{C}_{24:0}$), oleic acid ($\text{C}_{18:1}$), linoleic ($\text{C}_{18:2}$), linolenic acid ($\text{C}_{18:3}$), and erucic acid ($\text{C}_{22:1}$). The fatty acid percent composition of tested seeds is shown in Table 1 and Table 2, respectively. The mean of total saturated fatty acid (SFA), monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA) and the values of polyunsaturated/saturated indexes (P/S) are shown in Table 3.

The results of the determination of FA composition (Table 1) indicate that the seeds of both canola varieties have low content of saturated FA with the predominant presence of C_{16:0} and C_{18:0}. The level of C_{18:0} in the seeds of the canola variety type 1 was 4.4%±1.4 of the total FA, and 6.9%±1.6 of the total FA in seeds of canola variety type 2. Although it is classified as a SFA, data accumulated during the past years indicate that C_{18:0} is unique among the SFAs in the food supply [6]. Unlike other predominant long-chain SFA: C_{16:0}, C_{14:0}, and C_{12:0} which increase blood cholesterol levels, C_{18:0} has been shown to have a neutral effect on blood total and low density lipoprotein (LDL) cholesterol levels [6,7].

Table 1. Saturated fatty acid composition of canola seeds of two varieties

Type of Canola variety	Mean ±SD									
	C _{6:0} (%)	C _{8:0} (%)	C _{10:0} (%)	C _{12:0} (%)	C _{14:0} (%)	C _{16:0} (%)	C _{18:0} (%)	C _{20:0} (%)	C _{22:0} (%)	C _{24:0} (%)
Canola type 1 (n=45)	<0.1	<0.1	<0.1	<0.1	<0.1	5.2±0.6	4.4±1.4	<0.2	<0.2	<0.1
Canola type 2 (n=55)	<0.1	<0.1	<0.1	<0.1	<0.1	10.5±2.5	6.9±1.6	<0.2	<0.2	<0.1

The obtained results for the unsaturated FA (Table 2) composition of the seed oil, showed the predominant presence of C_{18:1} in the seed oil of canola variety type 1 which was found to be 59.5% ± 1.91 of the total FA. It is known that the consumption of C_{18:1} is effective in lowering LDL cholesterol level. The predominant presence of C_{18:3} (44.0% ± 2.02) was found in the seed oil of canola variety type 2. One of the most interesting yet controversial dietary approaches has been the possible prophylactic role of dietary γ -linolenic acid (GLA) in treating various chronic diseases states. This strategy is based on the ability of diet to modify cellular lipid composition and eicosanoid (cyclooxygenase and lipoxygenase) biosynthesis [8].

C_{22:1} is a MUFA which is a major constituent of certain oils, such as rapeseed oil. Because it has been linked to cardiac muscle damage, oil such as canola oil that are low in C_{22:1} was developed. In our studies, the seeds of the both canola varieties had low content of C_{22:1}.

Table 2. Unsaturated fatty acid composition of canola seeds of two varieties

Type of Canola variety	Mean ±SD			
	C _{18:1} (%)	C _{18:2} (%)	C _{18:3} (%)	C _{22:1} (%)
Canola type 1 (n=45)	59.5 ± 1.91	18.8 ± 3.5	11.9 ± 1.1	0.11±0.05

Canola type 2 (n=55)	23.2 ± 2.9	15.2 ± 3.6	44.0 ± 2.02	0.18±0.09
----------------------	------------	------------	-------------	-----------

All samples presented total SFA content less than one fourth of the total FA content (Table 3). It is known that the excessive consumption of SFA is related to the increase of the plasmatic cholesterol and the obesity [9]. On the other hand, the consumption of PUFA and MUFA has been recommended to improve the lipid profile in relation to the saturated SFA. Yu-Poth *et al.* indicate that the rich diets in PUFA may provoke an increase in the LDL -cholesterol oxidation and the reduction of the HDL -cholesterol levels [10]. There is a tendency in increasing the recommendations of MUFA consumption, that seems not to affect the HDL levels, and also it may reduce the LDL and triacylglycerols blood levels, that make it more effective in prevention of hearth diseases. Canola oil variety 1 showed high content of MUFA (59.5 ± 1.91) and canola oil variety 2 showed hight content of PUFA (59.2 ± 1.1).

Table 3. The content of SFA, MUFA, PUFA and the values of P/S indexes in two types of canola varieties

Type of Canola variety	Mean ±SD			P/S index
	SFA (%)	MUFA (%)	PUFA (%)	
Canola type 1 (n=45)	9.6 ± 0.56	59.5 ± 1.91	30.7 ± 1.7	3.2
Canola type 2 (n=55)	17.4 ± 0.67	23.2 ± 2.9	59.2 ± 1.1	3.4

The relationship between saturated and polyunsaturated FA content is expressed as P/S index. This value is an important parameter for determination of nutritional value of certain oil. Oils and fats with higher value of P/S index than 1 are considered to have nutritional value. Several studies indicate that higher value of P/S index means a smaller deposition of lipids in the body [11]. The results of our investigations (Table 3) showed that the both canola varieties had similar values of polyunsaturated/saturated indexes (P/S), which were found to be 3.2, and 3.4, respectively. This means that the both varieties of the oil had the same nutritional value.

CONCLUSIONS

The canola plant was developed in order to produce edible oil that contains much lower levels of erucic acid (C_{22:1}) than rapeseed oil. Canola oil derived from the seeds of the canola plant, the genetically engineered rapeseed plant or the canola hybrids, has been considered for human consumption. The types of fatty acids determine whether a vegetable oil is used for edible or industrial purposes.

In our investigations the FA composition of the seed oil obtained from two types of canola hybrids was analyzed. The canola variety type 2, was found to be high linolenic with the average content of linolenic acid (C_{18:3}) of 44.0% ± 2.02. The canola variety type 1, was found to be high oleic with the average content of oleic acid (C_{18:1}) 59.5% ± 1.91. The average content of (C_{22:1}) was below 0.2% in the both types.

The canola variety type 1 contained lower mean value of the total SFA in comparison with canola variety type 2, which had higher mean value of the total SFA. The canola variety type 1, had higher content of the total MUFA unlike the canola variety type 2. Besides the differences in the FA composition, as well as, the total SFA, MUFA, and the PUFA content, the both canola varieties had similar values of polyunsaturated/saturated indexes (P/S). This means that the both varieties of the oil had the same nutritional value.

REFERENCES

1. J.E. BRANDLE, P.B.E. Mc VETTY (1989): *Effects of inbreeding and estimates of additive genetic variance within seven summer oilseed rape cultivars*. Genome, vol. 32, pp. 115-119.
2. Z.P. KONDRA, B.R. STEFANSSON (1970): *Inheritance of the major glucosinolates of rapeseed (Brassica napus) meal*. Can. J. Plant Sci., vol. 50, pp. 643-647.
3. S. GOWERS (1980): *The production of hybrid oilseed rape using self incompatibility*. Cruciferae. Newsletter, vol. 5, pp. 15-16.
4. G.C. BUZZA (1995): *Plant Breeding. Brassica Oilseeds: Production and Utilization*. Edited by D.S. Kimber and D.I. McGregor. Cab International, pp. 153-175.
5. BS EN ISO 5508 (1995). *Animals and vegetables fats and oils. Analysis by gas chromatography of methyl esters of fatty acids*.
6. B.F. HAUMANN (1998): *Stearic acid: a 'different' saturated fatty acid*. INFORM (American Oil Chemists' Society), vol. 9(3), pp. 202-208.
7. R.P. MENSINK (2005): *Effects of stearic acid on plasma lipid and lipoproteins in humans*. Lipids, vol. 40, pp. 1201-1205.
8. Y. Y. FAN, R.S. CHAPKIN (1998): *Importance of Dietary γ -Linolenic Acid in Human Health and Nutrition*. J. Nutr., vol 128 (9), pp. 1411-1414.
9. V. RISTIC, G. RISTIC (2003): *Role and importance of dietary polyunsaturated fatty acids in the prevention and therapy of atherosclerosis*. Med. Pregled, vol. 56 (1-2), pp.50 - 53.
10. S. YU-POTH., T. D. ETHERTON, C. C. REDDY, T.A., PEARSON, R. REED., G. ZHAO.(2000): *Lowering dietary saturated fat and total fat reduces the oxidative susceptibility of LDL in healthy men and women*. Journal of Nutrition, vol. 130 (9), pp. 2228 - 2237.
11. C. L. LAWTON., H. J. DELALGRY, J. BROCKMAN, R.C. SMITH, J. E. BLUNDELL (2000): *The degree of saturation of fatty acids influences in post ingestive satiety*. British Journal of Nutrition, vol. 83 (5), pp. 473 - 482.