

Comparison of Dissolution Test for Metronidazole Tablets in International and American Pharmacopoeia

(poster presentation)

Violeta Koneva¹, *Biljana Gjorgjeska*², *Sofija Petkovska*³

¹ Student, University "Goce Delcev" Stip, Faculty of Medical Science, Krste Misirkov bb, P.fax 201, 2000 Stip, Republic of Macedonia

² Full Professor, University "Goce Delcev" Stip, Faculty of Medical Science, Krste Misirkov bb, P.fax 201, 2000 Stip, Republic of Macedonia

³ Assistant, University "Goce Delcev" Stip, Faculty of Medical Science, Krste Misirkov bb, P.fax 201, 2000 Stip, Republic of Macedonia

+389 32 550 434, e-mail: biljana.gorgeska@ugd.edu.mk

Abstract

The tablets are commonly used dosage forms and as such are subject to tests to prove their quality. As inevitable here is the test that determine dissolution of active component from the tablet. Over the years the test suffered a series of improvements. At the end it become an official test in pharmacopoeias and mandatory testing for tablets quality. Depending on the properties of the active ingredient, pharmaceutical form and effect of the drug used specific standardized equipment, appropriate medium, and method of analysis is proscribed in pharmacopoeia monographs.

The purpose of this paper was to make a comparison of the tests for metronidazole tablets in official monographs in International Pharmacopoeia and US Pharmacopoeia in order to determine similarities and differences.

Comparison of monographs in both pharmacopoeias was made in terms of apparatus used, the medium, the speed of rotation of the apparatus and the tolerances allowed when performing the test.

Although the dissolution test prescribed in general part of both pharmacopoeias is the same, in the individual monographs the visible differences exist in terms of the prescribed equipment, medium, speed and tolerance as shown in Table 1.

Table 1 – Dissolution testing for metronidazole tablets, comparison of monographs in International pharmacopoeia and US Pharmacopoeia

	International pharmacopoeia	US Pharmacopoeia
Apparatus	paddle	basket
Medium	500 ml buffer pH= 6,8	900 ml 0,1N HCl
Rotation speed	75 rpm	100 rpm
Sampling time	30 min	60 min
Tolerances	not less then 80% of labeled amount	not less then 85% of labeled amount

It can be concluded that the dissolution test of active component in metronidazole tablets is an important parameter for testing the quality, despite the differences that exist in performing the test in various pharmacopoeias. Using the dissolution test in routine control of the production process proves compliance with the required quality of each batch produced. This is particularly important in terms of ensuring the quality of medicines and meeting the standards of good manufacturing practice.

Keywords: apparatus, dissolution test, International Pharmacopoeia, European Pharmacopoeia, US Pharmacopoeia

1. EUROPEAN PHARMACOPOEIA 5.0, 2.9.3. Dissolution test for solid dosage forms, 01/2005:20903 pp. 228
2. The International Pharmacopoeia - Fifth Edition, 2015 Metronidazole tablets (Metronidazoli compressi)
3. USP 29-NF 24, Metronidazole tablets, page1426
4. Faisal Asif, Tania Sultana, Md. Didaruzzaman Sohel , Md. Helal Uddin Sumon, Md. Hassan Kawsar, Md. Ashrafur Islam, *In Vitro* Dissolution Pattern of Metronidazole Film Coated Tablet in Presence of Fruit Juice, *American Journal of Pharmacological Sciences*. 2014, 2(2), 32-36
5. Amber Nawab, Samina Alam, Nousheen Shoukat, IN VITRO DISSOLUTION STUDY OF METRONIDAZOLE FILM COATED TABET 400 MG (FLAGYL) IN THE PRESENCE OF DIFFERENT TEA, *WORLD JOURNAL OF PHARMACY AND PHARMACEUTICAL SCIENCES*, 2015, Volume 4, Issue 12, 102-11