

МАКЕДОНСКО ГЕОЛОШКО ДРУШТВО

ТРЕТ КОНГРЕС

на

Геолозите на Република Македонија

ЗБОРНИК НА ТРУДОВИ

-КНИГА 2-



Уредници:

Лепиткова, С. & Боев, Б.

Струга, 2016

*Посебно издание на
Geologica Macedonica, № 4*

МАКЕДОНСКО ГЕОЛОШКО ДРУШТВО

**ТРЕТ КОНГРЕС
на
Геолозите на Република Македонија**

ЗБОРНИК НА ТРУДОВИ

-КНИГА 2-

Уредници:
Лепиткова, С. & Боев, Б.

Струга, 2016

Издавач: Македонско геолошко друштво

Главни и одговорни уредници: Проф. д-р Соња Лепиткова и
Проф. д-р Блажо Боев

Уреднички одбор: Проф. д-р Тодор Серафимовски (Р.Македонија), Проф. д-р Блажо Боев (Р.Македонија), Acad. Prof. Vladimir Bermanec PhD (Croatia), Акад. проф д-р Владица Цветковиќ (Србија), Acad. prof. Ivan Zagorchev PhD (Bulgaria), Prof. Tadej Dolenec PhD (Slovenia), Prof. David Alderton PhD (Great Britain), Prof. Wolfgang Todt PhD (Germany), Акад. проф. д-р Николај С. Бортников (Русија), Prof. Clark Burchfield PhD (USA), Prof. Thierry Auge PhD (France), Проф. д-р Тодор Делипетров (Р.Македонија), Проф. д-р Милорад Јовановски (Р.Македонија), Проф. д-р Споменко Михајловиќ (Србија), Проф. д-р Драган Миловановиќ (Србија), Проф. д-р Дејан Прелевиќ (Germany), Prof. Albrecht von Quadt (Switzerland) PhD.

Технички уредник: Доц. д-р Игор Пешевски

Печати: Печатница "2-ри Август С" -Штип

Тираж: 300 примероци

Организационен одбор на Третиот Конгрес на Геолозите на Република Македонија

Претседател: Проф. д-р Соња Лепиткова
Секретар: д-р Златко Илијовски

Технички секретар: Доц. д-р Игор Пешевски

Членови: Проф. д-р Блажо Боев
Проф. д-р Тодор Серафимовски
Проф. д-р Милорад Јовановски
Проф. д-р Орце Спасовски
Проф. д-р Војо Мирчовски
д-р Коста Јованов
м-р Флорент Чиче
Кирил Филев

Финансиска поддршка:

ДПТУ „Бучим“ ДОО-Радовиш
АДОРА ИНЖЕНЕРИНГ ДООЕЛ – Скопје
Рудник “САСА” ДООЕЛ – Македонска Каменица
Градежен Институт „Македонија“ АД – Скопје
ГЕИНГ Кребс унд Кифер Интернешнл и др. ДОО – Скопје
„Мермерен комбинат“ АД – Прилеп
Простор ДОО – Куманово
„Геохидроконсалтинг“ ДООЕЛ – Скопје
„Геохидроинженеринг“ ДООЕЛ – Скопје
Хидроинженеринг ДООЕЛ– Битола
Градежен факултет – Скопје, Катедра за геотехника
„ГЕОМАП“ ДОО – Скопје
БУЛМАК ГРУП ДООЕЛ – Скопје
ЕУРОМАКС РЕСОУРЦЕС ДОО – Скопје
САРДИЧ МЦ ДООЕЛ – Скопје
МАРКОВСКИ КОМПАНИ БОРЧЕ ДООЕЛ – Битола
DIWI Македонија ДООЕЛ – Скопје
ВАРДАРГРАДБА ДОО – Скопје

КНИГА 2
СОДРЖИНА
CONTENTS

2. Фундаментална Геологија

PETROLOGICAL CHARACTERISTICS OF CLASTIC SEDIMENTARY ROCKS IN SV. BARBARA MINE IN RUDE NEAR SAMOBOR, SAMOBORSKA GORA MT. <i>Šime Bilić, Vesnica Garašić</i>	369
PETROLOGY OF GNEISSES FROM THE VRŠAC MOUNTAINS <i>Dragan Milovanović, Danica Srećković-Batočanin, Emin Memović</i>	371
МИНЕРАЛОШКО – ПЕТРОГРАФСКИ И ХЕМИСКИ КАРАКТЕРИСТИКИ НА ГРАНИТОИДНИТЕ КАРПИ ОД ЛОКАЛИТЕТОТ ПЕШТАНИ, ЗАПАДНА МАКЕДОНИЈА <i>Сашо Стојков, Даниел Спасовски, Орце Спасовски</i>	373
МИНЕРАЛОШКО-ПЕТРОГРАФСКИ И ГЕОХЕМИСКИ КАРАКТЕРИСТИКИ НА СКАРНОВИТЕ ОД НАОЃАЛИШТЕТО ИБЕРЛИ <i>Кика Шпритова</i>	379
GARNET-ANDALUSITE/SILLIMANITE-BIOTITE XENOLITHS FROM THE DACITE OF SLAVKOVICA (LJIG, SW SERBIA) <i>Nada Vasković, Danica Srećković – Batočanin, Suzana Erić, Vesna Matović</i>	389
BLADED QUARTZ TEXTURE AND ITS RELATIONSHIP TO ELECTRUM MINERALIZATION IN THE EOCENE, LOW-SULFIDATION KUKLITSA GOLD DEPOSIT, SE BULGARIA PRELIMINARY DATA <i>Irina Marinova, Elena Tacheva</i>	393
ASSOCIATION OF OXIDE MINERALS – CONCENTRATORS OF CHALCOPHILE ELEMENTS (Pb, Zn, Sb) FROM THE “MIXED SERIES” NEAR NEŽILOVO VILLAGE, MACEDONIA <i>Simeon Jancev, Nikita V. Chukanov, Vera N. Ermolaeva</i>	401
LORANDITE AND ORPIMENTE FROM EDIT-25 NORTH PART OF ALSHAR DEPOSIT <i>Ivan Boev, Blažo Boev</i>	405
NEW INVESTIGATIONS ON DUNJE PEGMATITE, MACEDONIA I: THE CONTRIBUTION TO THE KNOWLEDGE ABOUT ALKALI FELDSPARS AND ITS MINERAL PARAGENESIS <i>Vladimir Zebec, Snježana Mikulčić Pavlaković, Marin Šoufek, Blažo Boev, Ivan Boev, Vladimir Bermanec</i>	413

NEW INVESTIGATIONS ON DUNJE PEGMATITE, MACEDONIA II: RELATION TO HOST METAMORPHIC ROCKS AND ADJACENT GRANITE INTRUSIONS	
<i>Nenad Tomašić, Andrea Čobić, Blažo Boev, Ivan Boev, Vladimir Bermanec</i>	417
MOISSANITE METEORITE IN TERRITORY OF THE REPUBLIC OF MACEDONIA	
<i>Blažo Boev, Velo Markovski, Ivan Boev</i>	421
МИНЕРАЛОШКО-ПЕТРОЛОШКИ КАРАКТЕРИСТИКИ НА МИКАШИСТИТЕ ОД ОКОЛИНАТА НА С.БОНЧЕ, ПРИЛЕП	
<i>Филип Јованоски, Тена Шијакова-Иванова, Блажо Боев, Виолета Стефанова</i>	425
ИДЕНТИФИКАЦИЈА НА МИНЕРАЛИТЕ ОД НАОЃАЛИШТЕТО ЗА ОЛОВО И ЦИНК "ЗЛЕТОВО" СО ПРИМЕНА НА РЕНДГЕНСКО ДИФРАКЦИОНА МЕТОДА	
<i>Елена Наунова, Тена Шијакова-Иванова, Блажо Боев</i>	431
GEOCHEMICAL FEATURES OF SILURIAN - DEVONIAN SECTION OF PELAGONIANE ZONE IN ALBANIA	
<i>Irakli Prifti, Ilir Alliu, Agim Ymeri</i>	437
GEOMICROBIOLOGICAL OBSERVATION IN MAJDANSKA REKA, ALLCHAR, MACEDONIA	
<i>Vladimir Bermanec, Jasna Hrenović, Željka Fiket, Ladislav Palinkaš, Ivan Boev, Blažo Boev</i>	447
RADIONUCLIDES IN SOIL, MOSSES, AND MUSHROOM OF THE PRAŠNIK RAINFOREST (CROATIA)	
<i>Gordana Mednuić, Gordana Marović, Jasminka Senčar</i>	451
SCANNING ELECTRON MICROSCOPY STUDIES OF PARTICLES (PM-10) FROM THE TOWN OF KAVADARCI AND VILAGE VOZARCI , REPUBLIC OF MACEDONIA	
<i>Ivan Boev, Sonja Lepitkova, Blažo Boev</i>	453
КВАЛИТЕТ НА АМБИЕНТАЛЕН ВОЗДУХ- СУСПЕНДИРАНИ ЧЕСТИЧКИ (ПМ-10) ВО ОБЛАСТА ТИКВЕШ	
<i>Иван Боев, Дејан Мираковски, Марија Хаџи Николова, Блажо Боев</i>	459
GEO THERMOBAROMETRIC INVESTIGATIONS OF HERCYNIAN GRANITOIDS OF EAST SERBIA	
<i>Dragana Bosić, Suzana Erić, Kristina Šarić, Bojan Kostić, Vladica Cvetković, Dragan Jovanović</i>	467

MIGRATION OF MICROELEMENTS B, NI, MO, AS, V IN COAL FORMATION SIBOVČ FIELD OF KOSOVO BASIN <i>Agim Ymeri, Çerçiz Durmishi, Irakli Prifti, Adil Januzi</i>	469
СЕИЗМОТЕКТОНСКИ ЗОНИ И СЕИЗМИЧКИ ХАЗАРД ВО РЕПУБЛИКА МАКЕДОНИЈА <i>Никола Думуриџанов, Зоран Милутиновиќ, Радмила Шалиќ</i>	477
MAP OF THE MOHO DISCONTINUITY OF THE REPUBLIC OF MACEDONIA <i>Todor Delipetrov, Krsto Blazev, Blagica Doneva, Risto Popovski</i>	493
ТЕКТОНСКА РЕОНИЗАЦИЈА И СЕИЗМИЧНОСТ НА РЕПУБЛИКА МАКЕДОНИЈА <i>Благица Донева, Тодор Делипетров, Ѓорѓи Димов, Зоран Панов, Радмила К. Стефановска</i>	497
ПРОЕКТ ЗА ИЗРАБОТКА НА ДИГИТАЛНИ ГЕОФИЗИЧКИ КАРТИ НА РЕПУБЛИКА МАКЕДОНИЈА ВО ГИС ТЕХНОЛОГИЈА СО ТОЛКУВАЧИ <i>Новица Столиќ, Ивица Андов</i>	503
НОВИ СОЗНАНИЈА ЗА ГЕОМАГНЕТНОТО ПОЛЕ НА РЕПУБЛИКА МАКЕДОНИЈА <i>Марјан Делипетрев, Владимир Маневски, Крсто Блажев</i>	513
МЕТОД НА КОМБИНАЦИЈА НА ГЕО – ЕЛЕКТРИЧНО СОНДИРАЊЕ И КАРТИРАЊЕ <i>Владимир Маневски, Марјан Делипетрев, Ивица Коцев, Благој Делипетрев</i>	521
KOENIGSBERGER RATIO AND TOTAL MAGNETIC FIELD ANOMALY REDUCTION TO THE POLE FOR THE AREA OF MACEDONIA <i>Vesna Cvetkov, Dragana Đurić, Vesna Lesić, Miroslav Starčević, Mirko Petković, Snežana Petrović</i>	529
2D GEOPHYSICAL MODELS OF DEMIR KARIJA ORHOLITE COMPLEX <i>Dragana Đurić, Vesna Cvetkov, Ivana Vasiljević, Spomenko Mihajlović, Vladica Cvetković</i>	535
ГРАВИМЕТРИСКИ МРЕЖИ НА РЕПУБЛИКА МАКЕДОНИЈА <i>Новица Столиќ, Мирослав Старчевић, Сашо Димески</i>	539
COMPARING GEOMAGNETIC FIELD DAILY ANOMALIES AND GEOSPATIAL SEISMICITY AND ATMOSPHERE DATA IN BALKAN COUNTRIES DURING THE <i>BLACKSEAHAZNET PROJECT</i> <i>Milena Cukavac, Strachimir Cht. Mavrodiev, Lazo Pekevski, Spomenko J. Mihajlovic</i>	547

APPLICATION OF IP/RESISTIVITY “REAL SECTION” TECHNIQUE IN SEARCH FOR SULPHIDE MINERALIZATION IN SERBO – MACEDONIAN MASSIF, KOSOVO <i>Përparim Alikaj, Altin Karriqi, Erjon Çollaku</i>	553
ORE MINERALIZATION AT STUDIED GEOMAGNETIC ANOMALIES ON MT. GOLIIJA <i>Jovan Kovačević, Boris Vakanjac, Nikolić Dušan, Mihajlo Pandurov</i>	559
РЕГИСТРАЦИЈА НА ШУМАНОВ РЕЗОНАНС (ПЛ. ПЛАЧКОВИЦА) <i>Лазо Пекевски, Ристо Поповски, Зоран Панов, Страшимир Маєродиев</i>	569

3. Геологија и Економија

STRUCTURAL-METALLOGENIC MAP OF THE REPUBLIC OF MACEDONIA: PRINCIPLES AND CRITERIA <i>Todor Serafimovski, Alexandr Volkov, Goran Tasev</i>	573
THE AU/AG RATIO IN EPITHERMAL DEPOSITS <i>Alexander Volkov, Irina Chizhova, Anatoly Sidorov</i>	581
NEW DATA ON THE CENOZOIC VOLCANISM AND ORE MINERALIZATIONS IN THE PETROSHNITSA RIVER VALLEY, NW PART OF THE KRATOVO- ZLETOVO VOLCANIC AREA, REPUBLIC OF MACEDONIA <i>Slavcho Ivanov Mankov, Manol Stoyanov Antonov, Dmytro Rostislavovich Siroshstan, Valentin Yordanov Grozdev</i>	589
CHARACTERISTICS OF CHROMITE MINERALIZATION ON MT. JELICA <i>Predrag Mijatović, Boris Vakanjac, Dragan Jovanović, Božidar Luković</i>	597
3D MODELING OF SOME COPPER DEPOSITS IN THE REPUBLIC OF MACEDONIA <i>Todor Serafimovski, Christos Christidis, Dalibor Serafimovski, Goran Tasev, Mitko Ligovski, Igor Ivanovski, Lazar Gjorgjiev</i>	605
THE VRSHNIK ORE BODY A POSSITIVE EXAMPLE FOR EXPLORATION, EXPLOITATION AND FILLING IN THE BUCHIM COPPER MINE, EASTERN MACEDONIA <i>Kiril Filev, Todor Serafimovski, Lazar Gjorgjiev, Goran Tasev, Mite Mitev, Metodi Stojanov</i>	613
THE OCCURRENCE OF IRON MINERALIZATION IN VICINITY OF OSTENJAK (ARANĐELOVAC), SERBIA <i>Nemanja Pantelić, Bojan Kostić, Predrag Vulić</i>	619

OVERVIEW OF THE NATURAL PARAMETERS FROM THE GEOLOGICAL-ECONOMICAL EVALUATION OF THE BOROV DOL ORE DEPOSIT, REPUBLIC OF MACEDONIA <i>Lazar Gjorgjiev, Todor Serafimovski, Kiril Filev, Goran Tasev</i>	621
COPPER ORECLASTS OF OLISTOSTROME ORIGIN AT BOR, SERBIA <i>Ivan Antonijević</i>	629
РЕЗУЛТАТИ ОД ПРЕЛИМИНАРНИТЕ ПРОСПЕКЦИСКИ ИСПИТУВАЊА НА ЗЛАТО ВО АЛУВИОНОТ НА ПЕКЉАНСКА РЕКА, ВИНИЦА <i>Виолета Стефанова, Марин Александров, Тодор Серафимовски, Горан Тасев, Војо Мирчовски</i>	635
SUPERGENE PROCESSES IN THE COPPER MINERALISATION AT THE KRALJIČIN ZDENAC ON THE MEDVEDNICA MT. <i>Ladislav A. Palinkaš, Danijela Šmajgl, Andreja Čobić, Vladimir Bermanec</i>	643
THE CRVEN DOL ARSENIC-THALIUM MINERALIZATION IN ALSAR DEPOST IN THE REPUBLIC OF MACEDONIA <i>Ivan Boev, Blazo Boev, Sonja Lepitkova</i>	649
PREBAIKAL FORMATION ON THE TERRITORY OF THE REPUBLIC OF MACEDONIA AS SIGNIFICANT BEARERS OF QUARTZ RAW MATERIALS <i>KrstoBlazev, Gorgi Dimov, Blagica Doneva, Marjan Delipetrov</i>	659
GENESIS OF OIL IN THE DEEPEST MIOCENE SOURCE ROCKS IN THE NORTH-WEST PART OF SAVA DEPRESSION <i>Snježana Blažeković Smojić, Vesna Hrženjak, Darko Tomašić, Tamara Troskot-Čorbić, Marina Mužina</i>	665
КВАЛИТАТИВНИ КАРАКТЕРИСТКИ НА АМФИБОЛСКИТЕ ШКРИЛЦИ ОД ЛОКАЛИТЕТОТ “ПОЧИВАЛО“ ИСТОЧНА МАКЕДОНИЈА КАКО ОСНОВА ЗА НИВНА УПОТРЕБА КАКО АРХИТЕКТОНСКО – ГРАДЕЖЕН КАМЕН <i>Орце Спасовски, Даниел Спасовски</i>	675
DIMENSIONAL STONE DEPOSITS IN WESTERN MACEDONIA <i>Ljupche Kulakov</i>	681
МОЖНОСТИ ЗА КОРИСТЕЊЕ НА ОНИКСОТ И ТРАВЕРТИНОТ ОД ЛОКАЛИТЕТОТ ДЕКОВА ДАБИЦА КАКО АРХИТЕКТОНСКИ КАМЕН <i>Орце Спасовски, Даниел Спасовски</i>	689

ТЕКТОНСКАТА ПОВРЗАНОСТ НА РАСЕДНАТА ЗОНА ЖИВОЈНО -БРОД ГНЕОТИНО- СУВОДОЛ <i>Љупчо Петрески, Марија Манева, Анита Мартиновиќ</i>	695
ГЕОЛОШКИ РЕЗЕРВИ НА ЈАГЛЕН ВО ПЕ “РУДНИЦИ”- РЕК БИТОЛА <i>Љупчо Петрески, Елизабета Митревска, Марија Манева</i>	701
ЕНЕРГЕТСКА ПОТЕНЦИЈАЛНОСТ НА БИТОЛСКИОТ ДЕЛ ОД ПЕЛАГОНИСКАТА КОТЛИНА НА ПРОСТОРОТ ПОМЕЃУ СЕЛАТА ЛОЗНАНИ И КАНАТЛАРЦИ НА СЕВЕР И ОРИЗАРИ И НОВАЦИ НА ЈУГ <i>Никола Богатиновски, Стојанче Николов, Петре Пасков</i>	709
MINING / MINERAL SUPPORT SERVICES PROJECT <i>Duška Rokavec, Tina Benda</i>	719

3D MODELING OF SOME COPPER DEPOSITS IN THE REPUBLIC OF MACEDONIA

¹ Todor Serafimovski, ² Christos Christidis, ¹ Dalibor Serafimovski, ¹ Goran Tasev,
³ Mitko Ligovski, ⁴ Igor Ivanovski and ⁵ Lazar Gjorgjiev

¹Faculty of Natural and Technical Sciences, University "Goce Delcev"-Stip, , Republic of Macedonia

²National and Kapodistrian University of Athens, Faculty of Geology and Geoenvironment, Department of Economic Geology and Geochemistry, Panepistimiopolis, Zographou, 157 84, Athens, Greece

³Euromax Resources Skopje, Blvd Partizanski Odredi No.14, 1/2-3, Skopje, R. Macedonia

⁴Genesis Resources Ltd., Lermontova 3/2 Skopje, 1000 Skopje, R. Macedonia

⁵Kaltun Madencilik DOO Skopje, Boulevard Marx and Engels 1-5/2, Skopje, R. Macedonia

Abstract

This paper aims at presenting the 3D models of respective geological settings, deformation structures and geochemical distributions, in terms of better understanding of the related ore-genetic processes and exploration potential. Since the beginnings of mining industry the challenge of mineral exploration is to approach new exploration targets. In that direction, 3D and 4D modelling are the new exploration tools that can help the mineral explorers to visualise, interpolate and interpret geological data, which are critical time- and money-saving methods. In this study 3D-models had been constructed for several Macedonian Cu±Au polymetallic deposits Kadiica, Plavica, Kazan Dol and Borov Dol. Our 3D models were built by mapping geochemical variations in the Move software by Midland Valley for Kadiica Cu-porphyry deposit and Borov Dol Cu-Au porphyry deposit (gives the opportunity to the user to create slices of the 3D grid and take a closer look to the spatial variation of an assay) as well as in Target 3D for ArcGIS system for Cu-Au high-sulphidation epithermal Plavica deposit and Cu vein type deposit Kazan Dol.

Key words: copper deposits, 3D model, software, modeling, valorization

INTRODUCTION

During the last decade and half the extensive detailed geological exploration of particular ore deposits in the Republic of Macedonia took place. As it is already known, the territory of the Republic of Macedonia is a constitutional part of the geotectonic setting of the Serbo-Macedonian metallogenic province, which facilitates the formation of polymetallic mineralisations. This zone hosts many known deposits having been discovered and located in the areas of Kadiica deposit near the Pehcevo, Plavica deposit in the close vicinity of Kratovo and Probitip, Kazan Dol deposit near the Valandovo and Borov Dol deposit near the city of Radovis.

Both software represent a solid basis for modelling of certain copper and gold ore bodies within different deposits (Kadiica, Plavica, Kazan Dol and Borov Dol). Without any doubts a better understanding of the geometry and intensity in 3D will aid further detailed geological exploration at depth for all above mentioned ore deposits and may potentially lead to new discoveries. For each

aforementioned deposit were used lithological and geochemical data from exploration drill holes, which allowed construction of the 3D grid that depicts the distribution of each assayed element (Ag, As, Au, Mo, Pb, Zn, Cu). Also, there were created slices of the 3D grid and we were able to take a closer look to the spatial variation of an assay. For the 3D modelling of particular deposits were applied software packages that allow definition of morphological shapes of supposed ore body (ore bodies) at depth, which practically can suggest directions for their further exploration, especially if responsible person(s) decide that the exploration should continue with angled exploration drill holes.

REGIONAL AND LOCAL GEOLOGICAL SETTING

The metallogeny of the southern Balkan Peninsula is determined by the geodynamic evolution of the Tethys-Eurasian metallogenic belt (TEMB) and ancient crystalline massifs (Jankovic 1997). The southern Balkan Peninsula, including the Republic of

Macedonia, is a sector of the central TEMB. The formation of ore deposits was related here to certain chronological stages and geodynamic settings: (1) Late Permian-Early Triassic intracontinental rifting along the northern boundary of Gondwana and/or its already detached fragments; (2) Jurassic intracontinental rifting marked by ophiolitic complex; (3) suprasubduction magmatism; and (4) postcollision (orogenic) continental setting and related formation of various base- and noble-metal deposits (Pb-Zn-Ag, Cu-Au, Au-As-Sb-Tl, AuAg). The VZ extends in the NW-SE direction between the SMM and Pelagonian Massif. The SMM is located between the Carpathian Mountains and Balkanides in the east and the VZ in the west. The collision of the African and Eurasian plates was accompanied by closure of the ocean and subduction of the oceanic crust beneath the SMM. Deep faults that formed at the time at the active continental margins have important metallogenic implications. Collision, postcollision processes, and Tertiary magmatism were crucial for localization of base-metal ore mineralization. Other important ore-localizing structural elements are volcanic calderas, domes, and dikes. Three variously oriented fault systems are recognized. The first, NW-trending fault system coincides in the strike with major tectonic units. The second and third systems extend in the northeastern and meridional directions, respectively. The third system has important implications for the development of Tertiary magmatism and related iron skarn and porphyry copper mineralization. Volcanic activity progressed beginning from the late Oligocene, and ore mineralization formed in the Miocene. The geotectonic setting of Serbo-Macedonian Zone (SMZ) facilitates the formation of numerous copper mineralisations (Kadiica, Plavica, Buchim, Ilvotza and Osogovo in the Republic of Macedonia). The SMZ metallogenic belt consists of strongly deformed metamorphic rocks of Palaeozoic age superimposed by Tertiary magmatic activity. Substantial copper mineralisations and deposits occur associated with acidic stocks of Tertiary age, making shallow intrusions into SMZ Palaeozoic metamorphic formations. The mineralized and host-rocks belong to Oligocene-Miocene calc-alkaline complexes of dioritic, granodioritic and syenitic composition. It is

likely that the widespread calc-alkaline igneous suites resulted from an anatectic partial smelting of the lowermost continental crust, in post collision activity processes.

Local Geology of Kadiica – Bukovik/Kadiica region.- Bukovik-Kadiica deposit is located in the eastern part of the Republic of Macedonia in a hilly region close to the border with Bulgaria and 2 km northeast of the town of Pehcevo. The deposit can be described as one of a number of dacite plugs of Neogene origin that exist within the SMZ as part of the volcanic complex intruding into Palaeozoic sediments, andesites and gabbros. On a larger scale the Kadiica region is underlain by metamorphic rocks (metadiabases, schists, gabbro, diorite and younger granitoids) of Upper Proterozoic to Palaeozoic age (Tasev, 2010; Tasev et al., 2012). The Bukovik-Kadiica ore district has been located in the most eastern parts of the Besna Kobila-Osogovo-Tassos metallogenic zone (Aleksandrov, 1992; Janković et al., 1995) and has been characterized by complex polymetallic mineralisation. Within the same ore district there were determined ore body systems and intersected dykes of quartz-latites with an absolute age of 24–12 Ma (Serafimovski et al., 2001; Serafimovski T. et al., 2010). According to Singer et al. (2008) the ore of Kadiica deposit has an 1x0.6 km axis dimension and covers an area of 0,5 km². The ore veins are also variable, ranging from several centimeters up to 2 m, and rarely over 5 m in size. Investigations done so far have determined about 16 veins of NW-SE to NNW-SSE and ENE-WSW to E-W extension most commonly subparallel to each other. Occasionally they bend and intersect (commonly under a sharp angle) such as ore veins 2, 3, 4, 6 and 7 (Figure 1). Flank branching of main ore veins is common (ore vein 1) which results in formation of apophyses of small dimensions.

Local geology of the Plavica deposit.- As a part of well known Kratovo-Zletovo ore area in general it is characterized with a complex composition of mainly Tertiary volcanic-sedimentary complex imposed on the foundation of the Serbian-Macedonian massif, which represents constituting part of the western Tethyan belt. The district is host to vein-type Pb-Zn mineralisation, Cu-Au stockwork and disseminated mineralisation, and high-sulphidation epithermal Au depo-

stis. The Probistip Formation is comprised of andesitic tuff, sandstone, bituminous siltstone and volcanoclastic breccias. The Probistip Formation overlies the Zletovo Formation, which is composed of ignimbrite, volcanic breccia, and lava flows of dacitic and andesitic composition. The deposit is hosted by a sequence of andesitic to dacitic volcanic units of mixed sub-aerial and sub-aqueous nature, dipping approximately 30° south. Mineralisation occurs within steeply dipping, vuggy silica bodies up to 500 m long, and between 100 m and 150 m wide. At surface, the vuggy silica zones form prominent ridges trending W-WNW. The Plavica contains many of the features inherent in high sulphidation epithermal deposits such as: vuggy silica, advanced argillic alteration, in association with pyrite and sulphides and sulphosalts. High content of gold is associated with silica bodies, (Alderton and Serafimovski, 2007) and the advanced argillic alteration indicates existence of fluids with low pH.

Local geology of the Borov Dol ore deposit.- The Borov Dol orefield is localized at the margin of a volcanic caldera in the southern part of the ore district within the VZ. The NW-trending faults control the Neogene latite and quartz latite dikes, necks, and extrusions. The Borov Dol deposit is related to this volcanic-plutonic complex. The ore-bearing volcanic rocks are metasomatically altered. The volcanic rocks belong to andesite, latite, trachyte, and rhyolite, including transitional varieties. At the Borov Dol deposit, porphyry copper mineralization occurs as a ring around the subvolcanic andesitic body that cuts through older volcanic rocks. The intensely altered volcanic rocks host the ore. Potassic and quartz-sericite alteration are noted at the surface. Down the dip, the orebody morphology is conformable to the andesitic stock. The ore mineralization is stringer-disseminated; ore disseminations (70%) prevail over ore veinlets (30%) (Tudzarov and Serafimovski, 1995). Chalcopyrite as the major mineral in primary ore is accompanied by pyrite, molybdenite, magnetite, gold, bornite and sporadic enargite, famatinite, galena, and tennantite (Stefanova and Serafimovski, 2006).

Local geology of the Kazan Dol ore deposit.- The geological setting in the closest vicinity of the Kazan Dol deposit has been built by Precambrian metamorphic rocks

(gneiss in westernmost parts of the terrain as well as two-mica cordierite gneiss). Paleozoic products are represented by grey-greenish sericite-chlorite schist in southernmost parts of the area while the Mesozoic has been represented by Furka granites (central and south-eastern parts of the area), quartz and quartzless porphyries (along tectonic lines of NNW-SSE direction), quartz keratophyre and keratophyre. These granites have determined absolute age of 155±5 m.y. by Rb/Sr method, (Šoptrajanova, 1967) and 156±6 m.y. (Borsi et al., 1966). Magmatic rocks as diabase occur southern of the Tarla Tepe hill and are represented with NW-SE extension. At the contact with marble there were determined carbonate-limonitic breccia while at the contact with gneiss, silicification and feldspatization occurred. Preliminary information indicated that at the Kazan Dol locality were determined copper mineralizations in the oxidation zones (Ivanov, 1966) related to the Jurassic granitoids intruded into the crystalline schist, as well as copper veins (Janković et al., 1997) related to chlorite-mica schist that have formed narrow zone. Also, at the Kazan Dol locality were determined typical copper tetrahedrite veins, which define Alpine metallogeny in this part of the Republic of Macedonia (Serafimovski, 1995), as well as impregnations along the crushing zones intercalated with quartz veins (Serafimovski et al., 1997). Localized areas of quartz veining are associated with copper mineralization with individual veins rarely exceeding 150 metres in strike (Čifliganec, 1993). Veins are usually narrow ranging from millimetre stock-work veins to individual veins up to a few metres thick. Primary mineralization consists of rarely observed chalcopyrite and very minor pyrite while locally pyrrotite, sphalerite and tetrahedrite-tennantite have been reported within narrow veins. Secondary copper minerals are very widespread on the Kazan Dol area. The main oxide copper minerals are tenorite, malachite and locally cuprite, although azurite, native copper, covellite, chrysocolla, limonite and manganese minerals are widespread in the Kazan Dol mineralized area.

RESULTS AND DISCUSSION

The Kadiica ore deposit.- Using the lithological and the geochemical data from

11 drill holes at the Kadiica deposit, a 3D grid was created that depicts the distribution of each assay (Ag, As, Au, Mo, Pb, Zn, Cu; Fig. 1a). The software gives the opportunity to the user to create slices of the 3D grid and take a closer look to the spatial variation of an assay. From the grid showing the Cu distribution, an ore body with dimensions of 1000 x 600 m and 50 m thick, could be indi-

cated. From the geochemical distributions it could be illustrated that high values of As are followed by high values of Au and Pb and low values of Zn. Different to Au and Pb the Cu mineralization is located at deeper levels. Also, higher-grade Cu mineralization appears in the western part of the main ore body (Fig.1b).




Fig. 1 a) 3D geochemical distribution in Kadiica deposit for the following assays: Ag, As, Au, Mo, Pb, Zn and b) 3D geochemical distribution of Cu along E-W geological sections across Kadiica deposit.

The Plavica ore deposit.- A comprehensive and in many ways detailed study conducted at the Plavica, beside geochemistry and geophysics, included respectable array of 195 exploration boreholes totaling 47300 m.

We used all the boreholes, including the negative one so we can get more representative situation in the 3D modelling. The study results have shown interesting gold and copper concentrations in certain mineralized

zones, which are the basis for construction of ours 3D model with use of professional computer software package Target 3D for ArcGIS in integration with data in form of MapInfo tables and 2D DXF files. Although mineralisation estimation of domains was modeled based only on metal grade, with definition of cut-off grade and top cut grade values for the Plavica deposit, the ore body was 3D modelled, with respect to intersecting fault zones, overall geological setting and the compositional distribution of metal grades (Figure 2a/b). Setting up the complete 3D geological model of the Plavica ore deposit all available data and sources of information were compared and integrated as it was case with 3D deposits elsewhere (Arvanitidis et al., 2013).

The contours of the ore bodies, both for gold and copper, have shown strong correlation to the fault systems in the area (being of N-S, NW-SE and sporadically SW-NE direction and respectively reflecting their spatial position. Following sound geological reasoning during the creation of the model, a considerable amount of knowledge regarding the mineralized area and the ore body was specifically updated (mineralization extending up to 300 m at depth); some new N-S striking and cross-cutting fault structures were interpreted. For completion of this 3D model two different methods are used. The first is projection using block with help of kriging method and displayed volume through 30 m blocks. Figure 2a represents model of Au with 0.2g/t cutoff grade. Figure 2b represents Cu model with 0.12% cutoff grade.




Fig. 2 a) 3D model of Au (seen from SE); **b)** 3D model of Cu (Seen from SE); **c)** 3D model for Au using “solid” display (Seen from SE); **d)** 3D model for Cu using “solid” display (Seen from SE)

This 3D model displays visible concentration of Au in the NE part of the deposit. In the western part there is also smaller ore body deeper underground. Looking at this model one can clearly see that Cu mineralization is

not connected with Au mineralization. Cu is present through all the area, slightly deeper than Au. We would like to stress out that the actual 3D model of the Plavica polymetallic deposit we can see that it strongly reflects the

Трет Конгрес на Геолозите на Република Македонија
Third Congress of Geologists of Republic of Macedonia

geochemical Au-Cu pattern already seen at the geochemical geochemical cross sections (Serafimovski et al., 2014). The second method used for 3D modelling is the method of isosurface where “solid” bodies are formed. In our case we used different cutoff grade for projecting the solid bodies with cutoff grade of 0.4g/t for Au and 0.2% for Cu (Figure 2c/d).

The Borov Dol ore deposit.- After an extensive geological exploration programme at the Borov Dol ore deposit was finished, based on geochemistry, geophysics and exploration data of around 100 exploration drill holes totaling more than 20 000 m we proceeded with construction of an appropriate 3D model. The study results have

shown interesting copper concentrations between levels 630m and 300 m, which represents the basis for construction of ours 3D model. The drill holes were selected on positive vs. negative basis and were taken only those that positively contribute to the copper body contouring. When plotted in 3D, the majority of the geochemical groupings have a strong association with the lithologies of the deposit sequence (Figure 3e/f). Newly defined morphological contours of the ore bodies within the Borov Dol deposit basically are of stockwork-impregnated type (Figure 3c/d), which in general mimic inverted glass cup or cone that at depth deforms.




Fig. 3 a) Surface and the drill holes; b) Variation of copper in the drill holes; c)/d) cross sections from west to south on the grid with the variation of copper; e)/f) cross sections, striking NNW-SSE and NW-SE, cutting through Borov Dol Cu mineralization

The Kazan Dol ore deposit.- Detailed study conducted at the Kazan Dol locality beside geochemistry, geophysics and exploration trenches included respectable array of 53 exploration drill holes totaling 3442 m with an average depth of 90 m and 3052 samples. The study results have shown interesting copper concentrations in oxidation zones, which represents the basis for

construction of ours 3D model. The drill holes were selected on positive vs. negative basis and were taken only those that positively contribute to the copper body contouring. When plotted in 3D, the majority of the geochemical groupings have a strong association with the lithologies of the deposit sequence (Figure 4).




Fig. 4 a) 3D model of the Kazan Dol deposit with terrain surface displayed; b) 3D model of the deposit

The biggest advantages of constructing 3D volumes instead of 2D cross sections is the possibility to define ore grade volume gains and losses of the entire mineralized systems (Chmielowski et al., 2013). Construction of 3D model of this particular ore deposit will be used to constrain key features pertinent to ore genetic models, such as net gains or losses of elements within the entire ore system as well as the source of metals in the deposits as it was mentioned elsewhere

CONCLUSION

At the end of this compilation study some conclusions stand out itself. At the Bukovik-Kadiica deposit 3D model suggests Cu distribution (within an ore body) with dimensions of 1000 x 600 m and 50 m thickness. Geochemical distributions illustrated that high values of As are followed by high values of Au and Pb and low values of Zn, while opposite to Au and Pb the Cu mineralization is located at deeper levels. The Borov Dol 3D model confirmed interesting Cu concentrations between levels 630 m and 300 m, as well as that the deposit basically is of stockwork-impregnated type, which in general mimic inverted glass cup or cone that at depth deforms.

(Fisher et al., 2013; Chmielowski et al., 2013). During the creation of the model, a considerable amount of knowledge concerning the mineralization area and the ore body was specifically updated (thickness of more than 160 m); some new N-S striking and cross-cutting fault structures were interpreted, but further investigations are required to determine their metallogenic relationship to any of the mineralization processes.

At the Plavica deposit construction of the 3D model have revealed its isometric to lens/lensoid morphology, which swerves and distorts at depth. It has been proved with the 3D model spatial position of gold and copper mineralization within the Plavica deposit between hypsometric levels 1305 and 600 m. The present exploration at the Kazan Dol continues in the southern parts of the deposit where exist positive Cu and Zn anomalies as a direct reflection of oxidized Cu-veins enclosed in crushing zones. Displayed 3D model defines the spatial position of Cu mineralization in the Kazan Dol deposit between hypsometric levels 340 and 180 m.

REFERENCES

- Alderton, H. M. D. and Serafimovski, T., 2007. The geology and genesis of the Plavica copper-gold deposit, Macedonia. *Applied Earth Science (Trans. Inst. Min. Metall. B)*, Vol. 116, No. 2, p. 94–105 (2007).
- Aleksandrov, M., 1992. Metallogenetic features of the polymetallic ore field Sasa-Eastern Macedonia. Ph.D. Thesis, Faculty of Mining and Geology-Štip, Macedonia, 264 p.
- Arvanitidis, N. D., Michael, C., Christidis, C., Weihed, P., Gaal, G., Royer, J. J., Perantonis, G., Bakalis, V. and Ballas, D., 2013. Using 3D/4D modelling tools in exploration for epithermal gold potential areas in Eastern Rhodope zone (Western Trace, NE Greece)-Mineral deposit research for a high-tech world, *12th SGA Biennial Meeting, Proceedings, Volume 1*, Sweden, pp 58–61.
- Borsi, S., Ferrara, G. and Mercier J., 1966. Mesures d'âge par la methode Rb/Sr des granites filoniens de Karathodhoro (Zones internes des Hellenides, Macedoine Centrale Greece). *Proceedings 6th Counseling, part II Geochemistry and ore deposits*, pp. 5–10.
- Chmielowski, M. R., Jansson, N., Persson, MF., Fagerström, P. and Weihed, P., 2013. 3D geochemical modelling of hydrothermal alteration related to 1.89 Ga VHMS-type deposits, Kristineberg area, Skellefte District, Sweden. *Proceedings of the 12th Biennial SGA Meeting*, Uppsala, Sweden, pp. 66–69.
- Čifliganec, V., 1993. Copper Mineralization in the Republic of Macedonia: Types and Distribution Patterns. University “Sts. Cyril and Methodius” Skopje, Faculty of Mining and Geology-Štip, Spec. Issue, 1, 303 p. (in Macedonian with extended summary in English)
- Fisher, A. L., Cleverley, S. J., Pownceby, M. and MacRae, C., 2013. 3D representation of geochemical data, the corresponding alteration and associated REE mobility at the Ranger uranium deposit, Northern Territory, Australia. *Mineralium Deposita*, v. 48, no. 8, pp 947-966.
- Ivanovski, T., 1966. Guide for basic geological map, scale 1:100,000, Sheet Gevgelija. Federal Geol. Surv. Belgrade, Yugoslavia, 52 p.
- Janković, S., 1997. The Carpatho-Balkanides and adjacent area: a sector of the Tethyan Eurasian metallogenic belt. *Mineralium Deposita*, v. 32, pp. 426–433.
- Jankovic, S., Serafimovski, T., Aleksandrov, M., 1995. The Besna Kobila-Osogovo Metallogenic zone. *Geologica Macedonica*, Vol. 9, No. 1, 39-50, Štip.
- Jankovic, S., Serafimovski, T., Jelenkovic, R. and Cifliganec, V., 1997. Metallogeny of the Vardar zone and Serbo-Macedonian Mass. Symposium-Annual Meeting, Proceeding, Dojran, pp. 29-67.
- Serafimovski, T., 1995. Copper Deposits of the Vardar zone SE Europe: Types and Distribution Patterns. International Earth Sciences Colloquium on the Aegean Region. Izmir, Turkey, Proceed., Vol. II, pp. 623-637.
- Serafimovski, T., Tomson, I. N. and Kocneva, N. T., 1997. Alpine orogenic structures and metallogeny in the Serbo-Macedonian massif and the Vardar zone of the territory of Macedonia. Symposium-Annual Meeting, Proceeding, pp. 113-117.
- Serafimovski, T., Cifliganec, V. and Tasev, G., 2001. The porphyry Cu-Au mineral deposits related with small volcanic intrusions in the Republic of Macedonia. ABCD-GEODE workshop Vata Bai, Romania. Vol. 79, suppl. 2, pp 114a.
- Serafimovski, T., Tasev, G., Blažev, K. and Volkov A., 2010. Major alpine structures and Cu-porphyry mineralization in the Serbo-Macedonian massif. *Geologica Macedonica*, Vol. 24 (1), pp. 39–48.
- Serafimovski, T., Aleksandrov, M., Tasev, G., Stefanova, V., Paterson, J., Stevanović, M., Ristovski, V. and Ivanovski, I., 2014. Elaborate of detailed geological study, with calculation of geological ore reserves, of copper and gold in the Plavica, Kratovo locality. Faculty of Natural and Technical Sciences, University “Goce Delcev” for Genesis Resources International, Skopje, 431 p. (in Macedonian).
- Singer, D.A., Berger, V.I. and Moring B.C., 2008. Porphyry copper deposits of the world: Database and grade and tonnage models, 2008: U.S.G.S. Open-File Report 2008-1155, 45p.
- Šoptrajanova, G., 1967. Petrological and geochronological characteristics of some granitoids of Macedonia. PhD thesis, Faculty of Mining and Geology, University of Belgrade, 105 p.
- Tasev, G., 2010. Metallogeny of the polymetallic ore-bearing system Bukovik-Kadiica. PhD Thesis, University “Goce Delcev”-Štip, Faculty of Mining and Geology, 206 p.
- Tasev, G., Serafimovski T., Stefanova V., 2012. New data of Fluid Inclusions Studies of the Bukovik-Kadiica deposit, Republic of Macedonia. SCOPES Project-Internat. Conference.
- Tudzarov, N. and Serafimovski, T., 1995. Geotectonic setting and Metallogenetic features of the Borov Dol Porphyry Copper Deposit. *Geologica Macedonica*, Štip, Vol. 9, No. 1, pp. 89-96.