



UNIVERZITET U
KRAGUJEVCU
AGRONOMSKI FAKULTET U
ČAČKU



UNIVERSITY OF
KRAGUJEVAC
FACULTY OF
AGRONOMY
CACAK

XXII SAVETOVANJE O BIOTEHNOLOGIJI

sa međunarodnim učešćem

- ZBORNIK RADOVA 1 -



Čačak, 10 - 11. Mart 2017. godine

XXII SAVETOVANJE O BIOTEHNOLOGIJI

sa međunarodnim učešćem

- Z b o r n i k r a d o v a 1 -

ORGANIZATOR I IZDAVAČ

**Univerzitet u Kragujevcu,
Agronomski fakultet u Čačku**

Organizacioni odbor

prof. dr Gordana Šekularac, dr Pavle Mašković, dr Milun Petrović, dr Gorica Paunović, prof. dr Milomirka Madić, dipl. ing. Srđan Bošković

Programski odbor

prof. dr Leka Mandić, prof. dr Vladeta Stevović, prof. dr Dragutin Đukić, prof. dr Snežana Bogosavljević-Bošković, prof. dr Tomo Milošević, prof. dr Milica Cvijović, prof. dr Radojica Đoković, prof. dr Milomirka Madić, prof. dr Goran Dugalić, prof. dr Aleksandar Paunović, prof. dr Radoš Pavlović, prof. dr Milena Đurić, prof. dr Gordana Šekularac, prof. dr Biljana Veljković, dr Nikola Bokan, dr Drago Milošević, dr Lenka Ribić-Zelenović, dr Vladimir Kurćubić, dr Goran Marković, dr Ljiljana Bošković-Rakočević, dr Gorica Paunović, dr Milun Petrović, dr Milan Lukić, dr Slavica Vesković

Tehnički urednici

dr Milun Petrović, dipl. ing. Dušan Marković, dipl. ing. Srđan Bošković

Tiraž: 180 primeraka

Štampa

Grafička radnja štamparija Bajić, V. Ignjatovića 12, Trbušani, Čačak

PREDGOVOR

Pored osnovne obrazovne delatnosti na osnovnim, master i doktorskim akademskim studijama, kao i realizaciji posebnih programa za stalno stručno usavršavanje, Agronomski fakultet u Čačku obavlja i niz drugih delatnosti kao što su: izvođenje naučno-istraživačkih projekata samostalno i/ili u saradnji sa drugim organizacijama iz zemlje i inostranstva, primena tehničko-tehnoloških rešenja u praksi, publikovanje naučnih i stručnih radova kroz sopstvenu izdavačku delatnost i organizovanje naučnih i stručnih skupova.

Značajna aktivnost Agronomskog fakulteta u Čačku, preko dvadeset godina, je organizovanje naučno-stručnog skupa Savetovanje o biotehnologiji. Osnovni cilj savetovanja je upoznavanje šire naučne i stručne javnosti kao i poljoprivrednih proizvođača i prerađivača sa rezultatima najnovijih naučnih istraživanja domaćih i inostranih naučnih radnika iz oblasti primarne poljoprivredne proizvodnje i prerade. Na taj način fakultet nastoji da omogući primenu naučnih rezultata široj proizvodnoj praksi.

Zbornik radova XXII Savetovanja o biotehnologiji sa međunarodnim učešćem sadrži 112 radova iz oblasti: Ratarstva, Povrtarstva i krmnog bilja, Vinogradarstva i voćarstva, Stočarstva, Prehrambene tehnologije i Zaštite bilja, proizvoda i životne sredine.

Pored naučnih radnika iz gotovo svih visokoobrazovnih i naučnih institucija Republike Srbije, na ovogodišnjem savetovanju učestvuju i naučni radnici iz Bosne i Hercegovine, Makedonije, Bugarske i Crne Gore.

XXII Savetovanje o biotehnologiji sa međunarodnim učešćem održće se pod pokroviteljstvom Ministarstva za prosvetu, nauku i tehnološki razvoj Republike Srbije, uz materijalnu pomoć grada Čačka i donatorskih firmi iz oblasti primarne poljoprivredne proizvodnje i prerade.

Programski i organizacioni odbor XXII Savetovanja o biotehnologiji izražava veliku zahvalnost svim institucijama, organizacijama i kolegama koji su direktno učestvovali ili na bilo koji način pomogli u organizaciji ovog skupa.

U Čačku, marta 2017. godine

Programski i Organizacioni odbor
XXII Savetovanja o biotehnologiji

SADRŽAJ

Sekcija: Ratarstvo, povrtarstvo i krmno bilje

<i>Miodrag Jelić, Gordana Šekularac, Vera Đekić, Goran Dugalić, Aleksandar Paunović, Milomirka Madić, Ivica Đalović: SADRŽAJ I MOBILNOST GVOŽĐA U KISELIM ZEMLJIŠTIMA CENTRALNE SRBIJE</i>	11
<i>Desimir Knežević, Aleksandar Paunović, Mirjana Menkovska, Pavle Mašković, Danijela Kondić, Milica Zelenika, Milomirka Madić, Veselinka Zečević: PROTEINI GLUTENA – DETERMINANTA KVALITETA PŠENICE</i>	17
<i>Ivica Đalović, Yinglong Chen, Zed Rengel, Srđan Šeremešić, Željana Prijić, Miodrag Jelić: ARHITEKTURA KORENOVOG SISTEMA KUKURUZA ZA EFIKASNIJE USVAJANJE FOSFORA: NOVIJA SAZNANJA</i>	23
<i>Anja Dolapčev, Slaven Prodanović, Vladimir Sikora, Tomislav Živanović, Sanja Vasiljević, Dura Karagić, Snežana Katanski: ANALIZA ODNOSA VISINE BILJKE I DUŽINE METLICE U NS-KOLEKCIJI LINIJA OPRAŠIVAČA SIRKA ZA ZRNO</i>	31
<i>Livija Maksimović, Dušan Adamović, Larisa Merkulov-Popadić, Borivoj Pejić, Vera Popović: UTICAJ SISTEMA GAJENJA U PROMENLJIVIM VREMENSKIM USLOVIMA NA PRINOS BOSILJKA</i>	37
<i>Gorica Cvijanović, Svetlana Roljević, Nenad Đurić, Gordana Dozet, Vojin Đukić, Vojin Cvijanović: UTICAJ RAZLIČITIH VARIJANTI ĐUBRENJA NA KVANTITET I KVALitet ZRNA PŠENICE.....</i>	45
<i>Svetlana Roljević Nikolić, Gorica Cvijanović, Jelena Marinković: PRINOS ZRNA RAZLIČITIH SORTI PŠENICE U ORGANSKOJ PROIZVODNJI U ZAVISNOSTI OD RIZOSFERNE MIKROFLORE</i>	53
<i>Ljubiša Kolarić, Tihomir Gujančić, Goran Jaćimović, Vera Popović, Jela Ikanović, Ljubiša Živanović: UTICAJ GUSTINE USEVA NA PRINOS I KVALITET KORENA ŠEĆERNE REPE</i>	61
<i>Gordana Dozet, Gorica Cvijanović, Slobodan Milenković, Nevena Ninkov, Ljubiša Kostov, Dragana Kaluđerović: KOMPONENTE PRINOSA PASULJA U ZAVISNOSTI OD PRIMENE GUANITA I MIKROBIOLOŠKIH ĐUBRIVA.....</i>	69
<i>Ljubica Šarčević - Todosijević, Ljubiša Živanović: MIKROBIOLOŠKA AKTIVNOST ZEMLJIŠTA TIPA ČERNOZEM U ZEMUN POLJU</i>	75
<i>Gordana Dozet, Vojin Đukić, Svetlana Balešević-Tubić, Nenad Đurić, Zlatica Miladinov, Jovica Vasin, Snežana Jakšić: UTICAJ PRIMENE VODENIH EKSTRAKATA NA PRINOS U ORGANSKOJ PROIZVODNJI SOJE</i>	81
<i>Aleksandar Paunović, Milomirka Madić, Desimir Knežević, Miodrag Jelić, Vladanka Stupar: SPECIFIČNOSTI ĐUBRENJA JEČMA</i>	87
<i>Nebojša Gudžić, Miroljub Aksić, Slaviša Gudžić, Miodrag Jelić, Aleksandar Vuković: EFEKAT TRI NIVOJA KALCIFIKACIJE NA SADRŽAJ MOBILNOG AI I KISELOST KOD DISTRIČNO SMEĐEG ZEMLJIŠTA</i>	93
<i>Zoran Jovović, Dragan Mandić, Novo Pržulj, Ana Velimirović, Željko Doljanović: GENETIČKI RESURSI PŠENICE (<i>Triticum sp.</i>) U CRNOJ GORI</i>	99

<i>Zoran Jovović, Milana Šilj, Ana Velimirović, Novo Pržulj, Dragan Mandić: GENETIČKI RESURSI JEĆMA (<i>Hordeum sativum</i> Jess.) U CRNOJ GORI</i>	109
<i>Dalibor Tomić, Vladeta Stevović, Dragan Đurović, Nikola Bokan, Milomirka Madić: UTICAJ KALCIZACIJE ZEMLJIŠTA NA PRINOS SORTI ŽUTOG ZVEZDANA U TREĆOJ GODINI PROIZVODNJE</i>	117
<i>Dragica Spasova, Adrijana Burovska, Biljana Atanasova, Dusan Spasov, Mite Ilievski: ANALYSIS OF THE QUALITY OF OATS (<i>AVENA SATIVA</i> L.) GROWN IN CONDITIONS OF ORGANIC PRODUCTION</i>	123
<i>Dragoslav Đokić, Rade Stanisljević, Dragan Terzić, Jasmina Milenković, Goran Jevtić, Saša Barać, Bojana Milenković: ANALIZA KVALITETA I KOLIĆINE SEMENA VIŠEGODIŠNJIH TRAVA PRI PROCESU DORADE</i>	131
<i>Dragutin Đukić, Leka Mandić, Aleksandra Stanković-Sebić, Vladeta Stevović, Marijana Pešaković, Milica Zelenika, Vesna Đurović: INFICIRANJE LEGUMINOZNIH BILJAKA SA KVRŽIČNIM BAKTERIJAMA</i>	137
<i>Goran Perković, Aleksandra Govedarica-Lučić, Andrija Tomić: UTICAJ RAZLIČITIH NAČINA GAJENJA PAPRIKE NA DEBLJINU PERIKARPA PLODA</i>	147
<i>Goran Perković, Siniša Berjan, Branka Govedarica, Igor Đurđić, Radomir Bodiroga, Andrija Tomić: ORGANSKA POLJOPRIVREDA U FUNKCIJI ODRŽIVOG RAZVOJA RURALNIH PODRUČJA REPUBLIKE SRPSKE</i>	153
<i>Gorica Delić, Snežana Branković, Mirjana Staletić, Milivoje Milovanović: UTICAJ SOLI NATRIJUMA NA KLIJANJE SEMENA I RAZVOJ KLIJANCA JEĆMA (<i>HORDEUM VULGARE</i> L.), JARE SORTE JADRAN</i>	159
<i>Hristofor Kirchev, Stefka Dobreva, Angelina Muhova: PRODUCTIVITY AND QUALITY OF DURUM WHEAT (<i>TRITICUM DURUM</i> DESF.) AT INCREASING RATES OF NITROGEN FERTILIZATION UNDER LONG-TERM ACCUMULATION OF NUTRIENTS IN PELIC VERTISOLS</i>	165
<i>Ivan Yanchev: PRODUCTIVITY AND QUALITY OF BULGARIAN LAVENDER VARIATIES</i>	171
<i>Jordan Marković, Snežana Andelković, Tanja Vasić, Ivica Kostić, Snežana Babić, Dragoslav Đokić, Mirjana Petrović: PRINOS SUVE MASE LUCERKE U ZAVISNOSTI OD SORTE, FAZE RAZVIĆA I OTKOSA</i>	177
<i>Ljubiša Živanović, Vera Popović, Jela Ikanović, Ljubiša Kolarić: UTICAJ KOLIĆINE I OBLIKA AZOTA NA PRODUKTIVNOST OZIME PŠENICE</i>	183
<i>Marijana Dugalić, Ljiljana Bošković-Rakočević, Nikola Bokan, Aleksandar Paunović, Fejzo Begović: UTICAJ RAZLIČITE DOZE I NAČINA PRIMENE MINERALNIH ĐUBRIVA NA PRINOS KROMPIRA</i>	189
<i>Milomirka Madić, Aleksandar Paunović, Miodrag Jelić, Desimir Knežević, Dragan Đurović: PRINOS I KOMPONENTE PRINOSA ZRNA OZIMOG DVOREDOG JEĆMA GAJENOG NA KISELOM ZEMLJIŠTU</i>	195
<i>Svetla Kostadinova, Galia Panayotova: REACTION OF BARLEY VARIETIES TO NITROGEN-POTASSIUM FERTILIZATION</i>	201
<i>Vera Đekić, Jelena Milivojević, Mirjana Staletić, Jelić M., Vera Popović, Snežana Branković, Terzić D: UTICAJ MINERALNE ISHRANE NA PRINOS OZIME PŠENICE (<i>Triticum aestivum</i> L.)</i>	207

Vera Popović, Vladimir Sikora, Ljubiša Živanović, Milić Čurović, Dragan Terzić, Ljubiša Kolarić, Vera Rajićić, Jela Ikanović: SORTA FACELJE NS PRIORA ZA PROIZVODNJU BIOMASE U CILJU DOBIJANJA VOLUMINOZNE STOČNE HRANE 213

Sekcija: Voćarstvo i vinogradarstvo

Mlađan Garić, Vera Vukosavljević, Ivana Radojević, Miloš Ristić: PRINOS I KVALITET KLONOVA SORTE KABERNE SOVINJON U NIŠKOM REJONU 223

Tatjana Popović, Armin Kalač, Danijela Raičević, Radmila Pajović: RODNOST I KVALITET GROŽĐA SORTE ŽIŽAKU PODGORIČKOM VINOGORJU 231

Mirko Kulina, Gorica Paunović, Mirjana Radović, Andrijana Mitrović: FIZIČKO-HEMIJSKA SVOJSTVA PLODA NEKIH SORTI KRUŠKE (*Pyrus communis* L.) NA PODRUČJU SARAJEVSKE REGIJE 239

Dijana Stojanov: STANJE, PROBLEMI I PERSPEKTIVA PROIZVODNJE VOĆA U ZLATIBORSKOM OKRUGU 247

Jaćimović Vučeta, Božović Dina: POMOLOŠKE OSOBINE SORTI ORAHA U AGROEKOLOŠKIM USLOVIMA BIJELOG POLJA 253

Томо Милошевић, Небојша Милошевић, Иван Глишић: НЕКИ АТРИБУТИ ВЕГЕТАТИВНОГ РАСТА, ПРИНОСА И КВАЛИТЕТА ПЛОДА КАЈСИЈЕ (*Prunus armeniaca* L.) У ЗАВИСНОСТИ ОД СОПТЕ И ПОДЛОГЕ 257

Slavko Mijović, Saša Ilić, Ranko Popović, Tatjana Popović: UTICAJ FOLIJARNE PRIMJENE MAKRO I MIKRO ELEMENATA NA AGROBIOLOŠKE KARAKTERISTIKE SORTE MUSKAT ITALIJA NA ĆEMOVSKOM POLJU 267

Nebojša Milošević, Ivana Glišić, Milan Lukić, Milena Đorđević: BIOLOŠKO-POMOLOŠKE OSOBINE AUTOHTONIH SORTI ŠLJIVE U AGROEKOLOŠKIM USLOVIMA ČAČKA 273

Blaga Radovanović, Jasmina Dimitrijević, Aleksandra Radovanović, Jelena Mladenović, Nedeljko Manojlović: KINETIC STUDY OF OXIDATION DEGRADATION OF POLYPHENOLS IN SOUR CHERRY AND BLACKBERRY EXTRACTS DURING STORAGE 281

Ivan Glišić, Miloš Perišić, Tomo Milošević, Gorica Paunović: UTICAJ UKLANJANJA PRVIH SERIJA IZDANAKA NA VEGETATIVNI RAST, PRINOS I KVALITET PLODA KUPINE (*Rubus fruticosus* L.) 289

Sekcija: Zaštita bilja, proizvoda i životne sredine

Goran S. Marković, Jelena B. Popović-Dorđević, Nebojša D. Pantelić, Biljana P. Dojčinović, Aleksandar Ž. Kostić: ISPITIVANJE ODABRANIH FIZIČKO-HEMIJSKIH OSOBINA I SADRŽAJA MAKRO- I MIKROELEMENATA U VODI JEZERA GRUŽA 297

Grujica Vico, Danijel Mijić, Radomir Bodiroga: PRIMJENA POSLOVNE INTELIGENCIJE ZA ANALIZU PODATAKA I PODRŠKU ODLUČIVANJU U POLJOPRIVREDI 303

Slobodan Vlajić, Stevan Maširević, Rade Barać, Renata Iličić, Jelica Gvozdanović – Varga, Vladimir Božić: BOLESTI KUPUSA TOKOM 2016. GODINE 309

Markola Saulić, Ivica Đalović, Dragana Božić, Sava Vrbničanin: REZERVE SEMENA KOROVSKIH BILJAKA U KONVENTIONALNOM SISTEMU GAJENJA KUKURUZA. 315

Drago Cvijanović, Nataša Simić, Svetlana Vukotić: UPOTREBA ALTERNATIVNIH IZVORA ENERGIJE U POLJOPRIVREDI U CILJU ZAŠTITE ŽIVOTNE SREDINE 321

ANALYSIS OF THE QUALITY OF OATS (*AVENA SATIVA L.*) GROWN IN CONDITIONS OF ORGANIC PRODUCTION

Dragica Spasova¹, Adrijana Burovska¹, Biljana Atanasova¹, Dusan Spasov¹, Mite Ilievski¹

Abstract

During 2015 and 2016 the research of 11 oats genotypes is conducted, in conditions of organic production. Three of the oats genotypes are domestic populations, *Krivogastani*, *Trebenista* and *Kuceviste*, and the others are introduced varieties: *Rajac*, *Slavuj* and *Lovken* from Serbia, *Kupa*, *Baranja*, *Eksplorer*, *Sampionka* and *Istra* from Croatia. The phenological surveys, during the research showed that the vegetation period of the spring oat genotypes, grown in condition of organic production, is 100 – 110 days. The oats was sown in March, and the full maturity was reached in July, in each year of the research. In both years of research, the highest absolute mass of the grain had variety *Istra* (34,60 g in 2015 and 29,60 in 2016), while, the lowest had the population *Krivogastani* (12,3 g in 2015 and 14,90 g in 2016). There are very significant statistical differences between the varieties. The hectoliter mass of the grain is statistically different in different varieties, in the both years of the research. *Rajac* variety showed the lowest hectoliter mass in both experimental years (23,66 kg/hl in 2015 and 31,5kg/hl in 2016). The highest hectoliter mass in 2015 had *Istra* variety (42,05 kg/hl), and in 2016, it was *Krivogastani* population (36 kg/hl). The highest oats plants, in both years of research, were *Krivogastani* population (116,7 cm in 2015 and 143,3 cm in 2016). The lowest plants were *Kupa* variety (78,3 cm in 2015 and 83,4 cm in 2016). The comparison between varieties showed that there are statistically significant differences.

Key words: Oats, organic production, absolute mass, hectoliter mass, height.

Introduction

The oats (*Avena sativa L.*) originates from the Middle East. During the Bronze Age (1500 – 1700 BC) the oats was introduced in the colder and more humid parts of Europe and is therefore considered as "European wheat". It is thought that oats originates mainly from two species: the wild oats (*Avena fatua L.*) and the red wild oats (*Avena sterilis L.*) (Suttie, 2000).

In the past it was used for production of bread, but with increasing the standards the oats began to be used for food production (oat flakes, flour, cereals, muesli). According to (Vasilcenko et al. 1985), (Panajotova et al. 2003) and (Savova et al. 2005), one of the most quality features of oats, that makes it important raw material in the food industry,

¹"Goce Delcev" University, Faculty of Agriculture, Stip – Republic of Macedonia
biljana.atanasova@ugd.edu.mk, dragica.spasova@ugd.edu.mk

is the high protein content and the high absolute mass. The quality of the grain and the modest demands for cultivation give the oats special importance.

Compared to other cereals the oats contain up to three times more fat, and the protein are significantly digestible. The absolute and the hectoliter mass are in medium positive dependencies with the proteins, but in much stronger positive dependencies with the β glutens (Popov et al., 2009). The hectoliter mass is the basic physical indicator for determining the quality of the grain, which depends on the ratio between the essential substances that build grain (carbohydrates, proteins and fats), the absolute mass, the size and the fulfillment of the grain. It serves as indicator for the potential yields and quality or, the output of the flour (Zorovski et al., 2014).

Today, the great number of modern diseases is treated with healthy food. The organic production excludes the use of chemical means and allows protection of the plants only with natural resources. Studies of certain oats varieties in terms of organic farming, which committed (Galie et al. 2004), showed that oats is very suitable for organic production, taking into account the high yields he got, which ranged from 4 to 5 t/ha. Similar results got (Konstantinos, 2007), which cultivated new varieties of barley and oats in terms of organic production, which showed great stability in yield, good productivity and resistance to disease.

On the other hand, the development of farming and livestock production conditioned increasing of areas under crops, including oats. In the research of (Ilievski et al. 2015), it is concluded that the total production of oats in 2014 increased for 2 135 tons or 54.7%, compared with 2012.

Taking the above in mind, the goal of our research was to determine some important morphological and physical properties of oats grown in organic production, as well as the differences between the tested varieties and populations. The analysis will also determine which one is most suitable for growing in organic production in the Strumica region, i.e. which variety or population will guarantee high quality.

Material and methods

The research for organic production of oats began in 2015 and end up in 2016. The trials were conducted in field and laboratory conditions. The field trials were set on the experimental field of the "Goce Delchev" – University, Faculty of Agriculture – Strumica, and laboratory tests were conducted in the laboratories of the Faculty of Agriculture in Stip.

11 oats genotypes were analyzed, from which three were domestic populations (*Krivogastani*, *Trebenista* and *Kuceviste*), and the others are introduced varieties: *Rajac*, *Slavuj* and *Lovken* from Serbia, *Kupa*, *Baranja*, *Eksplorer*, *Sampionka* and *Istra* from Croatia.

The experiments, in both years of the research, were set in three repetitions, deployed by the method of random block system, with dimension of basic plot of 5 m².

The distance between the variants was 0,50 m, and between repetitions – 1,0 m. The distance between rows was 20 cm. The sowing rate was 550 grains per 1m², i.e. 5.5 million grains per 1ha. The preparation of the soil, in both years of the research was identical. The autumn basic processing of the soil was at a depth of 35 cm. Then, there

was additional processing and fertilization with 30 t/ha bio-fertilizer, according to the regulations for organic production. The sowing, in both years of the research was performed during March (23.03.2015 and 28.03.2016). Sowing was performed manually, in rows, at a depth of 5-6 cm. The harvest was in July (03.07.2015 and 07.07.2016). Before harvest, material from 1 m² was taken from each parcel, for laboratory tests (absolute and hectoliter mass of the grain, by the international methods of ISTA).

During vegetation the most important growing stages were monitored: emergence, tillering, booting, tasseling and maturity, by the method of Cooperman (1955).

Also, before harvest, the height of 10 plants from each parcel (30 plants of each variant) was measured.

The results were processed by analysis of variance method, and the differences were tested on LSD – test.

Results and discussion

1. Vegetative growing – phenological observations

The vegetative growth of oats goes through more stages, in which, the formation and growth of certain vegetative parts occur. Duration of the individual growing stages of oats depends on a number of factors: oats variety, soil fertility, climate and agrotechnique.

In our trials, the following growing stages were registered: emergence, tillering, booting, tasseling and maturity. The results are shown in Table 1.

From the results in the Table 1, could be seen that the sawing of oats in 2015 was made earlier compared with 2016. The reason was frequent rainfall that in turn allowed moist medium for the seed and immediate germination. The yield of oats grain is the most important quantitative feature of oats controlled by the influence of a number of genes with less individual impact, to whose expression largely affects external factors i.e. the term of sowing (Jukic et al., 2011). The period from sowing to emerging, in both years of the trials and all the tested variants, was 10 or 11 days and is coincided with the results of the surveys of (Spasova et al.2013), in which she examines the impact of the farming systems on the vegetative growth and reproductive development of oats.

The tasseling phenophase, in both years of the trials, began in the first half of June. From the results in the Table 1 could be noticed that variety *Istra*, both in 2015 and 2016, began the earliest this phenophase, i.e. 01.06. In 2015, the phenophase tasseling began with in a period of 8 days, at all examined variants, while in 2016, for the same phenophase it took 15 days.

All 11 variants, the full maturity phenophase reached at the end of June, or in the beginning of July, within a period of 2 – 3 days. In 2015 three of the variants reached the full maturity stage on 31. 06., while in 2016, the first signs of this phenophase are registered on 04.07. Up to 02.07. 2015 and 07.07.2016 all the variants have reached the full maturity. The yield, in the first year of examination was made on 03.07., and in 2016, on 07.07., i.e. when there were optimal conditions. Genotypes features and the climatic conditions, especially temperature and the humidity of the air, largely affect on the time of occurrence and duration of the vegetative stages.

Table 1. Phenological observations of oats per year in organic production

Variety / Population	Phenophases				
	Date of sowing	Date of emergency	Date of tasseling	Date of maturity	Date of harvest
2015					
<i>Krivogastani</i>	23.03	03.04	05.06	30.06	03.07
<i>Trebenista</i>	23.03	03.04	06.06	02.07	03.07
<i>Kuceviste</i>	23.03	03.04	08.06	02.07	03.07
<i>Rajac</i>	23.03	03.04	05.06	02.07	03.07
<i>Slavuj</i>	23.03	03.04	07.06	01.07	03.07
<i>Lovken</i>	23.03	03.04	07.06	30.06	03.07
<i>Kupa</i>	23.03	03.04	02.06	02.07	03.07
<i>Baranja</i>	23.03	03.04	05.06	01.07	03.07
<i>Eksplorer</i>	23.03	03.04	04.06	02.07	03.07
<i>Sampionka</i>	23.03	03.04	08.06	02.07	03.07
<i>Istra</i>	23.03	03.04	01.06	30.06	03.07
2016					
<i>Krivogastani</i>	28.03	07.04	15.06	07.07	07.07
<i>Trebenista</i>	28.03	07.04	08.06	04.07	07.07
<i>Kuceviste</i>	28.03	07.04	15.06	06.07	07.07
<i>Rajac</i>	28.03	07.04	04.06	04.07	07.07
<i>Slavuj</i>	28.03	07.04	13.06	07.07	07.07
<i>Lovken</i>	28.03	07.04	14.06	07.07	07.07
<i>Kupa</i>	28.03	07.04	08.06	06.07	07.07
<i>Baranja</i>	28.03	07.04	13.06	06.07	07.07
<i>Eksplorer</i>	28.03	07.04	02.06	04.07	07.07
<i>Sampionka</i>	28.03	07.04	14.06	05.07	07.07
<i>Istra</i>	28.03	07.04	01.06	04.07	07.07

2. Physical and morphological characteristics of the oats

The absolute mass is a mass of 1000 air dry grains expressed in grams. It is characteristic of the species and variety, and it could be different at the same variety, depending on the conditions of production.

The hectoliter mass means a mass of some seed in a volume of 100 liters or a mass of a one hectoliter expressed in kilograms. The hectoliter mass is a feature of the species and the variety, but it depends of the production conditions of the seed (agrotechnique, soil and climatic conditions).

The height of the plants depends mostly on the genotype, soil and climatic conditions and used agrotechnique. The height of the oats has great impact on the lodging of the plants. The higher stem is, so is more prone to lodging. The most suitable for production are varieties with low to medium high and firm stem (Spasova, 2008). In optimal climatic conditions the differences in the plant height at the different oats genotypes are minimal (4 – 5 cm, or 7%), (Georgieva, 1995).

These three features that are measure for the quality and yield of oats grain were analyzed in our biannual research, and the results are showed in Table 2.

Table 2. Absolute mass of the grain, hectoliter mass of the grain and plants height in oats in organic production by years

Variety/Population	Absolute mass of the grain (g)	Hectoliter mass of the grain (kg/hl)	Plants height (cm)
2015			
<i>Krivogastani</i>	12,3	26,80	116,7
<i>Trebenista</i>	24,7	32,00	88,1
<i>Kuceviste</i>	20,1	25,80	97,8
<i>Rajac</i>	17,8	23,66	95,7
<i>Slavuj</i>	20,5	29,30	94,6
<i>Lovken</i>	25,6	30,80	88,1
<i>Kupa</i>	26,4	34,00	78,3
<i>Baranja</i>	24,3	34,00	94,0
<i>Eksplorer</i>	27,7	33,50	87,0
<i>Sampionka</i>	22,0	34,40	90,8
<i>Istra</i>	34,6	42,10	87,4
LSD 0,05	2,77	1,80	6,17
0,01	5,50	3,50	8,80
2016			
<i>Krivogastani</i>	14,9	36,00	143,3
<i>Trebenista</i>	19,9	33,70	101,3
<i>Kuceviste</i>	20,0	31,90	103,1
<i>Rajac</i>	23,3	31,50	107,8
<i>Slavuj</i>	18,5	31,70	115,3
<i>Lovken</i>	19,3	32,30	100,1
<i>Kupa</i>	22,9	34,50	83,4
<i>Baranja</i>	16,0	31,90	139,5
<i>Eksplorer</i>	19,3	32,40	100,4
<i>Sampionka</i>	17,6	32,80	117,6
<i>Istra</i>	29,6	35,60	107,5
LSD 0,05	2,65	1,41	30,30
0,01	3,78	2,01	/

The results from our biannual research of oats organic production showed that, regardless the year of examination, the highest absolute mass of the grain has variety *Istra* (34,6g in 2015 and 29,6g in 2016). The lowest absolute mass has population *Krivogastani* (12,3g in 2015 and 14,9g in 2016). There is significant statistical variation between the varieties. The studies of other researchers present increasing of the absolute mass, at different varieties (Đekić et al., 2010).

The highest hectoliter mass in oats organic production, in 2015, has variety *Istra* (42,10kg/hl), and in 2016 it was population *Krivogastani* (36 kg/hl). The lowest hectoliter mass, in both years of examination reached variety *Rajac* (23,66 kg/hl in 2015 and 31,50 kg/hl in 2016). Our results does not coincide with the results of *Cnacosa*

(2008), in which the variety *Rajac*, grown in conditions of organic production reached the highest average hectoliter mass (37,88 kg/hl), and population *Krivogastani* had the lowest average hectoliter mass (33,61 kg/hl).

The oats plants height (Table 2) in both years of research is statistically different at different varieties. In both years of research the highest are plants of population *Krivogastani* (116,7 cm in 2015 and 143,3 cm in 2016). The lowest height in both years of research has variety *Kupa* (78,35cm in 2015 and 83,375 cm in 2016).

The difference in the plants height, compared by years, using the same agrotechnique, at the same genotypes, is due to the variety specificity, i.e. the specificity of the genetic characteristics that possess tested varieties and population (Spasova, 2008). The results of our research coincide with the results of the respected author.

Conclusion

Based on our biannual research on some morphological and physical properties of oats, in terms of organic production the following conclusions can be drawn:

The phenological surveys in both years of the research showed that the vegetation period of the spring oats genotypes tested in organic production is 100 – 110 days.

Among the tested varieties we can mention the Croatian variety *Istra* which entered the earliest into all phenophases of growth and development.

In both years of research the highest absolute mass of the grain has variety *Istra* (34,6g in 2015 and 29,6g in 2016). The lowest absolute mass has population *Krivogastani* (12,3g in 2015 and 14,9g in 2016). Consequently, we can expect that variety *Istra* to achieve the highest yield and quality.

The lowest hectoliter mass, in both years of examination reached variety *Rajac* (23,66 kg/hl in 2015 and 31,50 kg/hl in 2016). The highest hectoliter mass in oats organic production, in 2015, has variety *Istra* (42,10kg/hl), and in 2016 it was population *Krivogastani* (36 kg/hl).

In both years of research the highest are plants of population *Krivogastani* (116,7 cm in 2015 and 143,3 cm in 2016). The lowest height in both years of research has variety *Kupa* (78,35cm in 2015 and 83,375 cm in 2016). Compared between varieties, there are statistically significant differences.

Acknowledgment

The research presented in this article is part of project *Organic production of spring oats: required variety features, No. 0307-252/4 from 18.02.2015* financially supported by "Goce Delcev" University – Stip, Republic of Macedonia

References

Georgieva T. (1995): Проучване основните звена от технологията на отглеждане на зимуващ овес. Автограферат на дисертация за получаване на научната степен "Кандидат на селскостопанските науки", Висш селскостопански Институт - Пловдив, катедра Растениевъдство

- Galie Z., Bleidere M., Skrabule I., Vigovskis J. (2004): The first steps in variety testing for organic agriculture in Latvia: oats and potatoes. Proceedings of the First World Conference on Organic Seed. Challenges and Opportunities for Organic Agriculture and the Seed Industry, FAO Headquarters, Rome, 173-174, Italy.
- Zorovski P., Georgieva T., Savova T., Gotcheva V., Spasova D., (2014): Productivity potential of wintering oat genotypes (*Avena sativa* L.) under the south Bulgaria agroecological conditions. Journal of Mountain Agriculture on the Balkans, 17 (1). pp. 80-92. ISSN 1311-0489
- Ilievski M., Spasova D., Mihajlov Lj., Markova Ruzdik N., Ilieva V., Sofijanova E. (2015): Production and balance among of cereal plants in Republic of Macedonia. Yearbook - Faculty of Agriculture, 13 (1). pp. 129-138. ISSN 1857-8608.
- Jukić G., Varnica I., Šunjić K., Mijić Z., Beraković I. (2011): Utjecaj roka sjetve na prinos kultivara jare zobi. *Sjemenarstvo* 2011. Vol. 28 No. 1/2 pp.17-23 ref.9
- Cooperman F.M., Dvorjankin F.A., Rostovceva Z.P., Rzanova E.I. (1955): Этапы формирования органов плодоношения злаков. Издательство Московского Университета
- Popov N., Georgieva T. (2009): Study of the effects of treatment with growth regulators on quality characteristics of oats, Journal of Mountain Agriculture on the Balkans, vol. 12, 5 (990-1002), Troyan
- Savova T., Panajotova G., Georgieva T. (2005): Оценка на изходен материалот овес във връзка със селекцията по качество на зърното. Field Crops Studies, Vol. II-2, 209-215.
- Spasova D. (2008): Сортна специфичност на овесот во услови на органско и конвенционално производство. Докторска дисертација. Универзитет „Св. Кирил и Методиј“ – Скопје. Факултет за земјоделски науки и храна.
- Spasova D., Spasov D., Atanasova B., Ilievski M., Kukutanov R., Georgieva T. (2013): Impact of the system of cultivation on the vegetative growth and reproductive development of oats. Bulgarian Journal of Agricultural Science, 19 (No 5) 2013, 1047-1055. Agricultural Academy
- Sutie J.M. (2000): Haycrops legumes and pulses. In: Hayand Straw Conservation For Small Scale Farming and Pastoral Conditions. FAO Plant Production and Protection Series No. 29, FAO, Rome
- Đekić V., Glamočlija Đ., Milovanović M., Staletić M. (2010): Uticaj godine na prinos i kvalitet zrna kragujevačkih sorti ozime pšenice. Zbornik Naučnih radova XXIV Svetovanja agronoma, veterinara i tehnologa, 24-25.februar, Beograd, Vol. 16, br. 1-2, 43-50.