

The European branch of the Special Interest Group for Drug Utilization Research (SIGDUR)

### EUROPEAN DRUG UTILISATION RESEARCH GROUP CONFERENCE 2017

Patients, Medicines, Bytes: Drug Utilisation Research and E-health



# **ABSTRACT BOOK**

Wednesday 15th - Friday 17th November 2017 Technology & Innovation Centre University of Strathclyde, Glasgow UK







The European branch of the Special Interest Group for Drug Utilization Research (SIGDUR)

### **Table of Contents**

### Oral presentations

P1W	Antimicrobial Use and Resistance	4
P2W	ePrescribing and eDispensing Systems	8
P3W	Adherence	9
P4T	Prescribing and de-Prescribing in Old Age	13
P5T	Gender Difference and Safety in DU	17
P6T	Decision Support to Improve Drug Use	19
P8T	Drug Use in Pregnancy and Pediatrics	23
P9T	Evolving Methods in DUR	25
P10F	Inform and Evaluate Health Policy	31
P11F	Citizen Science and DUR	35
P12F	Cardiovascular Therapy	39
P13F	Patient Reported Outcome Measures	45
P14F	New Medicines: Management of Innovations	49
P15F	DUR Across the World	53

### Poster presentations

1.	Antibiotics A	60
2.	Cardiovascular A	70
3.	Cancer and Biologicals A	80
4.	Drug Policy A	89
5.	Elderly A	95
6.	Nervous System A	104
7.	Diabetes	111
8.	Interventions	119
9.	Respiratory and Safety	128
11.	Antibiotics B	138
12.	Cardiovascular B	147
13.	Cancer and Biologicals B	155
14.	Drug Policy B	163
15.	Elderly B	169
16.	Nervous System B	176
17.	Adherence	184
18.	Pregnancy and Pediatrics	189
19.	Surprising others	198

### P1W.4

## Impact of the national intervention programme on parental knowledge, attitudes and practice of antibiotic use for respiratory infections

<u>Dr Verica Ivanovska<sup>1,2</sup></u>, Prof Bistra Angelovska<sup>1</sup>, Dr Liset van Dijk<sup>3</sup>, Prof Hubert Leufkens<sup>2,4</sup>, Prof Aukje Mantel-Teeuwisse<sup>2</sup>

<sup>1</sup>Faculty of Medical Sciences, Stip , Macedonia <sup>2</sup>Utrecht University Netherlands, Utrecht, The Netherlands <sup>3</sup>Netherlands Institute for Health Services Research (NIVEL), Utrecht, The Netherlands <sup>4</sup>Medicines Evaluation Board, Utrecht, The Netherlands

### Background

Children are prone to high rates of antibiotic use due to greater frequency of respiratory tract infections, and concerns about possible complications. Recent studies raised awareness of high resistance levels and inappropriate antibiotic use, including OTC sales of antibiotics across South-Eastern Europe. Nation-wide multifaceted interventions to improve antibiotic use were undertaken in FYR Macedonia in September 2014. This study aimed to assess the parental knowledge and attitudes about antibiotics, and self-medication practices in children, and evaluate the impact of interventions on these parameters.

### Methods

Pre-post intervention surveys were conducted in May 2014, May 2015 and May 2016 in three administrative regions in the country. Data were collected by interviewing parents of children younger than 15 years of age through a questionnaire. The analysis of knowledge, attitudes and antibiotic use involved descriptive quantitative statistics. The effects of interventions were assessed by a logistic and linear regression analysis.

#### Results

Data from 1203 interviewees showed that 80% of parents knew that antibiotics could kill bacteria, while 40% believed antibiotics could kill viruses. One third of parents expressed potential dissatisfaction with doctors who would not agree with them on antibiotic use. More parents received information about not taking antibiotics unnecessary after the interventions, but the rates decreased one year later. At baseline, 20% of the parents and 10% of the children who received antibiotics in previous year, took them without prescriptions. Parental self-medication rates did not change over time, while children rates decreased only in 2015.

### Conclusion

The insignificant and short-term effects of the interventions demonstrate that interventions need to be implemented for a longer period, at a large scale, with active health providers' engagement, and accompanied by inspections to promote appropriate antibiotic use and discourage self-medication. Future research should combine other methods (observations and pharmacy sales data studies) to validate self-reported data.