

BIOACCUMULATION AND TRANSLOCATION FACTOR OF HEAVY METALS IN THE PLANTS *LINARIA* SP., *MORICANDIA* SP. AND *VIOLA LUTEA* HUDS FROM THE ALŠAR LOCALITY – REPUBLIC OF MACEDONIA

Tena Šijakova-Ivanova¹, Blažo Boev¹, Vesna Zajkova-Paneva², Ivan Boev¹, Elizabeta Karakaševa²

¹*Faculty of Natural and Technical Sciences, Institute of Geology, “Goce Delčev” University in Štip,
Blvd. Goce Delčev 89, 2000 Štip, Republic of Macedonia*

²*Faculty of Agriculture, “Goce Delčev” University in Štip,
Blvd. Krste Misirkov 10-A, 2000 Štip, Republic of Macedonia*

tena.ivanova@ud.edu.mk

Abstract: This work was conducted in order to assess the distribution of metal and metalloids in soil and plants (*Linaria* sp., *Moricandia* sp. and *Viola lutea* Huds) which appears at the Alšar locality – Republic of Macedonia, as well as bioaccumulation and translocation of the metals in different parts of the plant. For this purpose, the contents of metals in soil and parts (root, stem, leaf and flowers) of the plants were digested and then analyzed by ICP-MS. Obtained results showed that studied soil and plants are enriched with thallium and arsenic. The contents of these elements in soil exceeded the intervention value for soils (The New Dutch list). According to New Dutch list, natural background content of Tl is 1.0 mg/kg, while interventional value is 15 mg/kg. In this work, the total Tl content in the soil from Alšar site is 561 mg/kg. The content of arsenic in unpolluted soils presented in other papers ranges from 4 to 150 mg/kg. Elevated As content, up to 732 mg/kg, is reported for Slovakian soils. Natural background content for As is 29 mg/kg, and intervention value is 55 mg/kg reported by the New Dutch list. In soil from Alšar content of As is 5376 mg/kg. Bioconcentration factor (BCF), bioaccumulation coefficient (BAC) and translocation factor (TF) also were determined. *Linaria* sp. has the highest TF for Tl, Rb and Cs. TF for Tl is: stem/root – 1.39, leaf/root – 7.54, and flower/root – 1.52. TF for Rb and Cs is 1.0 and 4.4 leaf/root, respectively. BAC has the highest value for Tl stem/soil – 10.19, leaf/soil – 55.45 and flower/soil – 11.16. BCF is the highest for Tl root/soil – 7.35. *Moricandia* sp. has TF values > 1, in leaf/root for Cr (1.44), Co (1.01), As (1.23), Ni (1.0), Rb (2.0), Cs (1.1), Ba (1.0) and Cd (1.2), flower/root for Rb (3.8), Cs (4.5), Cu (1.3) and Cd (1.7). *Viola lutea* Huds has TF values > 1 in stem/root for Cu (1.57), Zn (1.5), Tl (1.47), Mo (1.1), Sb (2.42) and Cd (1.3), and in leaf/root for Cr (3.04), Co (3.87), Ni (5.86), Zn (1.23), Pb (7.09), Tl (10.63), As (6.21) and Mo (4.48). *Moricandia* sp. and *Viola lutea* Huds for BAC and BCF have values < 1. Considering the BCF, BAC and TF values for *Linaria* sp., this plant for Tl, Cs and Rb possessed the characteristics of hyperaccumulator. Because *Linaria* sp., *Moricandia* sp. and *Viola lutea* Huds from Alšar locality have high TF values, could be useful for phytostabilization of contaminated soils.

Key words: arsenic; thallium; *Linaria* sp.; *Moricandia* sp.; *Viola lutea* Huds; BAC; TF; BCF

INTRODUCTION

Metals are present in the soil in different chemical forms, which influence their reactivity and hence their mobility. The impact of these metals in soils is their possible transfer into water or plants, which is defined by the term “bioavailability”. On the other hand, specific properties of plants are very significant in determining the bioavailability of trace elements and are quite variable with changing soil and plant conditions. Different genotypes of the same species uptake different amounts of the trace elements. Many plants that

accumulate >1000 or >10000 mg kg⁻¹ of trace metals are categorized as metals hyperaccumulators (Ma et al., 2001; Rosselli et al., 2003; Yoon et al., 2006; Sinegani et al., 2007). Accumulation of heavy metals varied greatly among plants species. Uptake of elements by a plant is primarily dependent on the plant species, its inherent controls, and the soil quality. The metal and plant interact in a specific way, which depends on several factors such as soil type, plant, growth conditions and the presence of other ions. (Baker and Brooks, 1989).

At present, there are four standards to judge hyperaccumulators, including the threshold value standard (the metal concentration in dry shoots >1000 mg/kg for As), the bioconcentrations factor standard (the ratio of metal content in a plant to soil is greater than 1.0, sometimes up to 50–100), the translocation factor standard (the ratio of metal concentration in aerial parts of plant to roots is greater than 1.0) and the strong metal-endurance

standard (high capability to tolerate the toxicity of metals) (Sun Yuebing, 2007). Therefore, the purpose of this paper was to examine some plants (*Linnaria* sp., *Moricandia* sp. and *Viola lutea* Huds) which appears at the Alšar locality – Republic of Macedonia, like potentia hyperaccumulator plants.

Hyperaccumulator plants can play a key role in the fate of the pollutants of contaminated matrices via their root systems.

MATERIAL AND METHODS

Description of the sampling site

Alšar locality is situated in the Vardar zone near the Macedonian–Greek border. From the geological point of view, the surroundings of the Alšar locality consist of several geological formations arranged in five stratigraphic complexes: complex of Precambrian metamorphic rocks; complex of Mesozoic (Triassic–Jurassic) rocks; complex of Upper Cretaceous sediment rocks; complex of Pliocene sediments, pyroclastites and volcanic rocks; complex of Quaternary sediments. The locality Alšar is a unique deposit with a number of minerals of thallium which are sulphates of thallium with arsenic, antimony, iron and mercury. As a result of this mineralization the entire space of a few square kilometers is enriched with arsenic, thallium and antimony. Also there are increased concentrations of these elements in the soils, water and plants.

The mine mineralization has been presented with As–Tl minerals, Fe-sulfides, Sb and Pb sulphates, cinabarite, sulfur and gold (Janković 1997; Boev 1993).

Sample collection and preparation

Soil and plant samples were collected from three sampling sites, where this community is present is the nearby of the village of Majden, on the locations of Alšar. All plant samples were collected in three species, taking a representative sample of root, tree, leaf and flowers and stored in polyethylene bags in the field, and transferred to the laboratory as soon as possible for analysis. Root, tree, leaf and flowers of plants were thoroughly washed with tap water to remove dust and other particles, followed by Mili-Q water. The samples were dried in an oven for 48 h at 80°C, and finally ground to powder for chemical analysis.

Soil and plant analysis

The <500 µm fraction soil samples were used to determine the maximal environmentally available trace element in soil. Soil samples were digested with a mixture of HCl (37%) and HNO₃ (70%), in a ratio of 3:1 (v/v), at room temperature for 16 h and after at 130 °C for 2 h, under reflux conditions (ISO 11466).

Plant samples were digested in 5 ml acid mixture of HClO₄ (Merck) (60%) and HNO₃ (Merck) (85%, s.g. 1.42) (15:85 v/v) overnight. Digestion was completed by gradual increase of temperature from 60 to 195 °C (60 °C, 3 h; 100 °C, 1 h; 120 °C, 1 h; 150 °C, 30 min; 175 °C, 30 min; 195 °C, 2.5 h). After cooling, HCl (20%, 2.5 ml) was added, whirl mixed and warmed to 80°C for 30 min. The final volume was brought to 10 ml with double distilled water and re-warmed to 80°C for another 30 min. Samples were filtered through cellulose filter (0.2 µM) (Barazani et al., 2004). Standard materials were included for assurance control.

The soil and plant digested solutions were cooled to room temperature, filtered, transferred quantitatively to 50 and 25 ml volumetric flasks, respectively, made up to volume with distilled water, and kept in clean plastic vials before metal analysis. Trace metal content in soil and plant extracts were determined by atomic emission spectrometry with ICP-MS (Inductive Coupled Plasma Mass Spectrometry).

All chemicals were declared proanalysis, and all solutions were prepared with double-distilled water. Standard working solutions were prepared from original certified stock solutions (MERCK), concentration 1000 mg/l in 1% (v/v) HNO₃. All samples were analyzed at the Analytical laboratory of the University “Goce Delčev” in Štip. The qual-

ity control procedure consisted of reagent blanks, replicated samples and certified standard material (PS-3, COOMET № 0001-1999 BG, COD № 310a

98). (Zapranova et al., 2006) The average values and standard deviations (SD) were calculated using Microsoft Office Excel 2003 in a computer.

RESULTS AND DISCUSSION

According to the New Dutch list (New Dutch list, 2012), natural background content of Tl is 1.0 mg/kg, while interventional value is 15 mg/kg. The total content of Tl in uncontaminated soils from different countries is reported as follows: <0.04–0.52 mg/kg in Sweden (Erikson, 2001); 0.01–0.41 in Poland (Lukaszewicz et al., 1996); 0.39–0.59 in Japan (Takeda et al., 2004). In this work, the total Tl content in the soil from Alšar site is 561 mg/kg. The content of arsenic in unpolluted soils presented in other papers ranges from 4 to 150 mg/kg (Wang and Mulligan, 2006). Elevated As content, up to 732 mg/kg, is reported for Slovakian soils from some locations (Čurlík and Šefčík, 1999). Natural background content for As is 29 mg/kg, and intervention value is 55 mg/kg reported by the New Dutch list (Figure 1). In soil from Alšar content of As is 5376 mg/kg (Table 1).

Zn content of natural sources is at the range of 126–683 mg/kg (Davies et al., 2003). The content, of Zn in non-contaminated soils reported for several European countries varies between 7 and 89 mg/kg (Kabata-Pendias, 2001). In this work, the total Zn content in the soil from Alšar locality is 57.13 mg/kg.

The copper content of world soils ranges from 2 to 100 mg/kg with an average value of about 30 mg/kg. Most of this is in unavailable mineral form. According the New Dutch list, natural background content of copper in nature is 36 mg/kg, while interventional value is 190 mg/kg (Figure 1). Normal copper concentration in plants is within the range from 2 to 50 mg/kg (Kabata Pendias, 2011). In the sampling point of Alšar locality, the total Cu content is 28.1 mg/kg (Table 1).

The reported Ba average range for soils on the world scale is from 362 to 580 mg/kg (Faucon et al., 2007). Monitoring on 218 surface-soil samples from Poland gave the Ba range of 20–130 mg/kg (Terelak et al., 2002). Swedish soils contain Ba in the range from 383 to 778 mg/kg (Erikson, 2001). In soil from Alšar site content of Ba is 583 mg/kg (Table 1).

Rubidium content in soils is largely inherited from the parent rocks, as is indicated by the highest mean Rb content, 100–120 mg/kg, in soils over granites and gneisses. The lowest contents of Rb, 30–50 mg/kg, are in sandy and organic soils, 1–10 mg/kg (Market and Lieth, 1987). In soil from Alšar side the mean content of Rb is 95 mg/kg.

The Co content in soils is inherited mainly from parent materials. The worldwide mean value of Co in surface soils is calculated as 10 mg/kg. Higher Co contents in surface soils are found in arid and semiarid regions; for example, Egyptian soils contain Co from 16.5 to 26.8 mg/kg (Nassem and Abdalla, 2003). Swedish arable soils contain Co in the range of 0.4–14 mg/kg, at the average value of 7.1 mg/kg (Kabata-Pendias, 2011).

Naturally high Co contents are observed in soils over serpentine rocks, up to 520 mg/kg, and in soils around ore deposits, up to 85 mg/kg (Kabata-Pendias, 2011). Content of Co in some parts of the Alšar locality is 15 mg/kg (Table 1).

Table 1

Content of metals and metalloids in soil collected from Alšar locality

Mean content \pm SD, $n = 3$			
M (soil)	mg/kg	M (soil)	mg/kg
Na	7175 \pm 123	Co	15 \pm 2.1
Mg	26650 \pm 152	Ni	41 \pm 7.8
Al	54577 \pm 165	Cu	28 \pm 1.7
P	1354 \pm 87	Zn	57 \pm 4.4
K	21714 \pm 221	As	5376 \pm 114
Ca	39562 \pm 188	Pb	15 \pm 6.1
Ti	2629 \pm 76	Rb	95 \pm 14
Mn	1813 \pm 45	Sr	223 \pm 16
Fe	28626 \pm 112	Mo	10.2 \pm 4.0
Cr	55 \pm 12	Sb	8.3 \pm 2.7
Ba	583 \pm 43	Cs	34.1 \pm 5.8
Tl	561 \pm 49	Cd	1.0 \pm 0.7
V	57 \pm 8.0	Sn	9.0 \pm 2.4

The average natural level of zinc in the earth's crust is 70 mg/kg (dry weight), ranging between 10 and 300 mg/kg (Singh, 2005). Elevated

Content of Ni in soil from Alšar is 41 mg/kg. The natural background content of Ni in soil according to the New Dutch list is 35 mg/kg. Content of Pb in soil from Alšar is 15 mg/kg, and well be-

low natural background content is 85 mg/kg. The mean values for V, Cd, Sn, Cr, As, Pb, Co, Ni, Cu, Zn, Mo, Ba and Tl in soils (CS) in part of the Alšar site are shown on Figure 1.




Fig. 1. Mean values for V, Cd, Sn, Cr, As, Pb, Co, Ni, Cu, Zn, Mo, Ba and Tl in soils (CS) in part of the Alšar locality (ICP-MS method, in mg/kg), and intervention values (IV) in agricultural soils in some European countries (The New Dutch list)

*Distribution of metals and metalloids in different parts of the plants *Linaria sp.*, *Moricandia sp.* and *Viola lutea* Huds from the Alšar locality*

The uptake of metals from the soil depends on different factors such as their soluble content, soil pH, plant growth stages, types of species, etc. (Mench et al., 1994; Hammer and Keller, 2002; Marschner, 2002).

Different plant parts accumulate different levels of these metals. For example, higher metal concentrations were observed in the edible and inedible parts of vegetable species as reported by many authors (Overesch et al., 2007; Alloway, 2012; Ismail et al., 2014).

The specific properties of plants are very significant in determining the bioavailability of trace elements and are quite variable with changing soil and plant conditions.

Images from investigated plants (*Linaria sp.*, *Moricandia sp.* and *Viola lutea* Huds) are given in Figs. 2, 3 and 4, respectively.



Fig. 2. *Linaria sp.* (K1) from Alšar locality



Fig. 3. *Moricandia sp.* (K2) from Alšar locality




Fig. 4. *Viola lutea* Huds (K3) from Alšar locality

The total contents of the analyzed elements in the soil were analyzed in order to characterize possible soil pollution and to determine the transfer efficiency from the soil.

The contents of different elements in plants are highly associated with the chemical composition of growth media and plant species. Plant uptake of heavy metals from soil occurs either passively with the mass flow of water into the roots, or through active transport and crosses the plasma

membrane of root epidermal cells. Under normal growing conditions, plants can potentially accumulate some metal ions in order of magnitude greater than the surrounding medium (Kim et al., 2003; Jung and Thornton 1996).

The total contents and distribution of metals and metalloids accumulated in investigation plants:

Linaria sp, *Moricandia* sp. and *Viola lutea* Huds, are shown in Table 2.

Correlations between the elements contents in the investigated plants were established using bivariate statistics. The significant correlations are marked in bold in Tables 3 and 4.

Table 2

The presence of metals and metalloids (mg/kg) in root, stem, leaf and flower of the plants Linaria sp. (K1), Moricandia sp. (K2) and Viola lutea Huds (K3) from Alšar locality

	Mean content \pm SD, $n = 3$											
	K1 root	K1 stem	K1 leaf	K1 flower	K2 root	K2 stem	K2 leaf	K2 flower	K3 root	K3 stem	K3 leaf	K3 flower
Na	101 ± 32	62.0 ± 21	66.0 ± 19	49.0 ± 12	92.0 ± 18	53.1 ± 13	82.1 ± 15	54.1 ± 13	21.0 ± 75	23.1 ± 5.0	33.0 ± 6.0	29.0 ± 5.0
Mg	2875 ± 65	1947 ± 58	3365 ± 76	1367 ± 79	585 ± 46	336 ± 28	1096 ± 54	1045 ± 38	291 ± 21	603 ± 32	1494 ± 38	700 ± 43
Al	3246 ± 57	376 ± 24	507 ± 36	308 ± 38	637 ± 42	163 ± 28	1104 ± 51	213 ± 28	196 ± 18	67 ± 15	760 ± 31	155 ± 21
P	882 ± 44	1215 ± 65	957 ± 33	1229 ± 45	730 ± 23	1238 ± 45	1865 ± 46	1817 ± 32	478 ± 23	1084 ± 65	945 ± 28	1592 ± 76
K	20518 ± 421	25522 ± 555	13346 ± 267	10581 ± 286	17510 ± 411	27210 ± 167	24214 ± 176	27388 ± 211	11916 ± 322	36576 ± 387	17909 ± 432	21487 ± 312
Ca	18540 ± 512	9439 ± 321	13769 ± 454	3731 ± 366	8791 ± 657	4305 ± 132	18934 ± 566	5716 ± 654	2764 ± 165	3254 ± 122	13641 ± 321	2584 ± 176
Ti	79.0 ± 16	36.0 ± 12	53.0 ± 13	15.0 ± 5.0	36.1 ± 11	17.1 ± 7.0	77.0 ± 16	23.0 ± 6.0	12.0 ± 4.0	13.0 ± 4.0	55.0 ± 14.0	11.0 ± 3.0
V	2.26 ± 0.44	0.3 ± 0.03	0.46 ± 0.05	0.26 ± 0.03	2.46 ± 0.04	0.1 ± 0.7	2.27 ± 0.7	0.17 ± 0.04	0.21 ± 0.06	0.06 ± 0.01	1.78 ± 0.05	0.17 ± 0.04
Sn	0.14 ± 0.03	0.11 ± 0.03	0.14 ± 0.02	0.08 ± 0.02	0.06 ± 0.03	0.08 ± 0.02	0.07 ± 0.02	0.06 ± 0.01	0.06 ± 0.01	0.11 ± 0.01	0.08 ± 0.02	0.08 ± 0.02
Cr	2.48 ± 0.34	0.76 ± 0.21	0.71 ± 0.12	0.47 ± 0.11	0.72 ± 0.07	0.25 ± 0.3	1.0 ± 0.1	0.37 ± 0.2	0.27 ± 0.1	0.15 ± 0.2	0.83 ± 0.2	0.29 ± 0.1
Co	0.71 ± 0.2	0.11 ± 0.02	0.22 ± 0.03	0.09 ± 0.01	0.29 ± 0.03	0.06 ± 0.01	0.3 ± 0.1	0.07 ± 0.02	0.13 ± 0.0	0.05 ± 0.01	0.52 ± 0.1	0.12 ± 0.02
Ni	8.0 ± 2.5	2.0 ± 0.5	2.0 ± 0.6	1.0 ± 0.2	2.0 ± 0.6	0.7 ± 0.4	2.0 ± 0.4	0.89 ± 0.14	0.85 ± 0.21	0.77 ± 0.22	5.0 ± 2.1	2.0 ± 0.8
Cu	3.0 ± 0.6	2.0 ± 0.4	2.0 ± 0.4	2.0 ± 0.3	3.0 ± 0.3	3.0 ± 0.6	1.0 ± 0.3	4.0 ± 1.4	0.44 ± 0.2	0.7 ± 0.4	0.32 ± 0.06	2.0 ± 0.04
Zn	13.0 ± 2.1	8.0 ± 1.6	10.0 ± 3.2	5.0 ± 0.6	8.0 ± 1.6	4.0 ± 2.1	7.0 ± 1.6	7.0 ± 1.7	2.0 ± 0.4	3.0 ± 0.3	3.0 ± 0.3	10.0 ± 2.1
As	471 ± 36	30.0 ± 14	75.0 ± 32	44.0 ± 12	190 ± 24	29.0 ± 13	219 ± 32	33.013 \pm	70.0 ± 16	17.0 ± 4.6	433 ± 23	62.0 ± 12
Pb	0.87 ± 0.04	0.22 ± 0.01	0.36 ± 0.04	0.21 ± 0.01	0.95 ± 0.02	0.13 ± 0.01	0.89 ± 0.02	0.11 ± 0.01	0.13 ± 0.01	0.08 ± 0.01	0.95 ± 0.02	0.16 ± 0.01
Rb	73 ± 13	110 ± 32	75.0 ± 16	83.0 ± 20	47.0 ± 15	82.0 ± 25	94.0 ± 21	178 ± 32	24.0 ± 9.0	35.0 ± 8.0	28.0 ± 5.0	34.0 ± 6.0
Sr	15 ± 2.0	7.0 ± 2.0	7.0 ± 2.0	1.0 ± 0.2	16.0 ± 2.4	4.0 ± 0.6	11.0 ± 2.3	2.0 ± 0.4	11.0 ± 1.2	12.0 ± 1.2	27.0 ± 2.1	3.007 \pm
Sb	0.33 ± 0.04	0.04 ± 0.01	0.07 ± 0.01	0.05 ± 0.01	0.1 ± 0.02	0.03 ± 0.01	0.08 ± 0.02	0.03 ± 0.01	0.07 ± 0.02	0.17 ± 0.04	0.05 ± 0.01	0.26 ± 0.02
Cs	5.0 ± 0.8	2.0 ± 0.2	22.0 ± 2.5	5.0 ± 0.6	2.0 ± 0.2	0.22 ± 0.03	2.0 ± 0.4	9.0 ± 0.7	12.0 ± 0.6	14.0 ± 0.6	3.0 ± 0.4	8.0 ± 1.2
Ba	109 ± 21	54.0 ± 12	17.0 ± 2.9	3.0 ± 0.7	60 ± 17	65 ± 21	60 ± 18	12.0 ± 2.0	6.0 ± 1.0	3.0 ± 0.6	8.0 ± 1.2	0.5 ± 0.06
Tl	4127 ± 64	5718 ± 121	31122 ± 413	6265 ± 254	184 ± 35	52 ± 12	91 ± 21	146 ± 32	185 ± 33	272 ± 43	1967 ± 65	99.0 ± 21
Cd	2.0 ± 0.4	0.13 ± 0.02	0.38 ± 0.03	1.0 ± 0.02	0.06 ± 0.01	1.0 ± 0.1	0.7 ± 0.1	0.1 ± 0.01	0.07 ± 0.01	0.09 ± 0.02	0.07 ± 0.02	0.04 ± 0.01

Table 3

Matrix of correlation coefficients between the elements contents in the investigated plants from Alšar locality

	Li	Be	B	Na	Mg	Al	Ca	Ti	V	Cr
Li	1.000									
Be	0.986	1.000								
B	0.560	0.463	1.000							
Na	0.698	0.720	0.653	1.000						
Mg	0.613	0.512	0.692	0.480	1.000					
Al	0.995	0.989	0.535	0.690	0.559	1.000				
Ca	0.754	0.719	0.599	0.713	0.659	0.750	1.000			
Ti	0.784	0.755	0.593	0.721	0.642	0.784	0.998	1.000		
V	0.669	0.738	0.199	0.714	0.208	0.689	0.773	0.794	1.000	
Cr	0.971	0.953	0.649	0.759	0.612	0.974	0.855	0.881	0.743	1.000
Mn	0.823	0.830	0.482	0.772	0.555	0.827	0.951	0.963	0.898	0.896
Fe	0.975	0.983	0.447	0.695	0.548	0.981	0.803	0.834	0.791	0.966
Co	0.876	0.892	0.249	0.536	0.514	0.883	0.790	0.816	0.811	0.870
Ni	0.897	0.878	0.374	0.446	0.610	0.895	0.686	0.714	0.594	0.862
Cu	0.211	0.230	0.431	0.645	0.274	0.203	0.075	0.074	0.057	0.203
Zn	0.643	0.614	0.728	0.684	0.654	0.625	0.502	0.512	0.303	0.653
As	0.791	0.819	0.128	0.444	0.379	0.811	0.745	0.772	0.820	0.799
Tl	0.102	0.008	0.387	0.179	0.780	0.032	0.295	0.256	-0.136	0.087
Pb	0.632	0.687	0.158	0.637	0.278	0.646	0.800	0.814	0.978	0.709

Table 4

Matrix of correlation coefficients between the elements contents in the investigated plants from Alšar locality

	Mn	Fe	Co	Ni	Cu	Zn	As	Tl	Pb
Mn	1.000								
Fe	0.891	1.000							
Co	0.874	0.951	1.000						
Ni	0.726	0.915	0.931	1.000					
Cu	0.105	0.150	-0.044	0.010	1.000				
Zn	0.513	0.574	0.417	0.545	0.593	1.000			
As	0.829	0.897	0.979	0.888	-0.131	0.276	1.000		
Tl	0.188	0.038	0.049	0.076	0.176	0.351	-0.086	1.000	
Pb	0.898	0.771	0.843	0.628	-0.025	0.253	0.863	-0.043	1.000

Tables 3 and 4 show significant correlations between: a) V, Ti, Cr, Mn, Fe, Co, Ni; b) Li, Be, Na, Al, Ca. Correlation coefficients between other elements are not significant. Basic statistics for

elements contents in the plants *Linaria* sp. (K1), *Moricandia* sp. (K2) and *Viola lutea* Huds (K3) from Alšar locality are given in Table 5.

Table 5

Basic statistics for elements contents in the plants *Linaria* sp. (K1), *Moricandia* sp. (K2) and *Viola lutea* Huds (K3) from the Alšar locality (contents are given in mg kg⁻¹ of dried matter)

	Mean	Median	Range	Minimum	Maximum
Li (lithium)	0.047	0.025	0.272	0.0003	0.272
Be (beryllium)	0.018	0.006	0.098	0.0003	0.098
B (boron)	780	695	755	432	1187
Na (natrium)	55	53	80	21	101
Mg (magnesium)	1309	1071	3074	291	3365
Al (aluminium)	644	342	3179	67	3246
P (phosphorus)	1169	1150	1387	478	1865
K (potassium)	21181	21003	25995	10581	36576
Ca (calcium)	8789	7254	16350	2584	18934
Ti (titanium)	36	29	68	11	79
V (vanadium)	0,88	0,28	2,407	0,058	2,46
Cr (chromium)	0.723	0.593	2.321	0.156	2.477
Mn (manganese)	39,06	23,79	82	10	92
Fe (iron)	284	133	1144	33	1178
Co (cobalt)	0.224	0.129	0.652	0.0554	0.707
Ni (nickel)	2.345	1.799	7.556	0.702	8.258
Cu (copper)	1.895	2.234	3.209	0.318	3.526
Zn (zinc)	6,46	6,972	10	2,181	13
Ga (gallium)	1.355	0.539	3.798	0.248	4.046
Ge (germanium)	0.024	0.013	0.084	0.003	0.088
As (arsenic)	140	66.0	454	17.3	471
Se (selenium)	0.009	0.008	0.02	0.002	0.018
Rb (rubidium)	72.0	73.8	154	23.8	178
Sr (strontium)	9.54	8.61	25.18	1.36	26.54
Mo (molybdenum)	0.428	0.325	0.764	0.164	0.928
Pd (palladium)	0.092	0.019	0.611	0.012	0.623
Ag (silver)	1.72	0.007	20.5	0.002	20.5
Cd (cadmium)	0.4	0.094	1.612	0.042	1.655
Sn (tin)	0.091	0.08	0.087	0.058	0.145
Sb (antimony)	0.107	0.071	0.301	0.028	0.329
Cs (caesium)	12.04	6.45	63	0.227	3
Ba (barium)	32.96	14.72	108	0.507	109
Tl (thallium)	4186	228	31069	52	31121
Pb (lead)	0.423	0.212	0.874	0.081	0.954
Bi (bismuth)	0.009	0.002	0.097	-0.005	0.093
Th (technicium)	0.077	0.02	0.494	0.005	0.499
U (uranium)	0.006	0.003	0.016	0.001	0.018

Unlike the macroelements whose distribution depends on the type of the plant, trace elements mainly are concentrated in the leaves and root of plants in all the investigated species of herbs. The contents of Cr, Co, Ni, Pb, Sr, Ba and Cd in selected plants did not differ significantly among the different species of plants (Table 2). In terms of metals distribution in the plants *Linaria* sp. and *Moricandia* sp. average contents of these elements decrease in the order: root > steam > leaf > flower, with the exception of the plant *Viola lutea* Huds where the contents of these elements in the leaf were higher than those in the root (Figure 5).

The highest content of As is measured in roots of *Linaria* sp. (471 mg/kg) and *Moricandia* sp. (190 mg/kg) and in leaf of *Viola lutea* Huds (433 mg/kg). The contents of Rb, Cs and As in plants differ for selected species and for parts of plants. Increased levels of Rb are reported for plants from industrial regions, 3–150 mg/kg (Kosla et al., 2001). Cs distribution in plants is similar to K, which suggests that this metal can compete with K uptake and K binding sites in cells. On the basis of a few compiled data, Cs common range in various plants can be presented as < 0.1–3 mg/kg (Isaure et al., 2006). In this work, the highest content of Cs and Rb is measured in leaf of *Viola lutea* Huds (22 mg/kg and 75 mg/kg) and in flower of *Moricandia* sp. (9 mg/kg and 178 mg/kg respectively).

Distribution of chemical elements in soils and stream sediments in the area of Alšar mine was determined by Bačeva et al., 2014. The Tl content in plants seems to be a function of Tl concentrations in soils. There is an assumption that some plants, especially of the Cruciferae and Gramineae families, can serve as hyperaccumulators for phytoremediation of contaminated soils. McGrath (1998) reported that there are programs for screening plants (e.g., *Brassica* sp.) for their ability to phytoextract Tl in plants from soils impacted by a cement factory. The content of Tl in plants from the Alšar area is very high (31.122 mg/kg), especially in *Linaria* sp. It decrease in order: leaf > flower > steam > root for *Linaria* sp., root > flower > leaf > stem for *Moricandia* sp., and leaf > stem > root > flower for *Viola lutea* Huds. The highest content of Tl is measured in all parts of *Linaria* sp.

The average contents of Zn decrease in the order: root > leaf > stem > flower for *Linaria* sp., root > leaf > flower > stem for *Moricandia* sp., and flower > stem > leaf > root for *Viola lutea* Huds. Zinc contents in plants vary considerably, reflecting the different factors of various ecosystems and of the genotypes (Kabata-Pendias, 2001).




Fig. 5. Content of metals and metalloids in soil and parts of the plant (root, stem, leaf and flower) in investigated plants *Linaria* sp. (K1), *Moricandia* sp. (K2) and *Viola lutea* Huds (K3) from the Alšar locality

In plants from Alšar, the content of zinc is equally distributed between the organs of the plants. Translocation factor (TF), bioaccumulation coefficient and bioconcentration factor were described. The translocation factor was described as ratio of heavy metals in aerial part of the plants to that in plant root (equation 1) (Cui et al., 2007; Li et al., 2007). Plant responses to TF in soils depend on several factors, however there are some general trends, expressed by plant/soil. Translocation factor (TF) can be presented as generalized values: 10 (Cd); 1 (B, Br, Cs, Rb); 10^{-1} (Ag, Co, Cu, Ge, Hg, Mo, Pb, Sr, Te, Zn); 10^{-2} (Be, As, Li, F, I, Mn, Ni, Sb); 10^{-3} (Ba, Bi, Ga, Fe, Se, V, Tl, Zr) (Kabata-Pendias, 2001). Translocation factor was calculated by:

$$TF = \text{metals (aerial part)} / \text{metals (roots)}, \quad (1)$$

where metals (aerial part) and metals (roots) are metals contents in the aerial part of the plant (mgkg^{-1}) and in the root of the plant (mgkg^{-1}), respectively. $TF > 1$ represents that translocation of

metals effectively was made to the shoot from root (Malik et al., 2010; Moffat, 1995; Yoon et al., 2006).

Biological accumulation coefficient (BAC) was calculated as ratio of heavy metals in aerial part of the plants to that in the soil (equation 2) (Cui et al., 2007; Li et al., 2007):

$$BAC = \text{metals (aerial part)} / \text{metals (soil)}. \quad (2)$$

Metals (aerial part) and metals (soil) are metals contents in the aerial part of the plant (mgkg^{-1}) and in the soil (mgkg^{-1}), respectively. $BAC > 1$ indicates that the plant species has the ability to store metals from the soil into the aerial part of the plant (Baker and Brooks, 1989; Baker et al., 2000).

A plant's ability to accumulate metals from soils can be estimated using the BCF (biological concentration factor), which is defined as the ratio of metals content in the roots of plant to that in the soil. It is calculated as follows (equation 3) (Yoon et al., 2006):

$$BCF = \text{metals (roots)} / \text{metals (soil)}. \quad (3)$$

BCF, BAC and TF values >1 had been used to evaluate the potential of plant species for phyto-extraction and phytostabilization of metals in soil. By comparing BCF, BAC and TF we can compare the ability of different plants in taking up metals from soils and translocating them to the aerial part of the plant. Bioaccumulation coefficient (BAC), concentration factor (BCF) and translocation factor (TF) of Cr, Co, Ni, Cu, Zn, As, Pb, Rb, Cs, Sr, Sb, Ba, Tl and Cd in K1 (*Linaria sp.*), K2 (*Moricandia*

sp.) and K3 (*Viola lutea* Huds) are given in Tables 6, 7 and 8 such as in Figures 6, 7 and 8, respectively.

Bioaccumulation of heavy metals in two endemic *Viola* species (*Viola allcharensis* G. Beck, *Viola arsenica* G. Beck) and one Balkan endemic species (*Viola macedonica* Boiss. & Heldr.) from the soil in the vicinity of the As-Sb-Tl mine Alšar, Republic of Macedonia, was determined by Bačeva et al., 2014. It was found that the accumulation of As, Sb, and Tl in these endemic species is significantly high.

Table 6

TF, BAC and BCF for *Linaria sp.* from the Alšar locality

	TF Translocation factor			BAC Bioaccumulation coefficient			BCF Bioconcentration factor
	steam/root	leaf/root	flower/root	steam/soil	leaf/soil	flower/soil	root/soil
Cr	0.307	0.288	0.191	0.014	0.013	0.009	0.045
Mn	0.295	0.585	0.212	0.015	0.030	0.011	0.051
Fe	0.121	0.215	0.104	0.005	0.009	0.004	0.041
Co	0.156	0.315	0.132	0.007	0.015	0.006	0.047
Ni	0.270	0.265	0.155	0.054	0.053	0.031	0.202
Cu	0.793	0.848	0.873	0.077	0.082	0.085	0.097
Zn	0.595	0.741	0.402	0.133	0.166	0.090	0.224
Pb	0.247	0.416	0.239	0.014	0.024	0.014	0.058
Tl	1.386	7.542	1.518	10.189	55.453	11.163	7.353
As	0.063	0.160	0.094	0.005	0.014	0.008	0.088
Mo	0.373	1.228	0.270	0.029	0.095	0.021	0.077
Sb	0.125	0.219	0.159	0.005	0.005	0.007	0.043
Rb	1.513	1.021	1.138	1.160	0.784	0.872	0.768
Cs	0.465	4.358	0.951	0.070	0.656	0.146	0.151
Sr	0.477	0.436	0.091	0.030	0.029	0.006	0.067
Cd	0.079	0.229	0.632	0.232	0.673	1.858	2.939




Fig. 6. TF, BAC and BCF for *Linaria sp.* from the Alšar locality

Table 7

TF, BAC and BCF for Moricandia sp. from the Alšar locality

	TF			BAC			BCF
	Translocation factor			Bioaccumulation coefficient			Bioconcentration factor
	steam/root	leaf/root	flower/root	steam/soil	leaf/soil	flower/soil	root/soil
Cr	0.341	1.883	0.503	0.005	0.025	0.007	0.013
Mn	0.281	1.424	0.358	0.009	0.045	0.011	0.031
Fe	0.191	1.227	0.257	0.003	0.016	0.003	0.013
Co	0.188	1.017	0.240	0.004	0.020	0.005	0.020
Ni	0.445	1.282	0.564	0.017	0.049	0.022	0.039
Cu	0.846	0.466	1.216	0.087	0.048	0.125	0.103
Zn	0.629	1.060	1.076	0.075	0.126	0.128	0.118
Pb	0.141	0.944	0.120	0.009	0.060	0.008	0.063
Tl	0.284	0.497	0.792	0.093	0.041	0.260	0.328
As	0.150	1.151	0.176	0.005	0.163	0.006	0.035
Mo	0.577	1.798	0.936	0.023	0.072	0.038	0.040
Sb	0.296	0.849	0.280	0.004	0.011	0.004	0.013
Rb	1.736	1.984	3.759	0.864	0.988	1.871	0.499
Cs	0.142	1.259	5.885	0.007	0.059	0.278	0.047
Sr	0.226	0.640	0.113	0.017	0.047	0.008	0.074
Cd	19.189	1.189	1.774	1.948	0.121	0.180	0.102



Fig 7. TF, BAC and BCF for *Moricandia* sp. from the Alšar locality

Table 8

TF, BAC and BCF for Viola lutea Huds from the Alšar locality

	TF			BAC			BCF
	Translocation factor			Bioaccumulation coefficient			Bioconcentration factor
	steam/root	leaf/root	flower/root	steam/soil	leaf/soil	flower/soil	root/soil
Cr	0.574	3.038	1.076	0.003	0.015	0.005	0.005
Mn	0.517	2.977	0.747	0.006	0.032	0.008	0.011
Fe	0.352	5.003	0.974	0.001	0.017	0.003	0.003
Co	0.429	3.871	0.924	0.004	0.035	0.008	0.009
Ni	0.912	5.860	2.776	0.019	0.122	0.058	0.021
Cu	1.568	0.715	3.317	0.025	0.011	0.052	0.016
Zn	1.155	1.232	4.371	0.044	0.047	0.167	0.038
Pb	0.598	7.095	1.175	0.005	0.064	0.011	0.009
Tl	1.468	10.629	0.529	0.484	3.506	0.174	0.330
As	0.248	6.213	0.893	0.003	0.081	0.012	0.013
Mo	1.105	4.476	1.161	0.019	0.075	0.020	0.017
Sb	2.421	0.769	3.692	0.022	0.007	0.034	0.009
Rb	1.482	1.187	1.412	0.370	0.296	0.352	0.250
Cs	1.124	5.108	0.634	0.408	1.855	0.230	0.363
Sr	1.045	2.321	0.216	0.054	0.119	0.011	0.051
Cd	1.304	1.093	0.643	0.152	0.128	0.075	0.117

Fig. 8. TF, BAC and BCF for *Viola lutea* Huds from the Alšar locality

The accumulating species are characterized by a ratio of content of metals in the aerial part of the plant to the content in the soil greater than 1 (BAC > 1); species known as "excluder" having a ratio below 1 (Baker and Whiting, 2002). This

agrees with Rotkittikhun et al. (2006), who suggested that accumulation occurs when the content of an element in the aerial part of a plant is higher than in the soil or when the content ratio of leaf/soil should be greater than 1 (BAC). This test

reveals the ability of a species to accumulate metals. Secondly, the content of an element in the aerial parts of a plant should be 10 to 500 times higher than normal levels, and thirdly, the content ratio leaves/roots must be greater than 1 ($TF > 1$). The BCF values in the range of 0.1 to 0.50 (as potential phytostabilization) and values from 0.5 to 1.0 (as partial phytostabilization) should not be neglected. The species that show values of TF and especially $BCF < 1$ are inadequate to the phytoextraction (Fitz and Wenzel, 2002). In this work *Linaria* sp. has the highest TF for Tl. TF for Tl is 1.39 (steam/root), 7.54 (leaf/root) and 1.52 (flower/root). BAC has the highest values for Tl – 10.19 (steam/soil), 55.45 (leaf/soil) and 11.16 (flower/soil); Rb 1.16 (steam/soil), Cd 1.86 (flower/soil). *Linaria* sp. has the highest BCF values for

Tl – 7.35 (root/soil) and Cd (2.94). *Moricandia* sp. has TF values > 1 in steam/root for Rb (1.74) and Cd (19.19); in leaf/root for Cr (1.88), Co (1.02), As (1.15), Ni (1.28), Rb (1.98), Mn (1.42), Fe (1.23), Zn (1.06), Cs (1.26), Mo (1.80), Cd (1.19); in flower/root for Rb (3.76), Cs (5.89), Cu (1.21), Zn (1.08) and Cd (1.77). BAC is > 1 for Rb (1.95) (steam soil); Cd (1.87) flower/soil. *Viola lutea* Huds has TF values > 1 in steam/root for Cu (1.57), Zn (1.16), Tl (1.47), Mo (1.11), Sb (2.42), Rb (1.48), Cs (1.12), Sr (1.05) and Cd (1.30); in leaf/root for Cr (3.04), Mn (2.98), Co (3.87), Ni (5.86), Zn (1.23), Pb (7.10), Tl (10.63), As (6.21), Mo (4.48), Rb (1.19), Cs (5.11), Sr (2.32) and Cd (1.09); in flower/root for Cr (1.08), Ni (2.78), Cu (3.32), Zn (4.37), Pb (1.18), Mo (1.16), Sb (3.69) and Rb (1.41).

CONCLUSION

After summarizing the data collected in this research, we can confirm that accumulation of selected metals varied among the plants *Linaria* sp., *Moricandia* sp., *Viola lutea* Huds and uptake of an element by a plant is primarily dependent on the plant species and the soil quality. Significant correlation is between: a) V, Ti, Cr, Mn, Fe, Co, Ni; b) Li, Be, Na, Al, Ca. Correlation coefficient between other elements is not significant. *Linaria* sp. (K1) has the highest content of elements in the root, while in *Moricandia* sp. (K2) and *Viola lutea* Huds (K3) the highest content of elements is in leaves. The highest values for bioaccumulation

factor were obtained in case of Tl in steam, leaf and flower from *Linaria* sp. and *Moricandia* sp. *Linaria* sp. has the highest TF for Tl, Cs and Rb. *Moricandia* sp. and *Viola lutea* Huds for BAC and BCF have values < 1 . Considering the BCF, BAC and TF values for *Linaria* sp. this plant for Tl, Cs and Rb possessed the characteristics of hyperaccumulator. Because *Linaria* sp., *Moricandia* sp. and *Viola lutea* Huds from the Alšar locality have high TF values for As, Tl, Sb, Ni, Cu, Zn, Cs, Rb, Co, Mo, Cr and Pb, could be useful for phytostabilization of soils contaminated with these elements.

REFERENCES

- [1] Alloway B. J. (2012): *Trace Metals and Metalloids in Soils and Their Bioavailability: Heavy Metals in Soils*. London, Springer Press.
- [2] Bačeva K., Stafilov T., Šajn R., Tănăselia C., Makreski P.. (2014): Distribution of chemical elements in soils and stream sediments in the area of abandoned Sb-As-Tl Allchar mine, Republic of Macedonia, *Multidisciplinary Journal of Environmental Sciences, Ecology and Public Health*, Doi: 10.1016/j.envres.2014.03.045
- [3] Baker A. J. M., Whiting S. N. (2002): In search of the Holy Grail – A further step in understanding metal hyperaccumulation. *New Phytologist*. **155**, 1–4.
- [4] Baker A. J. M. and Brooks R. R. (1989): Terrestrial higher plants which hyperaccumulate metallic elements – A review of their distribution, ecology and phytochemistry. *Biorecovery*, **1**, 81–126.
- [5] Baker A. J. M., McGrath S. P., Reeves R. D. and Smith J. A. C. (2000): Metal hyperaccumulator plants: A review of ecology and physiology of a biological resource for phytoremediation of metal-polluted soils. In: *Phytoremediation of Contaminated Soil and Water*. Eds.: N. Terry and others. Boca Raton, FL7 Lewis Publishers, pp. 129–58.
- [6] Barazani O., Dudai N., Khadka U. R., Golan-Goldhirsh A. (2004): Cadmium accumulation in *Allium schoenoprasum* L. grown in an aqueous medium. *Chemosphere* **57**, 1213–1218.
- [7] Boev B., Stojanov R., Denkovski G. (1993): Geology of the polymetallic deposit Alšar, Macedonia, *Geologica Macedonica*, Štip, Vol. 7, No. 1, pp. 35–39.
- [8] Chen Y., Shen Z., Li X. (2004): The use of vetiver grass (*Vetiveria zizanioides*) in the phytoremediation of soils contaminated with heavy metals. *Appl. Geochem.* **19**, 1553–1565.
- [9] Cui S., Zhou Q. and Chao L. (2007): Potential hyper-accumulation of Pb, Zn, Cu and Cd in enduring plants distributed in an old smeltery, northeast China. *Environmental Geology*, **51**, 1043–1048.

- [10] Čurlík J., Šeřík P. (1999): *Geochemical Atlas of the Slovak Republic. Soils*. Ministry of the Environment of the Slovak Republic.
- [11] Davies B. E., Vaughan J., Lalor G. C., Vutchkov M. (2003): Cadmium and zinc adsorption maxima of geochemically anomalous soils (Oxisols) in Jamaica. *Chem. Spec. Bioavailab.* **15**, 59–66.
- [12] Eriksson J. E. (2001): Concentrations of 61 trace elements in sewage sludge, farmyard manure, mineral fertilizers, precipitation in oil and crops. Swedish EPA. Rep 5159. Stockholm Health, eds. Y. Zhu, N. Lepp, R. Naidu, 281–282, Tsinghua Univ. Beijing
- [13] Faucon M. P., Shutcha M., Meerts P. (2007): Revisiting copper and cobalt concentrations in supposed hyperaccumulators from SC Africa: Influence of washing and metal concentrations in soil. *Plant Soil*, **301**, 29–36.
- [14] Fitz W. J. & Wenzel W. W. (2002): Arsenic transformation in the soil–rhizosphere–plant system, fundamentals and potential application of phytoremediation, *Journal of Biotechnology*, **99**, pp. 259–278.
- [15] Hammer D., Keller C. (2002): Changes in the rhizosphere of heavy metal accumulating plants as evidenced by chemical extractants. *J Environ Qual*, **31**, 1561–9.
- [16] Isaure M. P., Frayse A., Deves G. et al. (2006): Microchemical imaging of cesium distribution in *Arabidopsis thaliana* plant and its interaction with potassium and essential trace elements. *Biochemie*, **88**, 1538–1590.
- [17] Ismail A., Riaz M., Akhtar S., Ismail T., Amir M., Zafar-ul-Hye M. (2014): Heavy metals in vegetables and respective soils irrigated by canal, municipal waste and tube-well waters. *Food Addit Contam B*. DOI:10.1080/19393210.2014.88878.
- [18] ISO 11466 (1995): Soil quality – Extraction of trace elements soluble in aqua regia.
- [19] Janković et al. (1997). *Magmatizam i tercierna mineralizacija vo kožuvskata metalogenetska oblast vo Makedonija so poseben osvrt na naogališteto Alšar*, Rudarsko-geološki fakultet, Posebno izdanie, br. 5, Štip, 262 p.
- [20] Jung M. C. & Thornton I. (1996): Heavy metal contamination of soils and plants in the vicinity of a lead-zinc mine, Korea, *Applied Geochemistry*, **11**, pp. 53–59.
- [21] Kabata-Pendias A., Pendias H. (2001): *Trace elements in soils and plants*. 3rd edition. CRC Press, Boca Raton, Florida, 413 p.
- [22] Kabata-Pendias A., Pendias H. (2011): *Trace elements in soils and plants*. 4rd edition. CRC Press, Boca Raton, Florida, 548 p.
- [23] Kim I. S., Kang H. K., Johnson-Green P. & Lee E. J. (2003). Investigation of heavy metal accumulation in *Polygonum thunbergii* for phytoextraction, *Environmental Pollution*, **126**, pp. 235–243.
- [24] Kosla T., Skibniewska E., Debski B., Urbanska-Slomka G. (2002): Rubidium in the trophic chain soil-plants-animals. *Trace Elements and Electrolytes* **19** (4), 171–176.
- [25] Li M. S., Luo Y. P. and Su Z. Y. (2007): Heavy metal concentrations in soils and plant accumulation in a re-stored manganese mineland in Guangxi, South China. *Environmental Pollution*, **147**, 168–175.
- [26] Łukaszewicz Z., Zembruski W., Piela A. (1996): Direct determination of ultratrace of thallium by flow injection-differential-pulse anodic stripping voltammetry. *Anal. Chem.* **318**, 159–165.
- [27] Ma L.Q., Komar K. M., Tu C., Zhang W., Cai Y. and Kennelley E. D. (2001): A fern that hyperaccumulates arsenic. *Nature*, **409**, 579–579.
- [28] Malik R. N., Husain S. Z., Nazir I. (2010): Heavy metal contamination and accumulation in soil and wild plant species from industrial area of Islamabad, Pakistan. *Pak. J. Bot.*, **42** (1), 291–301.
- [29] Markert B. and Lieth H. (1987): Element concentration cadasters in a Swedish biotype, *Fresenius Z. Anal. Chem.*, **327**, 716.
- [30] Marschner H. (2002): *Mineral Nutrition of Higher Plants*. London, UK. Elsevier, Academic Press.
- [31] McGrath S. P. (1998): Phytoextraction for soil remediation, In: *Plants that Hyperaccumulate Heavy Metals*, R. R. Brooks, ed., CAB Intern., 261.
- [32] Mench M., Vangronsveld J., Didier V., Clijsters H. (1994): Evaluating of metal mobility, plant availability and immobilization by chemical agents in a limed-salty soil. *Environ Pollut.* **86**, 279–86.
- [33] Moffat A. S. (1995): Plants proving their worth in toxic metal clean up. *Science* (Washington, DC) **269**, 302–303.
- [34] Nasseem M. G., Abdalla Y. H. (2003): Cobalt status in the North-Western Coast soils of Egypt in relation to cobalt content of barley for ruminants. *16 Int. Symp. Environ. Biogeochem.* Edinburgh.
- [35] New Dutch list (2012): <http://www.contaminatedland.co.uk/std-guid/dutchl.htm>, last accessed May, 2012
- [36] Overesch M., Rinklebe J., Broll G., Neue H. U. (2007): Metals and arsenic in soils and corresponding vegetation at Central Elbe river flood plains (Germany). *Environ Pollut.* **145**, 800–12.
- [37] Zapranova, P., Angelova V., Dospatliev L., Ivanov K., (2006): Comparative study of methods for the preparation of soil samples for analysis AAS of Cd, Pb, Cr and Ni, *Scientific Papers*, **34**, 5.
- [38] Rosselli, W., C. Keller and K. Boschi (2003): Phytoextraction capacity of trees growing on metal contaminated soil. *Plant Soil*, **256**, 265–272.
- [39] Rotkittikhun P., Kruatrachue M., Chaiyarat R., Ngernsarsaruay C., Pokethitiyook P., Pajitpraporn A., Baker A. J. M. (2006): Uptake and accumulation of lead by plants from the Bo Ngam lead mine area in Thailand. *Environ. Pollut.* **144**, 681–688.
- [40] Sinegani A. A. S. and Ebrahimi P. (2007): The potential of Razan-Hamadan highway indigenous plant species for the phytoremediation of lead contaminated land. *Soil Environ.*, **26**, 10–14.
- [41] Singh, V. P. (2005): *Metal Toxicity and Tolerance in Plants and Animals*, Sarup & Sons, New Delhi, 328 pages.

- [42] SUN Yuebing, ZHOU Qixing, WEI Shuhel, REN Liping, (2007): Growth responses of the newly-discovered Cd-hyperaccumulator *Rorippa globosa* and its accumulation characteristics of Cd and As under joint stress of Cd and As, *Front. Environ. Sci. Engin. China*, **1** (1), 107–113.
- [43] Takeda A., Kimura K., Yamasaki S-I. (2004): Analysis of 57 elements in Japanese soils, with special reference to soil group, and agricultural use. *Geoderma*, **119**, 291–307.
- [44] Terelak H., Motowicka-Terelak T., Maliszewska-Kordybach B., Pietruch Cz. (2002): *Monitoring of Chemical Properties of Arable Soils in Poland*. Bibl. Monit. Środowiska, Warsaw (Po).
- [45] Wang S., Mulligan C. N. (2006): Occurrence of arsenic contamination in Canada: Sources, behavior and distribution, *Sci. Total Environ.* **366**, 701–721.
- [46] Yoon Y., Cao X., Zhou Q. & Ma L. Q. (2006): Accumulation of Pb, Cu, and Zn in native plants growing on a contaminated Florida site, *Science of Total Environment*, **368**, pp. 456–464.

Резиме

БИОАКУМУЛАЦИОНЕН И ТРАНСЛОКАЦИОНЕН ФАКТОР НА ТЕШКИ МЕТАЛИ ВО РАСТЕНИЈАТА *LINARIA SP.*, *MORICANDIA SP.* И *VIOLA LUTEA HUDS* ОД ЛОКАЛИТЕТОТ АЛШАР – РЕПУБЛИКА МАКЕДОНИЈА

Тена Шијакова-Иванова¹, Блажо Боев¹, Весна Зајкова-Панева², Иван Боев¹, Елизабета Каракашева²

¹Факултет за природни и технички науки, Институт за геологија, Универзитет „Гоце Делчев“ во Штип, бул. Гоце Делчев 89, 2000 Штип, Република Македонија

²Земјоделски факултет, Универзитет „Гоце Делчев“ во Штип, ул. Крсте Мисирков 10-А, 2000 Штип, Република Македонија
tena.ivanova@ud.edu.mk

Клучни зборови: арсен; талиум; *Linaria sp.*; *Moricandia sp.*; *Viola lutea* Huds; BAC; TF; BCF

Целта на овој труд е да се оцени дистрибуцијата на металите и металоидите во почвата и растенијата *Linaria sp.*, *Moricandia sp.* и *Viola lutea* Huds од локалитетот Алшар – Република Македонија. Врз основа на добиените резултати се одредени факторот на биоконцентрација (BCF), коефициентот на биоаккумуляција (BAC) и факторот на транслокација (TF) во различните делови на растенијата. Содржината на металите во почвата и во деловите на растенијата (корен, стебло, лист и цвет) беше одредена со ICP-MS. Врз основа на добиените резултати беше утврдено дека *Linaria sp.* има највисок TF за Tl, Rb и Cs. TF за Tl е следниот: стебло/корен – 1,39; лист/корен – 7,54 и цвет/корен – 1,52. TF за Rb е 1,0 (лист/корен) и за Cs е 4,4 (лист/корен). BAC има највисока вредност за Tl (стебло/почва – 10,19, лист/почва – 55,45 и цвет/почва – 11,16). BCF исто така е највисок за Tl (корен/почва – 7,35). *Moricandia sp.*

има вредностите на TF > 1 во лист/корен за Cr (1,44), Co (1,01), As (1,23), Ni (1,0), Rb (2,0), Cs (1,1), Ba (1,0) и Cd (1,2), во цвет/корен за Rb (3,8), Cs (4,5), Cu (1,3) и Cd (1,7). *Viola lutea* Huds вредностите на TF има > 1 во стебло/корен за Cu (1,57), Zn (1,5), Tl (1,47), Mo (1,1), Sb (2,42) и Cd (1,3). во лист/корен за Cr (3,04), Co (3,87), Ni (5,86), Zn (1,23), Pb (7,09), Tl (10,63), As (6,21) и Mo (4,48). *Moricandia sp.* и *Viola lutea* Huds за BAC и BCF имаат вредности < 1. Според вредностите на BCF, BAC и TF за *Linaria sp.*, ова растение за Tl, Cs и Rb ги поседува карактеристиките на хиперакумулатор. Испитувањата покажаа дека проучуваната почва и растенијата се богати со талиум и арсен. Поради тоа што *Linaria sp.*, *Moricandia sp.* и *Viola lutea* Huds, имаат високи вредности на TF, тие би можеле да бидат користени за фитостабилизација на почви контаминирани со овие елементи.