GOCE DELCEV UNIVERSITY - STIP FACULTY OF AGRICULTURE



JOURNAL OF AGRICULTURE AND PLANT SCIENCES

YEAR 2018

VOLUME 16, Number1

EDITOR IN CHIEF

Liljana Koleva Gudeva, Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, liljana.gudeva@ugd.edu.mk

Editors

Ljupco Mihajlov, Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, ljupco.mihajlov@ugd.edu.mk Fidanka Trajkova, Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, fidanka.trajkova@ugd.edu.mk

Administrator

Biljana Atanasova, Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, <u>biljana.atanasova@ugd.edu.mk</u>

Technical Editing

Slave Dimitrov, Publishing Department, Goce Delcev University, Stip, Macedonia, <u>slave.dimitrov@ugd.edu.mk</u>

Language Editors

Biljana Ivanova, MA, Senior lecturer, Faculty of Philology, Goce Delcev University, Stip, Macedonia, <u>biljana.petkovska@ugd.edu.mk</u> – English language Editor

Danica Gavrilovska Atanasovska, Goce Delcev University, Stip, Macedonia, <u>danica.gavrilovska@ugd.edu.mk</u> – Macedonian language Editor

Journal of Agriculture and Plant Sciences, JAPS, Vol 16, No. 1 Successor of the Yearbook of Faculty of Agriculture of GDU, Vol 14

Editorial Office Faculty of Agriculture, Goce Delcev University - Stip, Krste Misirkov Str., No.10-A, P.O. Box 201, 2000 Stip, Republic of Macedonia japs@ugd.edu.mk http://js.ugd.edu.mk/index.php/YFA

> ISSN 2545-4447 print ISSN 2545-4455 on line Vol. 16, No. 1, Year 2018

GOCE DELCEV UNIVERSITY - STIP, REPUBLIC OF MACEDONIA FACULTY OF AGRICULTURE

UDC 63(058) ISSN 2545-4447 print ISSN 2545-4455 on line



Journal of Agriculture and Plant Sciences, JAPS, Vol 16, No. 1

Successor of the Yearbook of Faculty of Agriculture of GDU, Vol 14

YEAR 2018

VOLUME XVI, Number 1

EDITORIAL BOARD

Aco Kuzelov,

Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, <u>aco.kuzelov@ugd.edu.mk</u> **Danijela Raičević,** Biotechical Faculty, University of Montenegro, Mihaila Lalica bb., Podgorica,

Montenegro, nelar@mail.com

Dragan Skorić,

Serbian Academy of Sciences and Arts, Knez Mihajlova 35, 11000 Belgrade,

Serbia, draganskoric@sbb.rs

Dragomir Vlcev,

Institute of Agriculture – Karnobat, Bulgaria, vulchevd@abv.bg

Hatice Gülen,

Istanbul Bilgi University, Faculty of Engineering and Natural Sciences,

Department of Genetics and Bioengineering, Istanbul, Turkey,

hatice.gulen@bilgi.edu.tr

Jovica Vasin,

Institute of Field and Vegetable Crops, Novi Sad, Serbia, jovica.vasin@ifvcns.ns.ac.rs

Kiril Bahcevandziev,

Coimbra Agricultural School, 3045-601 Coimbra, Portugal, kiril@esac.pt

Klemen Lisjak,

Agricultural Institute of Slovenia, Hacquetova ulica 17, Ljubljana, Slovenia, <u>Klemen.Lisjak@kis.si</u> **Marijan Bubola**,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, <u>marijan@iptpo.hr</u> Maryna Mardar,

Odessa National Academy of Food Technologies, Odessa, 65039, Kanatnaya Str., Ukraine, <u>marinamardar2003@gmail.com</u>

Rubin Gulaboski,

Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, <u>rubin.gulaboski.@ugd.edu.mk</u> **Sanja Radeka**,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, <u>sanja@iptpo.hr</u> Sasa Mitrev,

Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, <u>sasa.mitrev@ugd.edu.mk</u> **Shuhe Wei**,

Institute of Applied Ecology, Chinese Academy of Sciences, China, <u>shuhewei@iae.ac.cn</u> Velichka Nikolova Rodeva,

Maritsa Vegetable Crops Research Institute, 32 Brezovsko shosse St. Plovdiv 4003, Bulgaria, <u>velirod@yahoo.com</u>

Violeta Dimovska,

Faculty of Agriculture, Goce Delcev University, Stip, Macedonia, <u>violeta.dimovska@ugd.edu.mk</u> **Wolfram Schnäckel**,

Anhalt University of Applied Sciences, Bernburger Straße 55, 06366 Köthen, Germany, Wolfram.Schnaeckel@hs-anhalt.de

CONTENT

Jane Aleksoski THE EFFECT OF BACKCROSS METHOD IN TOBACCO BREEDING9
Kata Angelova, Verica Ilieva, Natalija Markova Ruzdik,
Ilija Karov, Ljupco Mihajlov, Mite Ilievski
EXAMINATION ON YIELD AND SOME YIELD ASSOCIATED PARAMETERS
IN DIFFERENT RICE GENOTYPES
Olivera Bicikliski, Fidanka Trajkova, Ljupco Mihajlov
EVALUATION OF THE CURRENT STATUS IN ORGANIC AGRICULTURAL PRODUCTION
IN REPUBLIC OF MACEDONIA AND EUROPEAN COUNTRIES
Lence Buseva, Dragica Spasova, Biljana Atanasova
CHARACTERIZATION OF SOME DOMESTIC AND INTRODUCED VARIETIES OF
COTTON IN THE AGRO-ECOLOGICAL CONDITIONS OF STRUMICA REGION
Valentina Butleska Gjoroska, Marija Krstik, Ivana Jovanovska Klincarska,
Ana Cvetanovska, Lenka Cvetanovska, Liljana Koleva Gudeva
EVALUATION OF TOTAL PHENOLS IN ALFALFA (Medicago sativa L.)
COLLECTED FROM DIFFERENT LOCALITIES IN REPUBLIC OF MACEDONIA
Stefan G. Dragoev, Dessislava B. Vlahova-Vangelova,
Dessislav K. Balev, Kolyo T. Dinkov, Aco Kuzelov
PECULIARITIES OF THE HORSE MEAT AGING
Maryna Mardar, Galina Krusir, Rafaela Znachek, Larisa Agunova
BIOASSAY IN SAFETY ASSESSMENT OF NEW GRAIN PRODUCTS
Vesna Markoska, Kiril Lisichkov, Blazo Boev, Rubin Gulaboski
THE INFLUENCE OF THE PERLITE AS A SUBSTRATE FOR IMPROVING ON SOME WATER
PROPERTIES ON THE FLUVIAL SOIL WITH AN APLICATION OF RETENTIONAL CURVES
Milan Mitreski, Jane Aleksoski, Ana Korubin - Aleksoska,
Marjan Trajkoski, Jordan Trajkoski
VARIATION OF MORPHOLOGICAL PROPERTIES IN VIRGINIA TOBACCO TYPES
Efremco Nikolov, Violeta Dimovska, Fidanka Ilieva
THE EFFECT OF PRUNING ON FRIUTING CAPACITY OF MICHELE PALIERI TABLE
GRAPE VARIETY GROWING IN TIKVES VINEYARD
Nenad Petkovski, Ljupco Mihajlov, Natalija Markova Ruzdik
GENETIC AND ENVIRONMENTAL EFFECT ON THE GRAIN YIELD OF
SPRING BARLEY VARIETIES CULTIVATED IN THE REPUBLIC OF MACEDONIA
Mitre Stojanovski, Anita Čakarova, Aco Kuzelov, Elena Joshevska,
Gordana Dimitrovska, Dzulijana Tomovska, Katerina Bojkovska
CHANGES IN FATTY ACID COMPOSITION OF POULTRY MEAT AFTER HEAT TREATMENT

INTORDUCTION

In the past ten years the educational, research and applicative activities of the Faculty of Agriculture – Stip, Goce Delcev University – Stip, contributed to the development of agriculture sector in the country and broader region.

The Faculty of Agriculture organized the 1st International Meeting Agriscience & Practice (ASP 2018), giving an opportunity to the participants for presentation and discussion of original scientific and practical results in different fields of agriculture.

The 1st International Meeting Agriscience & Practice (ASP 2018), heled on 10-11 May at Faculty of Agriculture - Stip, was organized with intention to bring together all agricultural stakeholders for sharing their knowledge, experience and obstacles. One of the main aims was to link research and field work in agricultural sector in the country and broader, giving it an international dimension. All oral presentations as well as poster presentations at ASP 2018 were organized in several scientific sessions:

- Agricultural economics,
- Plant biotechnology,
- Plant production,
- Plant protection,
- Quality control and food safety,
- Soil science and hydrology,
- Viticulture, enology and fruit production.

The main goal of the Meeting was linking and promoting scientific achievements and practical knowledge, presented in different thematic areas, which were achieved in the Republic of Macedonia and wider in the region.

Journal of Agriculture and Plant Sciences Vol. 16, No. 1 contains the presented papers from the 1st International Meeting Agriscience & Practice (ASP 2018).

Editorial Board,

Editor in chief,

August, 2018

Prof. Liljana Koleva Gudeva, PhD

JOURNAL OF AGRICULTURE AND PLANT SCIENCES, JAPS, Vol 16, No. 1, 2018

Manuscript received: 28.05.2018 Accepted: 21.06.2018



In print: ISSN 2545-4447 On line: ISSN 2545-4455 UDC: 637.5'652/'658.043:641.522.2 Original scientific paper

CHANGES IN FATTY ACID COMPOSITION OF POULTRY MEAT AFTER HEAT TREATMENT

Mitre Stojanovski^{1*}, Anita Čakarova², Aco Kuzelov³, Elena Joshevska¹, Gordana Dimitrovska¹, Dzulijana Tomovska¹, Katerina Bojkovska¹

¹Faculty of Biotechnical sciences, St. "Kliment Ohridski", University Bitola, R. Macedonia ²Food Industry "Blagoj Gorev", Veles, R.Macedona ³Faculty of Agriculture, Goce Delcev University, Stip, R.Macedonia *Corresponding author: mitre.stojanovski@yahoo.com

Abstract

The purpose of the paper is to determine the changes in the fatty acid composition of fat in poultry meat after the heat treatment, after frying in different types and combinations of fats. Lard, sunflower oil, palm oil and a combinations of three parts of lard and one part of sunflower oil, three parts of lard and one part of palm oil and three parts of palm oil and one part of sunflower oil were used for the examination. After frying with lard in the meat, there is a greater change in the fatty acid composition, where the fatty acid C18:1n9cof 45.84%, which it had it before frying, increased to 49.05%, C18:O decreased from 10.18% to 7.45% and decreased C16:O from 29.55 % to 27.45%. After frying in sunflower oil and palm oil, there has a formation of new fatty acids in insignificant amounts. In combination of three parts of lard and one part of sunflower oil, minimal changes in the fatty acid C14:O from 0.88% to 23.81%, fatty acids C16:O decreased from 27.24% to 1.72%, C18:1n9c from 48.05 % to 15.16% and C18:2n6c from 12.44% to 0.20%. After the heat treatment of the meat in three parts of palm oil and one part of sunflower oil, there is an increase in the content of fatty acid from C18:2n6cfrom 8.42% to 11.62 %.

Key words: quality, frying, lard, palm oil, sunflower oil

INTRODUCTION

Thermal processing of meat is regular action in cuisine and in the meat industry. The objective of thermal processing of meat is to prepare the meat for consuming and to increase its sustainability. One of the most applied and fastest ways of thermal processing of meat for consuming in households is frying. Soft, boneless pieces of meat from the shoulder part, the thigh, the loin and the neck, and from which the visible fat tissue was removed, are usually used for frying. Sunflower oil and lard are used in the households to fry meat, and lately the palm oil became very popular. During frying with less or more oil, sensory characteristics which are very alike to those formed during grilling or roasting, are formed in the meat. During frying of the meat with less or more oil, attractive chestnut brown colour is being formed on its

surface, while golden yellow to chestnut colour is formed on the surface of the fat tissue. The chestnut brown colour of the meat surface is consequence of Maillard reaction, which occurs during thermal processing.

During thermal processing – frying of meat, the temperature on the meat surface reaches 180-150°C (Rahelič et al., 1980) and numerous physical and chemical reaction happen within the meat. As a result of the heat, the water evaporates from the meat, the fats are extruded and they leak into the frying pan.

Changes which occur in the oils in which the meat is fried are primarily result of heat denaturation of fats, as well as a result of reaction between fats in the meat and soluble proteins.

103

MATERIAL AND METHODS

Material samples

The poultry that were used for examination are purchased from trade network usually from freshly slaughtered broilers (meat from the chest). The meat was cut down to size and shape as for frying, 80-100 g.

Fats used for the heat treatment of the meat (frying) and for examining their chemical changes after the heat treatment are lard, sunflower oil, palm oil and mixture, lard and sunflower oil in a ratio of 3:1, palm oil and sunflower oil in a ratio of 3:1, lard and palm oil in a ratio of 3:1. Lard used for the heat treatment of meat and for the examination of chemical changes in fat after the heat treatment of the meat is obtained from the trade network. The sunflower oil used to test the chemical changes in fats after the heat treatment of the meat is commercial oil produced in the oil factory "Blagoj Gorev" from Veles, R. Macedonia.

Thermal processing of the meat

Thermal processing – frying of the meat is performed in laboratory. The meat samples are frying after the melting and reaching the boiling point. The meat is fried in a large amount of fat, that is, the level of fat in the plate in which the meat was fried was enough to cover them. The determination of fried meat was subjective, that is, the conclusion that the meat was sufficiently thermally treated was the formation of the attractive colour on its surface. After the heat treatment, the meat is taken out of the plate in which it is thermally processed and it is transferred to another plate in which a kitchen paper for absorption of fat is placed on the bottom. The changes in meat weight were monitored by measuring meat before and after thermal processing with electronic weighing scale accurate to 0.1 g. Prior to the heat treatment of the meat, one sample of the fat in the amount of 25 cm³ was taken or a total of 12 samples of fat were taken for testing the fatty

acid composition before the treatment. Fatsampling before heat treatment is done after fat fading. The fats are placed in glass bottles with a volume of 30 cm. After the heat treatment of the meat and the cooling of the fats, four more samples are taken to examine the changes that occur in the fatty acid composition of the fats.

Determination of fatty acids content

Examination of changes in fatty acid composition of the fats is conducted before and after thermal processing .The fats from the analyzed samples of the poultry meat were extracted by the Rose-Gottlieb, (AOAC, 1990) method (modified method of Secchiari and Coll. (2003). Butylated hydroxytoluene (BHT) (50 µg BHT/ml in methanol) was added in order to avoid oxidative degradation of the fatty acids. Methylation of the fatty acids was done by the AOAC Official Method 996.06 (2005).

The produced metal esters of the fatty acids (FAMES) were analyzed with gas chromatograph Hewlett Packard 5890 series II with flame ionization detector and capillary column HP 88 ($60m \times 0.250mm \times 0.20 \mu m$). Helium was the gas carrier, and the total duration of one cycle was 38.50 minutes. The FAMES produced from the analyzed samples of the fermented sausages were identified on the basis of the comparison of their retention time with the retention time of the standards of FAMES consisted in 37 component mixture (Supelco 37 component FAME mix). The content of each of the analyzed fatty acids is shown in percent (%).

Analyses of changes in fatty acid composition of the fats are examined in the laboratory of the Food Institute at the Faculty of Veterinary Medicine in Skopje.

Statistical analysis

All data were presented as mean value with their standard deviation indicated (mean \pm SD).

RESULTS AND DISCUSSION

Significant physical changes, which contribute to change of basic characteristics of meat and forming new sensory characteristics important to its quality, occur during thermal processing of meat. One of the basic changes which occur during thermal processing is weight loss – meat shrinkage. The weight loss of the meat which occurs after thermal processing could be seen best from data given in Table 1.

104

		Parameters				
Meat frying	Changes in meat weight	x	S	Cv	min	max
Lard	Before thermal processing (g)	84.80	8.060	9.504	70.00	94.00
	After thermal processing (g)	46.80	5.741	12.267	42.00	56.00
	Shrinkage (%)	44.00	8.367	19.015	34.00	55.00
Sunflower oil	Before thermal processing (g)	85.60	5.426	6.334	82.00	94.00
	After thermal processing (g)	55.20	5.741	10.400	48.00	64.00
	Shrinkage (%)	35.66	2.494	6.994	32.00	39.00
Palm oil	Before thermal processing (g)	92.40	13.529	14.642	76.00	106.00
	After thermal processing (g)	57.40	10.151	17.684	44.00	66.00
	Shrinkage (%)	38.20	2.315	6.061	35.00	42.00
Lard and sunflower oil	Before thermal processing (g)	83.60	10.072	12.047	70.00	100.00
	After thermal processing (g)	51.60	8.139	15.773	40.00	64.00
	Shrinkage (%)	38.60	3.262	8.450	36.00	43.00
Lard and palm oil	Before thermal processing (g)	104.00	11.662	11.134	92.00	120.00
	After thermal processing (g)	65.60	4.630	7.058	60.00	72.00
	Shrinkage (%)	36.80	3.059	8.314	33.00	42.00
Palm oil and sunflower oil	Before thermal processing (g)	86.80	6.013	6.928	76.00	94.00
	After thermal processing (g)	59.60	9.830	16.494	44.00	68.00
	Shrinkage (%)	31.80	9.064	28.504	23.00	49.00

Table 1. Weight Loss-meat shrinkage in poultry meat during thermal processing

As seen from data presented in Table 2, the average weight loss during thermal processing of poultry by frying with sunflower oil is 35.66%, by frying with lard is 44.00 % and by frying with palm oil is 38.20%. The frying in a mixture of lard and sunflower oil, lard and palm oil, palm oil and sunflower oil in 3:1 the average weight loss is 35.53%. Shrinkage in poultry meat during thermal processing fried in palm oil and sunflower oil is the lowest and it is 31.80%, and the highest in meat that is fried in combination of lard and sunflower 38.60%. The obtained results in our examinations are in relation to the results stated by other authors. The meat shrinkage during thermal processing depends on the reached temperature in the centre of the product. The weight loss of meat

will be as bigger as the temperature of oils in which the meat is thermally processed gets higher (Buchar and Frohlich, 1969; Rohman, A. and Che Man, Y. B. 2009a; Naz Shanina, et al., 2005; Gunston, F.D. 2004). Significant changes of fatty acid composition occur after thermal processing of meat, which is caused by the temperature during thermal processing, as well as the interaction of fats and soluble proteins and the interaction of other meat substances and oils used for thermal processing – frying of meat, (Rashood, et al., 1996; Marina, et al., 2010; Marikkar et al., 2011; Indrasti, et al., 2010).

The changes in the fatty acid content that occur during thermal processing of meat – frying with lard, sunflower oil and palm oil could be seen best from Table 2.

	Different types of fats					
	Lar	d	Sunflower oil		Palm oil	
Fatty acid	before	after	before	after frying	before	after
composition	frying	frying	frying		frying	frying
C14:0	1.22	1.43	-	-	0.65	0.75
C16:0	29.55	27.45	10.58	11.57	46.15	44.43
C16:1	2.36	2.79	-	-	-	0.36
C17:0	0.23	0.28	-	-	-	-
C17:1	0.15	0.23	-	-	-	-
C18:0	10.18	7.45	3.49	3.58	3.21	3.25
C18:1n9c	45.84	49.05	45.42	45.51	44.40	44.77
C18:2n6c	9.00	9.31	40.06	38.80	4.06	6.21
C18:3n6	0.39	0.45	-	0.20	-	0.23
C21:0	-	0.45	-	-	-	-
C18:3n3	0.45	0.86	-	0.18	-	0.21
C22:0	-	0.25	-	-	-	-
C20:5n3	_	-	-	0.15	_	_

Table 2. Changes in the fatty acid content before and after heat treatment of poultry meat - fried with different types of fats

It is acertained that two fatty acids are formed in poultry fried with lard, such as C21:0 in an amount of 0.45% and C22:0 in an amount of 0.25 %. During the thermal processing of the poultry, the content of C18:1n9c has increased from 45.84% to 49.05%, while C18:0 has decreased from 10.18% to 7.45%, C16:0 has also decreased from 29.55% to 27.45%. These changes are statistically significant at the level of (p<0.5).During frying of poultry with palm oil there is minimal decrease of the content of C16:0 from 46.15 % to 44.43 %, and at the expense of it new fatty acids appear, such as C18:3n3 andC18:3n6, which are either result of denaturation changes which occurred in the palm oil during thermal processing of the meat,

or extracted from poultry. The changes which occur in fatty acid composition of fats after thermal processing of the meat are minimal and do not show statistical significance.

Most probably, the new fatty acids which appear in fats after thermal processing (most noticeable after frying with sunflower oil) of the meat are either extracted from the meat or newly created fat acids from the existing ones, under influence of high temperature during thermal processing of the meat

The changes in the fatty acid content that occur during thermal processing of meat – frying with combinations of fats could be seen best from Table 3.

	Different mixture of fats					
	Lard : sunflower oil Lard : palm oil		Palm oil : sunflower oil			
Fatty acid	(3:1)	3:1) (3:		(3:1)	(3:1)	
composition	before	after	before	after frying	before	after
	frying	frying	frying		frying	frying
C14:0	0.88	23.81	0.88	23.81	0.59	0.67
C16:0	27.24	1.72	27.24	1.72	42.31	41.99
C16:1	2.23	-	2.23	-	-	0.20
C17:0	0.12	-	0.12	-	-	-
C17:1	-	7.65	0.21	7.65	-	-
C18:0	8.15	51.25	8.15	51.25	3.23	3.48

Table 3. Changes in the fatty acid content before and after heat treatment of poultry meat - fried with different mixture of fats

C18:1n9c	48.05	15.16	48.05	15.16	45.24	41.71
C18:2n6c	12.44	0.20	12.44	0.20	8.42	11.62
C18:3n6	0.36	0.21	0.36	0.21	0.21	0.23
C18:3n3	0.53	-	-	-	-	0.10

Under the influence of high temperature during the frying of poultry in a mixture of lard and sunflower oil, one more significant change is that the content of C14: O increases from 0.88% to 23.81%, and decreases the content of C16:O from 27.24% to 1.72%, C18:O is increased from 8.16 to 51.25%, which is a consequence of the transformation of cis oleic fatty acid. After the heat treatment of poultry meat in the mixture of lard and sunflower oil, a new fatty acid C17:1 occurs in an amount of 7.65%, which we believe has been squeezed out of the meat. During the frying of poultry in a mixture of lard and palm oil,C16: O in the amount of 27.24% after frying decreased to 1.72%, while in oleic fatty acid (C18: O) before frying was represented by 8.15%, and after frying, its presence in fat

was 51.25%. A similar situation exists with the fatty acid of C18: 2n6c where its quantity of 12.44% has decreased to 0.20%. After frying of the poultry in this combination of fats there is a new fatty acid C17: 1 which is represented by 7.65%. Other fatty acids have no significant changes, they are almost unchanged. In the third combination of fats pamitic fatty acid (C16: O), which in fats before frying was 42.31% after frying fell to 41.99%, then fatty acid C18: 1n9c which in fat before frying was present in quantity of 45.24% decreased to 41.71%, while fatty acid C18: 2n6c of 8.42% was increased to 11.62%. For other fatty acids in fat before and after frying of poultry meat there are no major changes.

CONCLUSIONS

Based on the investigations carried out on chemical changes that occur in fats after the heat treatment, the following conclusions can be made:

- The average shrinkage during heat treatment of poultry meat in sunflower oil is 35.66%, lard 44.00% and palm oil 38.20%;
- After frying of poultry in lard, there is an occurrence of C18:3n3 with a representation of 0.45% and C22:0 of 0.25% and after frying in palm fat there is a new fatty acid C16: 1 where its presence is 0.36%.

During frying the meat in the mixture of fats, changes that occur in the fatty acid composition of fat in which the meat is processed present the fact that some fatty acids have decreased, others have increased and there have been new fatty acids. The emergence of new fatty acids is considered to be primarily a consequence of the extraction of fatty acids from the meat, and others are a result of the denaturation of fatty acids and the transition into a cis or trans form of the corresponding fatty acid.

REFERENCES

- Gunston, F.D. (2004). Creation and evaluation of a two-dimensional contour plot of fatty acid methyl esters after off-line coupling of reversed-phase HPLC and GC/EI-MS. *Analytical and Bioanalytical Chemistry*, (396), 2695–2707.
- Indrasti, D., CheMan, Y.B., Mustafa, S. & Hashim, D.M. (2010). Lard detection based on fatty acids profile using comprehensive gas chromatography hyphenated with time-of-flight mass spectrometry. *Food Chemistry* (122), 1273 - 1277.

Marikkar, J. M. N., Ng, S. L. & Che Man, Y. B. (2011). Composition and thermal analysis of lipids from prefried chicken nuggets. *Journal of the American Oil Chemists' Society*, (88), 749–754.

Marina, A.M., Che Man, Y.B. & Ismail, A. (2010). Use of the SAW sensor electronic nose for detecting the adulteration of virgin coconut oil with RBD palm kernel olein. Journal of the American Oil Chemists' Society, (87), 263–270.

Rashood, K. A., Shaaban, R. R. A., Moety, E. M.

A. & Rauf, A. (1996). Triacylglycerolsprofiling by high performance liquid chromatography: a tool for detection of pork fat in processed foods. *Journal of Liquid Chromatography*, (18), 2661-2672.

- Rohman, A. & Che Man, Y. B. (2009a). Analysis of cod liver oil adulteration using Fourier Transform Infrared (FTIR) spectroscopy. *Journal of the American Oil Chemists' Society*, (86),1149-1153.
- Rahelic, S. & Puac, S. (1980–1981). Fibre types in Longissimus dorsi from wild and highly selected pig breeds. *Meat Sci*ence, (5), 439–450.
- Ramírez M.R., Estévez M., Morcuende D., Cava R. (2004). Effect of the type of frying

culinary fat on volatile compounds isolated in fried pork loin chops by using SPME-GC-MS. Journal of Agricultural Food Chemistry, 52(25),7637-43.

- NazShanina, R., Siddiqi Rina Sheikh, S., & AsadSaYeed (2005). Deterioration of olive, corn and soybean oils due to air, light, heat and deepfrying. *Food Research International*, (38), 127-134,
- Olsen, E., Vogt, G., Ekeberg, D., Sandbakk, M., Pettersen, J., & Ntlsson, A. (2005). Analysis of the early stages of lipid oxidation in freeze-stored pork back fat and mechanically poultry meat. *Journal Agricultural Food Chemistry*, (53:2), 338-348.

ПРОМЕНИ ВО МАСНОКИСЕЛИНСКИОТ СОСТАВ КАЈ ЖИВИНСКОТО МЕСО ПО ТЕРМИЧКАТА ОБРАБОТКА

Митре Стојановски¹, Анита Чакарова², Ацо Кузелов³, Елена Јошевска¹, Гордана Димитровска¹, Џулијана Томовска¹, Катерина Бојковска¹

¹ Факултет за биотехнички науки, Универзитет "Св. Климент Охридски" - Битола, Р. Македонија ² Прехранбена индустрија "Благој Ѓорев" - Велес, Р.Македонија ³ Земіоделски факултет, Универзитет "Гоие Делчев" - Штип, Р.Македониіа *Контакт автор: mitre.stojanovski@yahoo.com

Резиме

Целта на трудот е да се утврдат промените во маснокиселинскиот состав на мастите кај пилешко бело месо по термичката обработка и тоа по пржење во различни видови и комбинации на масти. За испитување се користеше свинска маст, сончогледово масло, палмина маст и комбинација на три дела свинска маст и еден дел на сончогледово масло, три дела на свинска маст и еден дел на палмина маст и три дела на палмина маст и еден дел на сончогледово масло. По пржењето со свинската маст кај месото има поголеми промени во маснокиселинскиот состав каде што масната киселина C18:1n9c од 45.84% која ја имаше пред пржење се зголеми на 49.05%, С18:О се намали од 10.18% на 7.45% и намалување на С16:О од 29.55% на 27.45%. По пржење во сончогледовото масло и палмина маст има формирање на нови масни киселини во незначително количество. Во комбинација од три дела свинска маст и еден дел на сончогледово масло констатирани се минимални промени во маснокиселинскиот состав кој изнесуваат под 1 %. При пржење со три дела свинска маст и еден дел на палмина маст масната киселина C14:О од 0.88% се зголеми на 23.81%, масните киселини C16:О се намалија од 27.24% на 1.72 %, C18:1n9c од 48.05% на 15.16% и C18:2n6c од 12.44% на 0.20%. По термичката обработка на месото во три дела на палмина маст и еден дел на сончогледово масло има зголемување на содржината на масната киселина С18:2п6с од 8.42% на 11.62%.

Клучни зборови: квалитет, пржење, свинска маст, палмина маст, сончогледово масло