

Vol.

61




2018

**UNIVERSITY OF MINING AND GEOLOGY "ST. IVAN RILSKI"
JUBILEE INTERNATIONAL SCIENTIFIC SESSION**



JOURNAL
OF
MINING AND GEOLOGICAL
SCIENCES


part I
GEOLOGY
AND
GEOPHYSICS



UNIVERSITY OF MINING AND GEOLOGY "ST. IVAN RILSKI"

JOURNAL
OF
MINING AND GEOLOGICAL SCIENCES

Volume 61
Part I: Geology and Geophysics



Sofia, 2018

Editor-in-chief

Prof. Dr. Pavel Pavlov

University of Mining and Geology "St. Ivan Rilski"
1, Prof. Boyan Kamenov Str., 1700 Sofia, Bulgaria
e-mail: srs@mgu.bg; <http://www.mgu.bg/nis>

EDITORIAL BOARD

Prof. Dr. Lyuben Totev

Deputy editor, UMG "St. Ivan Rilski"

Prof. Dr. Viara Pojidaeva

Deputy editor, UMG "St. Ivan Rilski"

Assoc. Prof. Dr. Stefka Pristavova

Deputy editor, UMG "St. Ivan Rilski"

Prof. Dr. Yordan Kortenski

UMG "St. Ivan Rilski"

Assoc. Prof. Dr. Elena Vlasseva

UMG "St. Ivan Rilski"

Assoc. Prof. Dr. Antoaneta Yaneva

UMG "St. Ivan Rilski"

Prof. Dr. Desislava Kostova

UMG "St. Ivan Rilski"

Part I: Geology and Geophysics

Prof. Dr. Yordan Kortenski

Chairperson of an editorial board, UMG "St. Ivan Rilski"

Prof. Dimitar Sinyovski, DSc.

UMG "St. Ivan Rilski"

Prof. Dr. Maya Stefanova

Institute of Organic Chemistry, Bulgarian Academy of Sciences

Assoc. Prof. Dr. Nikolai Stoyanov

UMG "St. Ivan Rilski"

Prof. Dr. Radi Radichev

UMG "St. Ivan Rilski"

Prof. Stoyan Groudev, DSc

UMG "St. Ivan Rilski"

Prof. Dr. Strashimir Srtashimirov

UMG "St. Ivan Rilski"

Prof. Dr. Ognyan Petrov

Institute of Mineralogy and Crystallography, Bulgarian Academy of Sciences

Technical secretary: **Kalina Marinova**

Printig: **Publishing House "St. Ivan Rilski"**

©All rights reserved. Reproduction in part or whole without permission is strictly prohibited.

ELEMENTS AND STRUCTURE OF THE NORMAL GEOMAGNETIC FIELD IN THE REPUBLIC OF MACEDONIA

Marjan Delipetrev¹, Blagica Doneva¹, Gorgi Dimov¹

¹University of Goce Delchev, Faculty of Natural and Technical Sciences, 2000 Shtip, Macedonia, blagica.doneva@ugd.edu.mk

ABSTRACT. This paper presents the research of the normal geomagnetic field of the Republic of Macedonia. The complex geological structure of the explored area of regional and local character, has an important impact on the geomagnetic field. The geomagnetic field is composed of two components: normal and anomalous field. The geomagnetic field represents an important segment in the regulation of the natural processes on the Earth, which makes its permanent observation and study of special importance. Conducted investigations of the geomagnetic field maintain the continuity of monitoring and studying of the field changes in the area of the Republic of Macedonia which started in 2000.

Keywords: geomagnetic field, total field, declination, inclination, normal field, geomagnetic models and maps

ЕЛЕМЕНТИ И СТРУКТУРА НА НОРМАЛНОТО ГЕОМАГНИТНО ПОЛЕ В РЕПУБЛИКА МАКЕДОНИЈА

Мариан Делипетрев¹, Благица Донева¹, Горги Димов¹

¹Универзитет в Гоце Делчев, Факултет по природни и технички науки, 2000 Штип, Македонија, blagica.doneva@ugd.edu.mk

РЕЗЮМЕ. Тази статија представя изследването на нормалното геомагнитно поле на Република Македонија. Комплексната геолошка структура на изследваната област от регионален и местен характер има важно влияние върху геомагнитното поле. Геомагнитното поле се състои от два компонента: нормално и аномално поле. Геомагнитното поле представлява важен сегмент в регулирането на естествените процеси на Земята, което прави неговото постоянно наблюдение и изследване особено важно. Проведените изследвания на геомагнитното поле поддържат непрекъснатостта на мониторинга и изучаването на промените в полето в областта на Република Македонија, започнали през 2000 г.

Ключови думи: геомагнитно поле, общо поле, деклинация, наклон, нормално поле, геомагнитни модели и карти

Introduction

The geomagnetic field reflects the Earth's structure, the processes within it, as well as the external influences. The field is presented as a sum of three components: (Delipetrov, 2003).

- Normal geomagnetic field which reflects the causes of magnetism in the core and the mantle. This component includes agents that are in the depths of the earth's crust, but they are spread throughout the territory of the Republic of Macedonia;
- Regional geomagnetic anomalous field affected by characteristic generators of the magnetic field in the separated neotectonic zones in the Republic of Macedonia;
- Local geomagnetic field which is in direct correlation with the geological structures in the upper part of the Earth's crust with expressed magnetic features (increased concentration of ferrous magnetic minerals).

The territory of the Republic of Macedonia, as part of the Alpine orogen in the Balkan Peninsula, is characterized by a very complex geological structure, where structural segments are separated - zones which are widespread in the Balkan Peninsula. In recent geological history, neotectonic phase, different geological processes have divided these areas in separated units. The different geological structure and processes in these regions have significant impact on the

structure of the geomagnetic field. There is no other such small area as the Republic of Macedonia with so complex geological structure and hence, with such structure of the local geomagnetic field in whole Europe.

Models and maps of the elements of the normal geomagnetic field in the Republic of Macedonia

Two methods of modeling of the geomagnetic field prevail today, one is the method of spherical harmonic analysis and the other is the method of polynomial analysis.

The normal field for a territory can be expressed by:

$$E(\Delta\varphi, \Delta\lambda) = a_1 + a_2 \cdot \Delta\varphi + a_3 \cdot \Delta\lambda + a_4 \cdot \Delta\varphi^2 + a_5 \cdot \Delta\lambda^2 + a_6 \Delta\varphi \cdot \Delta\lambda$$

where:

$E(\Delta\varphi, \Delta\lambda)$ – is the value for the normal field of the point with coordinates φ_1 and λ_1 ;

φ_1 and λ_1 – is the geographic latitude and longitude of the place;

φ_0 and λ_0 – is the geographic latitude and longitude of the point depending on which measurements are being reduced; for the Republic of Macedonia the central point is $\varphi_0 = 41,50^\circ$ and $\lambda_0 = 22^\circ$;

$\Delta\varphi = \varphi_1 - \varphi_0$ – is the difference of geographic latitudes in minutes;

$\Delta\lambda = \lambda_1 - \lambda_0$ – is the difference of geographic longitudes in minutes;

a_i – are the coefficients for corresponding differences in nT/min ($\gamma/\text{min.}$), i.e. minutes / minutes or gammas and minutes. Usually, the differences of latitude and longitude are calculated in terms of coordinates of the geomagnetic observatory located on that territory.

Observations of the geomagnetic field of a given territory (usually state) are carried out in two parallel procedures. One is permanent monitoring of the geomagnetic field in an observatory. Another procedure that is performed periodically every 3 to 5 years are terrestrial observations of the basic network of repeat stations, and observations of the secondary net within a period of 10 years.

Based on this data collection about the geomagnetic field every five years, maps and models of the elements of the normal geomagnetic field for a given epoch are made.

Model of the elements of the normal geomagnetic field in the Republic of Macedonia

Field measurements were conducted on the fifteen repeat stations in Macedonia. (Delipetrev, 2011)]

During the field measurements, the following instruments were used:

- DI Fluxgate magnetometer – an instrument that measures values of the geomagnetic declination D and inclination I. The instrument consists of a non - magnetic theodolite and fluxgate sensor mounted on the telescope, so the optical and magnetic axes are parallel (fig. 1).
- Proton - magnetometer - GEOMETRICS G-856.

A model of the elements of the normal geomagnetic field in the Republic of Macedonia (Table 3) is made on the basis of the measured data (Table 1).




Fig. 1. DI Flux magnetometer - theodolite with Fluxgate sensor on the top

Table 1. Measured data on the repeat stations

| Station | H (nT) | X (nT) | Y (nT) | Z (nT) |
|--------------------|----------|----------|---------|----------|
| Tetovo | 24233.26 | ? | ? | 39941.46 |
| Egri | 24751.17 | 24716.87 | 1302.71 | 39237.34 |
| Mavrovo | 24264.10 | 24231.20 | 1263.12 | 39704.21 |
| Plackovica | 24288.75 | 24251.75 | 1340.17 | 39822.61 |
| Slivnica | 24379.72 | 24337.21 | 1439.08 | 39790.33 |
| Vodno | 24207.50 | 24169.74 | 1351.72 | 39950.88 |
| Bajlovce | 23883.98 | 23853.08 | 1214.61 | 40156.53 |
| Island Gradot | 24530.80 | 24484.68 | 1503.55 | 39402.71 |
| Nikolic | 24540.18 | 24504.80 | 1317.27 | 39576.18 |
| Ponikva | 24110.89 | 24073.10 | 1349.27 | 40111.39 |
| St. Maria Precesna | 24350.92 | 24315.68 | 1309.65 | 39651.61 |
| Crna Skala | 24227.53 | 24189.96 | 1348.62 | 40140.25 |
| Luke | 23937.17 | 23898.48 | 1360.40 | 40464.26 |
| Galicica | 24727.66 | 24696.04 | 1250.18 | 39098.67 |
| Prilep lake | 24520.03 | 24485.53 | 1300.37 | 39665.69 |

Table 2. Coefficients of the model of the normal geomagnetic field

| Element | Coefficients | | | | | |
|---------|--------------|---------|----------|---------|---------|----------|
| | a_0 | a_1 | a_2 | a_3 | a_4 | a_5 |
| D | 3.6278 | 0.118 | -0.0386 | 0.078 | 0.0154 | -0.3455 |
| I | 58.088 | -0.037 | -0.1578 | 0.119 | 0.0928 | -0.4256 |
| T | 46550 | -24.449 | -172.278 | -20.585 | 197.784 | -449.635 |
| H | 24607 | 11.072 | 17.149 | -95.376 | 38.900 | 57.755 |
| X | 24558 | 7.939 | 18.218 | -97.322 | 37.095 | 66.862 |
| Y | 15560 | 46.635 | -15.966 | 23.628 | 13.308 | -145.088 |
| Z | 39514 | -36.462 | -217.088 | 33.564 | 210.554 | -567.011 |

Table 3. Values of the normal geomagnetic field calculated with the coefficient from Table 2

| Stations | D | I | T | H | X | Y | Z |
|--------------------|-------|--------|-------|-------|-------|------|-------|
| Galicica | 3.378 | 57.276 | 46271 | 25014 | 24970 | 1474 | 38927 |
| Egri | 3.465 | 57.367 | 46330 | 24984 | 24938 | 1510 | 39016 |
| St. Maria Precesna | 3.533 | 58.039 | 46453 | 24590 | 24543 | 1515 | 39411 |
| Mavrovo | 3.466 | 58.122 | 46451 | 24531 | 24486 | 1483 | 39445 |
| Prilep lake | 3.509 | 57.854 | 46567 | 24777 | 24731 | 1517 | 39428 |
| Vodno | 3.597 | 58.298 | 46671 | 24526 | 24477 | 1539 | 39707 |
| Bajlovce | 3.507 | 58.850 | 46675 | 24144 | 24099 | 1477 | 39945 |
| Ponikva | 3.721 | 58.603 | 46750 | 24355 | 24304 | 1581 | 39905 |
| Plackovica | 3.698 | 58.217 | 46575 | 24531 | 24480 | 1582 | 39591 |
| Luke | 3.737 | 59.005 | 46235 | 23809 | 23759 | 1552 | 39633 |
| Tetovo | 3.617 | 58.346 | 46683 | 24499 | 24450 | 1546 | 39738 |
| Island Gradot | 3.946 | 57.667 | 46327 | 24777 | 24719 | 1685 | 39144 |
| Nikolic | 3.499 | 57.823 | 46493 | 24759 | 24713 | 1511 | 39352 |
| Slivnica | 3.983 | 58.166 | 46607 | 24583 | 24524 | 1708 | 39596 |
| Crna Skala | 3.769 | 58.244 | 46903 | 24685 | 24632 | 1623 | 39881 |

Maps of the elements of the normal geomagnetic field of the Republic of Macedonia

The declination of the observed territory varies from 3,378° on the measuring point Galicica to 3,983° on the station Slivnica. Declination varies in the interval $\Delta D = 0,605^\circ$ (Fig. 2).




Fig. 2. Map of declination, D

Analysis of the declination field shows that on the observed territory the extreme stands in the central southern part, near the measuring station Island Gradot. The field is quiet in the west and northwest, which should be expected according to the geological structure.

From the observations, the inclination varies from Galicica $57.276^\circ \leq I \leq 59.005^\circ$ Luke. The mean value of inclination is $I_{sr} = 58,125^\circ$. The field of inclination I related to that of the declination is more homogenous and relatively quiet. There is a slight twisting of isolines in the central southern part (Fig. 3).




Fig. 3. Map of the inclination, I




Fig. 4. Map of total intensity, T

Values of the measured points for the total field vector T varies in the interval from 46 235 nT on the repeat station Luke to 46 903 nT on the point Crna Skala. The interval of variation is $\Delta T = 371,0$ nT. The field is relatively quiet with a twisting of the isolines in the central southern part and in the northeastern part (Fig. 4).




Fig. 5. Map of horizontal component, H

Analyzing the map of the horizontal component H (Fig. 5) from north in the area of Bajlovcce with a minimum value of 24150 nT toward south shows an increase to $H_{max} = 25000$ nT in the area of Ohrid and Prespa Lake, i.e. in the region of the repeat station Galicica. The average value of the field is $H_{sr} = 24575$ nT. The interval, in which the horizontal component on the observed territory varies, is $\Delta H = 850$ nT. The field is homogenous and relatively quite.

The field of the northern component X (Fig. 6) is relatively quiet and with a maximum value $X_{max} = 24900$ nT from southwest, decreasing toward north and in the northeastern part it has a minimum value $X_{min} = 24100$ nT. The average value of the northern component is $X_{sr} = 24500$ nT. The interval of variation is $\Delta X = 800$ nT.




Fig. 6. Map of the northern component, X

The field of the eastern component Y (Fig. 7) varies from $Y_{min} = 1470$ nT in the area of the repeat stations Galicica and Bajlovcce to $Y_{max} = 1700$ nT at the repeat stations Slivnica and

Island Gradot. The western part of the Republic of Macedonia has a quieter field related to the central and the eastern part. The expressed extreme is present in the southern part. The value of the component varies in the interval of $\Delta Y = 250$ nT. The average value is $Y_{sr} = 1595$ nT.




Fig. 7. Map of the eastern component Y

The field of the vertical component Z (Fig. 8) is relatively quiet with expressed twisting of the isolines in the northeastern and central southern part. The maximum value is in the area of the repeat station Bajlovce $Z_{max} = 39945$ nT, and minimum value is in the region of the repeat station Galicica $Z_{min} = 38927$ nT. The average value of the component is $Z_{sr} = 39435$ nT. The interval of variation is $\Delta Z = 1018$ nT.




Fig. 8. Map of the vertical component, Z

The fields of declination - D and eastern Y component are more complex in terms of other components of the field. In all maps, the western part of the territory of the Republic of Macedonia has a quieter field compared to the central and eastern part.

The comparison of maps of the relevant elements of the normal field from previous measurements in 1990 showed lower degree of similarity with these maps because of the use of a different network of repeat stations, the greater distance and difference in applied technology.

Structure of the geomagnetic field

The geomagnetic field on the Earth's surface is a sum of vectors of many different magnetic fields:

$$T = T_0 + T_M + T_a + T_e + \delta T.$$

The implemented geomagnetic field observation and the applied methodology of data processing have eliminated the influence of the local anomaly field T_{al} , the electromagnetic field T_e , and the impact of short time variations of the field δT .

In order to eliminate the influence of geomagnetic causes that are deep underground, and because their dimensions and area of influence are far beyond the territory of the investigated area, an IGRF (International Geomagnetic Reference Field) model is used as a filter that reflects the impact of normal geomagnetic field, in this case shown by two components: the field- T_0 and the axial dipole field- T_k on the continent.

Maps of the differences of the normal geomagnetic field's elements between the IGRF model and our model were created (Figs. 9 - 12). (Panovska et al., 2006)

The analysis of the components of the regional anomaly of the geomagnetic field T_{ar} reflects the neotectonic regionalization of the Republic of Macedonia in three mega blocks:

- Eastern Macedonian zone;
- Vardar zone;
- Western Macedonian zone including Pelagonian block.




Fig. 9. Map of ΔT between our measurements and IGRF model




Fig. 10. Map of ΔZ between our measurements and IGRF model




Fig. 11. Map of ΔD between our measurements and IGRF model

Conclusion

From the data presented on the elements and structure of the normal geomagnetic field in the Republic of Macedonia, it can be concluded the following:

- The maps and the elements' models of the normal geomagnetic field express the specificity of the area of the Republic of Macedonia;
- The presented data for the declination can have wide application in all activities where compass is used;
- Maps of the differences of the normal geomagnetic field elements between the IGRF model and our model were created.




Fig. 12. Map of ΔI between our measurements and IGRF model

References

- Delipetrov, T. Basics of geophysics. Faculty of mining and geology, Shtip, 2003. - pp. 49.
- Delipetrov, M. Structure of the geomagnetic field on the territory of the Republic of Macedonia, Doctoral thesis, Faculty of Natural Science and Mathematics, Skopje, 2011.
- Panovska, S., Delipetrov T., Delipetrov B. Correlation Between IGRF2000 Model and Measured Geomagnetic Data on the Territory of the Republic of Macedonia from 2003 and 2004 Measurements, XIIth IAGA Workshop on geomagnetic observatory instruments, data acquisition and processing, Belsk, Poland, June, 2006.