UDK: 631

SRPSKO DRUŠTVO ZA PROUČAVANJE ZEMLJIŠTA

ZEMLJIŠTE I BILJKA



ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 2013.

BEOGRAD, 2013.

ZEMLJIŠTE I BILJKA SOIL and PLANT – ПОЧВА И РАСТЕНИЕ

EDITOR-IN-CHIEF Dr Gligorije M. ANTONOVIĆ

PUBLICATION STAFF: Marija ŽIVANOVIĆ, Publication Editor

ADVISORY BOARD

Gligorije ANTONOVIĆ (Beograd), Srboljub MAKSIMOVIĆ (Beograd), Duro BOŠNJAK (Novi Sad), Vinsent van ENGELEN (Wageningen), Frederik HIL WING (Heuseden), Svjetlana Cupać (Beograd), Imre MOLNAR (Novi Sad), Josif MITRIKESKI (Skopje), Tatjana MITKOVA (Skopje), Olga NAJDANOVSKA (Skopje), Boris NIKOLOV (Sofija), Husnija RESULOVIĆ (Sarajevo), John STRATUS (Thessaloniki), Gradimir VASIĆ (Beograd), Anastasios VOULGAROPOULOS (Thessaloniki)

IZDAJE: Srpsko društvo za proučavanje zemljišta, Beograd - Zemun, Nemanjina 6, fah 44 Published by Serbian soil science society U troškovima publikacije učestvuje Republika Srbija MINISTARSTVO NAUKE, Beograd.

Štampa: Akademska izdanja, Zemun

CIP – Каталогизација у публикацији Народна библиотека Србије, Београд 631 ZEMLJIŠTE i biljka =Soil and plant = Почва и растение / editor-in-chief Gligorije M. Antonović. – God. 1, br. 1 (maj 1952) - Beograd - Zemun (Nemanjina 6) : Srpsko društvo za proučavanje zemljišta, 1952 – (Zemun : Akademska izdanja). – 24 cm. –

Tri puta godišnje ISSN 0514-6658 = Zemljište i biljka COBISS.SR-ID 338701

Official journal of the Serbian soil science society Editorial office: 11081 Zemun, Nemanjina 6, PO Box 44 UDK: 631

YU ISSN 0514-6658

SRPSKO DRUŠTVO ZA PROUČAVANJE ZEMLJIŠTA



ZEMLJIŠTE I BILJKA



ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 2013.

BEOGRAD, 2013.

ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 2013.

SOIL AND PLANT, Vol. 62, No. 1, 2013.

CONTENTS

Original scientific papers

SADRŽAJ

Originalni naučni radovi

- Dražen D.Vicić, Milovan M. Stoiljković, Branislav Jurišić, Svetlana Polavder, Nenad Č. Bojat: Odnos Mg:Ca u korenu, stablu i listovima vrste *Teucrium montanum* L. sa serpetinskih staništa Srbije 31

UDC 631.847 Original scientific paper

DETERMINATION OF BROCCOLI LEAF AREA AND STOMATAL NUMBER USING DIFFERENT APPLICATION METHODS OF MICROBIOLOGICAL FERTILIZER - SLAVOL

Olga NAJDENOVSKA¹, Snezana DJORDJEVIC², Elizabeta MISKOSKA-MILEVSKA¹, Daniela DIMOVSKA³ and Zoran POPOVSKI¹

¹Faculty of Agricultural Sciences and Food – Skopje ²Faculty of Agriculture Belgrade-Zemun, Serbia ³Centre for Applied Research and Permanent Education in Agriculture – Skopje

Najdenovska Olga, Snezana Djordjevic, Elizabeta Miskoska-Milevska, Daniela Dimovska and Zoran Popovski (2013): Determination of broccoli leaf area and stomatal number using different application methods of microbiological fertilizer - Slavol. – Zemljište i biljka, Vol.62, No.1, 1-13, Beograd.

Application of microbiological fertilizers in agriculture has positive effect on plant production and also solves problems regarding the environmental protection. The fertilizer research is currently focusing on the effects of microbiological fertilizers on living organisms and soil. The aim of the study was to investigate the effect of different methods of application of microbiological fertilizer – Slavol on leaf area and stomatal number in broccoli "Verdia F_1 " (*Brassica oleracea* L. var. *italica* Plenck.) plants. Two different methods of fertilizer application were used in this study: foliar

Corresponding author: Elizabeta Miskoska-Milevska, University Ss. Cyril and Methodius, Skopje, Faculty of Agricultural Sciences and Food, Bldv. Edvard Kardelj b.b. P.O. Box 297, Skopje R. of Macedonia; e-mail: miskoska@yahoo.com

For many years scientists have investigated the positive effects of microorganisms' activities on plant production (STOLILJKOVIĆ, 1988; GOVEDARICA et al., 1999; NAJDENOVSKA et al., 2002a; NAJDENOVSKA et al., 2002b; NAJDENOVSKA et al., 2002c; NAJDENOVSKA et al., 2004; NAJDENOVSKA, 2011; NAJDENOVSKA, 2012; ĐORĐEVIĆ et al., 2005a; ĐORĐEVIĆ et al., 2005b; BALEŠEVIĆ-TUBIĆ et al., 2011). A number of plant growth-promoting rhizobacteria (PGPR) belonging to various taxonomic groups contain the enzyme 1-aminocyclopropane-1-carboxylate (ACC) deaminase, which hydrolyses ACC, the immediate precursor of the plant hormone ethylene (GLICK et al., 1995 and BELIMOV et al., 2005). Lowers the levels of ethylene produced in developing or stressed plants promoting root elongation. A large array of heavy metaltolerant bacteria including species of Pseudomonas, Azotobacter, Klebsiella, Enterobacter, Alcaligenes, Arthrobacter Burkholderia, Bacillus and Serratia have reported to enhance plant growth (KLOPPER et al., 1989 and GLICK, 1995). Several bacteria improve plant growth through suppression of pathogens by competing for nutrients, by antibiosis, or by synthesizing siderophores, which can solubilize and chelate iron from the soil and inhibit the growth of phytopathogenic microorganisms (CABALLERO-MELLADO et al., 2007).

Also, the productivity and health of horticultural crops depends on the ability of the plant cover to intercept light energy. This ability is a function of the amount of leaf area, the architecture of the vegetation cover, and plants' ability to convert light energy. Leaves are complex plant organs upon which life depends. Their primary function is process of photosynthesis. Through the process of photosynthesis they are able to trap energy in the form of sugar molecules. In order to maximize the photosynthetic production, plants increase the photosynthesizing surface. Estimation of leaf area is valuable in studies of plant nutrition, plant competition, plant-soil-water relations, plant protection measures, respiration, light reflectance and heat transfer in plants (MOHSENIN, 1986), Also, leaf area is an important parameter in understanding photosynthesis, light interception, water and nutrient use and crop growth and yield potential (SMART, 1974; WILLIAMS, 1987). Leaf area can be measured by nondestructive or destructive measurements. Agronomists require rapid, inexpensive, reliable and nondestructive method for measuring leaf area. Nondestructive estimation of plant leaf area offers inexpensive and reliable alternatives in horticultural experiments.

Stomata control the movement of gases in and out of a leaf, making carbon dioxide available for photosynthesis and controlling the loss of water from the leaf through transpiration. The number of stomata is affected by ecclogical condition and physiological process (DŰZENLI and ERGENOĞLU, 1991). Individual plants can change the stomatal number to adjust for changes in the environment. According to RYUGO (1988) stomatal frequency differs greatly from

ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 1-13, 2013.

application and application through irrigation water (drip irrigation system). The experiment was carried out on alluvial soil in the private farm located in Jurumleri, near to Skopje, Macedonia. Obtained results indicated that maximum average leaf area was 361.39 cm^2 in plants with foliar application of Slavol and minimum average leaf area was 167.86 cm^2 in control plants. The leaf area of plants treated with Slavol through drip irrigation system was 262.68 cm^2 . It can be concluded that method of foliar application of Slavol obtained the highest values of leaf area. According to the LSD-test these values were significantly different (p<0.01).

Key words: broccoli, leaf area, microbiological fertilizer, stomatal number

INTRODUCTION

Broccoli (Brassica oleracea L. var. italica Plenck.) belongs to the genus Brassica and it is one of the major crop plants in this genus. It is an Italian vegetable, native to the Mediterranean region, cultivated in Italy in ancient roman times. In USA, the broccoli first appeared in 1806, but it commercially cultivation was started around 1923 (DECOTEAU, 2000). The broccoli is an important vegetable crop and has high nutritional and good commercial value (YOLDAS et al., 2008).

Fertilizers are compounds usually given to plants with the intention to increase the yield and quality of crops. Over the last few decades there has been an increase in the use of organic fertilizers due to the rise in popularity of organic food production. The development of microbiological fertilizer (biofertilizer) could be connected with green organic food safety and environment protection which display a vital role in the sustainable development of high-yield, highquality, high-efficiency agriculture (MENG et al., 2008). The use of chemical fertilizers in agriculture led to deterioration of soil conditions and environment. Biofertilizers are alternative fertilizer and one of the important components of organic farming that enhances the plant growth and yield. Namely, biofertilizer is a substance which contains living microorganisms which, when applied to seed, plant surfaces, or soil, colonizes the rhizosphere or the interior of the plant and promotes growth by increasing the supply or availability of primary nutrients to the host plant (VESSEY, 2003). Application of biofertilizers can be expected to reduce the use of chemical fertilizers and pesticides. From the economic and ecological aspects microbiological fertilizers are much more acceptable than the use of mineral fertilizers and chemicals (MILIC et al., 2003). Microbiological fertilizers may contain one type of microorganism or a mixture of different microorganisms. Natural bio-organic fertilizer - Slavol is made up of the combined nitrogen fixing and phosphorus mineralizing bacteria (Azotobacter chroococcum, A. vinelandi, Derxia sp., Bacillus megaterium, B. lichenformis and B. subtilis).

Olga NAJDENOVSKA et al .: BROCCOLI LEAF AND APPLICATION OF SLAVOL

ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 1-13, 2013.

species to species, ranging from 125 to over 1000 stomata/mm². In some resources, it has been known that stomatal density changes with characteristics such as drought, net photosynthesis production (BIERHUIZEN *et al.*, 1984), precipitation change (MISIRLI and AKSOY, 1994), vegetative developmental phases and grafts on different rootstocks (ÇAGLAR and TEKIN, 1999) and altitude (ÇAĞLAR *et al.*, 2004; ASLANTAŞ and KARAKURT, 2009).

No information for effect of microbiological fertilizer – Slavol on broccoli leaf area and stomatal number can be found in the literature. The aim of the present experiment was to study the effect of different methods of application of microbiological fertilizer – Slavol on broccoli leaf area and stomatal number.

MATERIALS AND METHOD OF WORK

The study was carried out in the private farm located in Jurumleri near to Skopje, Macedonia. Broccoli "Verdia F_1 " (*Brassica oleracea* L. var. *italica* Plenck.) plants were used as plant material. The influence of different methods of application of microbiological fertilizer – Slavol on broccoli leaf area and stomatal number was investigated. Two different application methods of fertilizer were used in this study: foliar application and application through irrigation water (drip irrigation system).

The seedlings were grown in a greenhouse. Firstly, the seedlings (about 45days old) were treated with Slavol (10 mL/L) for few minutes, than they were transplanted in rows. The broccoli plants were planted on alluvial soil and they were treated with Slavol using different application methods during the vegetation period, starting from 21th of August until 1th of October 2012. One group of broccoli plants was treated with Slavol through the leaf (foliary) in the dose of 10 mL/L in intervals of 7 days. The second group of broccoli plants was treated with Slavol through drip irrigation system (1mL/L) in intervals of 2 days. The flow capacity of drip irrigation system was 2L /min/m².

Leaf area was determinate by using 30 leaf samples taken from broccoli plants one month after last application of Slavol. In same time, leaf samples from not treated (control) plants were analyzed as a control for comparison. The leaves used in this research were randomly selected from each group of broccoli plants. After cutting, the leaves were placed in plastic bags and were transported on ice to the laboratory. The leaf width (cm) was measured from tip to tip at the widest part of the lamina and leaf length (cm) was measured from lamina tip to the point of petiole intersection along the lamina midrib.

Broccoli leaf area was determined using model of nondestructive estimation of broccoli leaf area by STOPPANI *et al.* (2003). According to these authors the broccoli leaf area estimation equation was found to be, LA $(cm^2) = 0.63$ (leaf length x leaf width at the widest point).

Stomata on upper and lower leaf surface were counted on the middle portion of the leaf by nail polish replicas (BREWER, 1992). The prepared slides were analyzed using light microscope by a 16x ocular and a 40x objective. The number of stomata was expressed per mm² of leaf area. Average stomatal number was calculated from 10 leaves per treatment using counts from three areas per leaf. Collected data were statistically processed using analysis of variance (ANOVA) and the significance was evaluated by least significant differences (LSD) at p < 0.05 and p < 0.01.

RESULTS AND DISCUSSION

In the present study, Slavol as liquid fertilizer was applied using two different methods: foliar application and application through irrigation water. Foliar application refers to the spraying of fertilizer solution on the foliage of growing plants. Application through irrigation water (fertigation) refers to the application of water soluble fertilizers through irrigation water. In the research was used drip irrigation system. This method saves water and fertilizer by allowing water to drip slowly to the roots of plants. The drip irrigation has some advantages and disadvantages. Namely, fertilizer and nutrient loss is minimized, water application efficiency is high, weed growth is minimized and foliage remains dry, reducing the risk of disease. On the other hand, this application method of fertilizers is expensive.

According to SARIC *et al.* (1989) feeding plant affects numerous physiological-biochemical processes that are important for plant growth, development and yield. Application of fertilizers in vegetable crops is important for obtaining higher yields and products which are characterized by better quality (MOLNAR, 1995). Leaf is one of the important parts of the plants because it is the one which is in charge of accomplishing the photosynthesis, the respiration and transpiration. According to VOLKENBURGH (1999) the leaf expansion depends of genetic constitution and environmental conditions.

Table 1 Average	broccoli leaf	area (cm²)	after usin	g different application
	me	thods of Sla	avol	

	Control plants	Plants with foliar application of Slavol	Plants treated with Slavol through drip irrigation system		
Average leaf area (cm ²)	167.86	361.39	262.68		
Minimum	60.48	144.9	90.72		
Maximum	264.6	595	468.72		

In this study, broccoli leaf area was estimated with nondestructive model by STOPPANI et al. (2003). Using new equipment and tools such as hand scanner or

Olga NAJDENOVSKA et al .: BROCCOLI LEAF AND APPLICATION OF SLAVOL

laser optical apparatuses for determination of leaf area is very expensive investments for basic and simple research. Development of mathematical models and equations from linear leaf measurements has been shown to be very useful in studying plant growth and development (ELSNER et al., 1988; RAJENDRAN and THAMBURAJ, 1987). Non-destructive leaf area estimation also saves time as compared with geometric reconstruction. Nondestructive leaf area or plant growth measurements are often desirable because continued use of the same plants over time can reduce variability in experiments as compared with destructive sampling (NESMITH, 1991; NESMITH, 1992; GAMIELY et al., 1991). Because leaf length or width are dimension that can be easily measured in the field, green-house and pod experiments, use of this model would enable researchers to make nondestructive measurements or repeated measurements on the same leaves (FALLOVO et al., 2008). According to same authors, the leaf area (length: width ration) may vary among different plant materials and it is necessary to be found good model of nondestructive leaf area that will be used in physiological study independently of the genetic materials.

Average leaf area in plants treated with Slavol was larger than in control plants (Table 1). According to the data in Table 2, the LSD-test showed that the differences among the average leaf areas in control and treated plants were statistically significant (p<0.01). The smallest value of average leaf area (167.86 cm²) was detected in control plants and the biggest value (361.39 cm²) was detected in plants with foliar application of Slavol.

Table 2.- Analysis of variance for Table 1.

Source of Variation	SS	df	MS	F	P-value	F crit
Rows	378853.2	29	13063.9	1.229365	0.247974	1.662901
Columns	561854.2	2	280927.1	26.43634	6.91E-09	3.155932
Error	616339.9	58	10626.55			
Total	1557047	89				

LSD_{0.01} 70.80706

......

This conclusion was in agreement with FRITZ (1978) who pointed out that a repeated application of small units of foliar fertilizers stimulates plant metabolism and an increased nutrient uptake via the roots can be observed. According to YILDIRIM *et al.* (2007) one of the benefits of foliar fertilization is the increased uptake of nutrients from the soil. This notion is based on the belief that the foliar fertilization causes the plant to release more sugars and other exudates from its roots into the rhizosphere. Beneficial microbial populations in the root zone are

stimulated by the increased availability of these exudates. In turn, this enhanced biological activity increases the availability of nutrients, disease-suppressive biochemicals, vitamins and other factors beneficial to the plant (YILDIRIM *et al.*, 2007). Also, KOLOTA and OSINSKA (2001) recognized that supplementary foliar fertilization during crop growth can improve the mineral status of plants and increase the crop yield.

Also, the plants treated with Slavol through drip irrigation system had bigger average leaf area (262.68 cm²) than in control plants (Table 1). According LSDtest difference between average leaf area of plants with foliar application of Slavol and plants treated with Slavol through drip irrigation system is significant (p<0.01) (Table 2). From obtained data it can be concluded that highest average values of leaf area were obtained using method of foliar application of microbiological fertilizer Slavol.

KANDIL and GAD (2009) researched effects of inorganic and organic fertilizers on growth and production of broccoli and found that using organic manure plus inorganic solution fertilizers gave a significant effect on plant growth, heads yield, chemical constituents and mineral composition of broccoli. Also, their values for broccoli leaf area were from 169 to 294 cm². In our research, obtained value for average broccoli leaf area in plants with foliar application of Slavol (361.39 cm²) was bigger than values for broccoli leaf area presented by KANDIL and GAD (2009).

UZUN and KAR (2004) found significant difference between leaf area of broccoli plants and planting times. According to GRANIER and TARDIEU (1998) variation in leaf size has been attributed to differences in cell number, or cell size, or combinations of the two. It is known that cell division in plants is correlated with carbohydrate supply, and that leaves starved for photosyntate or other nutrients will develop fewer cells than those growing amid plenty (DALE, 1988; CHAPIN, 1991). Many researchers (MEDINA, 1970; GULMON and CHU, 1981) have reported reducing leaf area under limiting nitrogen. Microbiological fertilizers which contain highly efficient strains of bacteria, fungi and algae provide plants with biogenous elements: nitrogen, phosphorus and potassium (GOVEDARICA *et al.*, 2002). In previous studies was confirmed positive effect of microbiological fertilizer – Slavol on plant production (DORDEVIĆ *et al.*, 2005a; JELAČIĆ, 2007; MISIMOVIĆ *et al.*, 2012).

Our data indicated that the nitrogen fixing and phosphorus mineralizing bacteria present in the microbilogical fertilizer - Slavol probably increase the supply of plant nutrients that improve the nutritional status of plants and increased leaf size in treated plants. This is in agreement with study of NAJDENOVSKA *et al.* (1998). Namely, according to these authors nitrogen fixing bacteria synthesize substances of growing so that they improve the growing and development of plants, as well as the resistance of the plants to illnesses and harmful insects.

6

ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 1-13, 2013.

According to JONES et al. (1987) stomatal dimension and especially frequency can change more than two fold in response to radiation, to water status or according to developmental stage. The leaves of broccoli are amphistomatous i.e. stomata are on upper and lower epidermis. According to ASTON (1978) stomata on upper epidermis are more sensitive to environment variables.

Table 3. - Average stomatal number of broccoli leaves (stomata/mm²) after using different application methods of Slavol

Average stomatal	Control plants	Plants with foliar application of Slavol	Plants treated with Slavol through drip irrigation system	
number on upper epidermis (stomata/mm ²) Average stomatal	145	102	140	
number on lower epidermis (stomata/mm ²) Total stomatal	250	232	242	
number (stomata/mm ²)	395	334	382	

As presented in Table 3, stomatal number on lower epidermis is bigger than on upper epidermis. Average stomatal number on upper and lower epidermis of control plants was 145 and 250 stomata/mm², respectively. Also, in plants treated with Slavol through drip irrigation system stomatal number on upper epidermis were 140 stomata/mm² and 242 stomata/mm² on lower epidermis. On the other hand, average stomatal number on upper and lower epidermis of plants with foliar

application of Slavol was 102 and 232 stomata/mm², respectively (Table 3). No significant difference was detected between stomatal number on upper epidermis in control plants and plants treated with Slavol through drip irrigation system (Table 3). The plants with foliar application of Slavol had the lowest stomatal number on upper epidermis compared with control plants and plants treated with Slavol through drip irrigation system. According to the LSD-test these differences were statistically significant (p<0.01).

Also, no statistically significant difference was found among treatments and control refer to stomatal number on lower epidermis.

The difference between the total stomatal numbers in control plants and plants treated with Slavol through drip irrigation system was not statistically significant. In plants with foliar application of Slavol was measured the lowest total stomatal number (334 stomata/mm²). According to the SDS-test, the difference between the control plants and plants with foliar application of Slavol was statistically significant (p<0.01). Also, analyses of total stomatal number in plants treated with Slavol through drip irrigation system and plants with foliar application of Slavol showed statistically significant difference (p<0.05) (Table

Olga NAJDENOVSKA et al .: BROCCOLI LEAF AND APPLICATION OF SLAVOL

Stomatal characteristics are one of the main factors that determine photosynthetic ability and the genetic basis for stomatal frequency is still poorly understood (ARMINIAN et al., 2008). According to these authors phenotyping may not have been sufficiently precise, there was a strong environmental influence or a large genotype by environment interaction was involved in variability of these traits. Also, these authors suggest that stomatal characteristics are not an appropriate approach to breed high yielding wheat varieties. KÜRSCHNER (1997) demonstrated empirically that a difference exists in stomatal density and stomatal index response during leaf expansion depending upon the position of the leaf in tree crown.

Table 4.- Analyses of variance table for Table 3 (total stomatal number)

ANOVA						
Source of Variation	SS	df	MS	F	P-value	F crit
Rows	134667.1	29	4643.694	0.775841	0.769357	1.662901
Columns	62787.22	2	31393.61	5.245058	0.008058	3.155932
Error	347151.4	58	5985.37			
Total	544605.8	89			•	

LSD_{0.05} 39.95714

LSD001 53.14053

From obtained results, no general conclusion can be given refer to effects of application methods of fertilizer - Slavol on stomatal number in broccoli leaf.

CONCLUSION

By this experiment it is proved that microbiological fertilizer -Slavol had positive statistically significant effect on leaf area in both group of plants where was used different method of application of microbiological fertilizer - Slavol. Also, it can be concluded that highest average values of leaf area were obtained using method of foliar application of microbiological fertilizer Slavol. The LSDtest showed that differences among plants treated with microbiological fertilizer -Slavol and control plants were statistically significant (p<0.01).

In plants with foliar application of Slavol were detected the lowest total stomatal number compared with control plants and plants treated with Slavol through drip irrigation system. According to the LSD-test the difference between ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 1-13, 2013.

control and plants with foliar application of Slavol was statistically significant (p<0.01). Also, the difference between plants treated with Slavol through drip irrigation system and plants with foliar application of Slavol was statistically

The results presented here support the hypothesis that microbiological fertilizers can improve plant grow but, more studies are needed to be better understand effect of microbiological fertilizers on living organisms and soil.

REFERENCES

ASLANTAS R. And KARAKURT H. (2009): The Effects of altitude on stomatal number and some vegetative growth parameters of some apple cultivars. Res. J. Agric. Biol. Sci. 5. 853-857. ARMINIAN A., HAUSHMAND S., KNOX R.E. and SHIRAN B. (2008): Stomatal Characteristics.

Heritability and Their Relationship to Grain Yield in Double Haploid Bread Wheat Population, 11th International Wheat Genetics Symposium 2008 Proceedings Volume 2: 633-

Aston M.J. (1978): Differences in the Behavior of Adaxial and Abaxial Stomata of Aphistomathous Sunflower Leaves: Inherent or environmental? Plant and Physiol. 5:211-218. BALEŠEVIĆ-TUBIĆ S., OUKIĆ V., MARINKOVIĆ J., DOZET G., PETROVIĆ K. and TATIĆ M. (2011):

Importance of Microbiological Fertilizer, Used in Soybean Production: Agronomical and Biological Aspects. African Journal of Microbiology Research. Vol. 5 (27),4909-4916.

BELIMOV A. A., HONTZEAS N., SAFRONOVA V.I., DEMCHINSKAYA S.V., PILLUZZA G., BULLITTA S. and GLICK B.R. (2005): Soil Biol.Biochem. 37:241.

BIERHUIZEN J.F., BIERHUIZEN J.M. And MARTAKIS G.F.P. (1984): The effect of light and CO2 on photosynthesis of various pot plants. Gartenbauswissenschaft. 49. 215-257. BREWER C.A. (1992): Response by Stomata on Leaves to Micro Environmental Conditions. In.

Goldman, C.A. (ed.), Tested Studies for Laboratory Teaching. Proceeding of the 13th Workshop/Conference of the Association for Biology Laboratory Education (ABLE). 13. 67-77.

CABALLERO-MELLADO J., ONOFRE-LEMUS J., ESTRADA-DE LOS SANTHOS P. and MARTINEZ-AGULIAR L. (2007): The tomato Rhizosphere, an Environment Rich in Nitrogen-Fixing Burkholderia Species with Capabilities of Interest for Agriculture and Bioremediation. Applied and Environmental Microbiology. 73. 16: 5308-5319.

CAGLAR S. and TEKIN H. (1999): The stomata density of pistachio cultivars on different pistachio rootstocks. Turkish J. Agric. For. 23. 1029-1032. CAGLAR S., SUTYEMEZ M. and BEYAZIT S. (2004): Stomatal density in some selected walnut

(Juglans regia L.) types. J. Akdeniz Univ. Agric. Fac. 17. 169-174. CHAPIN F.S. (1991): Integrated responses of plants to stress. Bioscience, 41, 29-36.

DALE J.E. (1988): The control of leaf expansion. Annual Review of Plant Physiology an Plant DECOTEAU D.R. (2000): Vegetable Crops. Upper Rever Company. New Jersey. USA.

DOZENLI S. and ERGENOGLU F. (1991): Studies on the density of stomata of some Vitis vinifera L. varieties grafted on different rootstocks trained up various trellis systems. Tr. J. Forestry, 15.

DORDEVIĆ S., ĐURIĆ N., TRKULJA V., NAJDENOVSKA O., GAVRILOVIĆ, M. (2005a): The influence of seed bacterial inoculation on productivity of various wheat varieties. Zbornik naučnih radova.

DORDEVIC S., MILOŠEVIC N., JARAK M., NAJDENOVSKA O. (2005b): The effects of seed inoculation in maize by the bacteria of Bacillus sp. Zbornik naučnih radova. Vol. 11, br. 1-2. 95-102.

Olga NAJDENOVSKA et al .: BROCCOLI LEAF AND APPLICATION OF SLAVOL

ELSNER E.A., GERALD L. JUBB JR., (1988): Leaf Area Estimation of Concord Grape Leaves from Simple Linear Measurements, Amer. J. Enol. And Vitic, 39(1), 95-97.

FALLOVO C., CRISTOFORI V., DE-GYVES E.M., RIVERA C.M., REA R. and FANASCA S. (2008): Leaf Area Estimation Model for Small Fruits from Linear Measurements. Hort Science 43 (7): 2263-2267.

- FRITZ A. (1978): Foliar Fertilization a Technique for Improved Crop Production. V. Africa Symp. Hort, Crops. Acta Hort, 84: 150-161.
- GAMIELY S., RANDLE W.M., MILLS H.A., SMITTLE D.A. (1991): A Rapid And Nondestructive Method For Estimating Leaf Area Of Onions. HortScience, 26(2):206.
- GLICK B.R., KARATUROVIC D.M. and NEWELL P.C. (1995): A Novel Procedure for Rapid Isolation of Plant Growth Promoting Pseudomonas. Can. J. Microbiol. 41:533-536.

GLICK B.R. (1995): The Enhancement of Plant Growth by Free-living Bacteria, Canadian Journal of Microbiology, 41:109-117.

- GOVEDARICA M., MILOŠEVIĆ N., JARAK M., DORDEVIĆ S., DURIĆ S., STAMENKOVIĆ-JOVANOVIĆ S., RADANOVIĆ Z., NAJDENOVSKA O. (1999). Uticaj sojeva Azotobacter chroococcum na mikrobiološku aktivnost zemljištu i prinos šećerne repe. Savremena poljoprivreda, V. 48, 3-4. 95-101, Novi Sad.
- GOVEDARICA M., JELAČIĆ Z., JARAK M., MILOŠEVIĆ N., ERIĆ N. (2002); Bacterization as a Measure for Increasing Yield and Microbiological Activity in Corn. Collection of Scientific works. Institute PKB Agroekonomik. Vol 8. no. 1. 105-109.
- GRANIER C., TARDIEU F. (1998): Spatial and temporal analyses of expansion and cell cycle in sunflower leaves. Plant Physiology. 116. 991-1001.
- GULMON S.L., CHU C.C. (1981): The effects of light and nitrogen on photosynthesis leaf characteristics and dry matter allocation in the chaparral shrub, Diplacus auranticus, Occologia. 49. 207-212.
- JELACIĆ S., BEATOVIĆ D., LAKIĆ N. and VUJOŠEVIĆ A. (2007); The Effect of Natural Biostimulators and Slow-disintegrating Fertilizer on the Quality of Rosmary Seedlings (Rosmarinus officinalis L.) Journal of Agricultural Sciences. 52, 2. 85-94.
- JONES H.G. (1987): Breeding for stomatal characters. In: Stomatal Function, Zieger E., Farguhar G.D. and Cowan I.R. (eds). Stanford University Press. Stanford (CA) pp. 431-443.
- KANDIL H. and GAD N. (2009): Effects of Inorganic and Organic Fertilizers on Growth and Production of Broccoli (Brassica oleracea L.). Factori și Procese Pedogenetice din Zona Temperată 8 S. 61-69.
- KLOPPER J.W., LIFSHITZ R., ZABLOTOWICZ R.M. (1989): Free-living bacterial inocula for Enhancing Crop Productivity, Trends Biotechnol, 7:39-43.
- KOLOTO E. And OSINSKA M. (2001): Efficiency of foliar nutrition of field vegetables grown at different nitrogen rates. In: Proc. IC Environ. Probl. N-Fert.Acta Hort. 563: 87-91.

KURSCHNER W.M. (1997): The anatomical diversity of recent and fossil leaves of the durma diver oak (Ouercus petraea Lieblein/O. pseudocastanea Goeppert) - Implications for their use as biosensors of paleoatmospheric CO2 levels; Review of Paleobotany and Palynology, v. 96: 1-30.

- MEDINA E. (1970): Relationships between nitrogen level, photosynthetic capacity, and carboxydismutase activity in Atriplex patula levels. Carnegie Inst Wash Year Book. 69. 655-662.
- MENG Y., XU F. And MENG Q. (2008): Current Application Status and Prospect of Microbiological Fertilizer in China. Chines Agric, Sci. Bull. 24 (6): 276-283.
- MILIĆ V., HRUSTIĆ M., VASIĆ M., STARČEVIĆ LJ., MARINKOVIĆ J. (2003): Use of Microbiological Fertilizers in Bean, Soyabean and Maize Production. Field Veg. Crop Res. 38. 259-270.
- MISIMOVIĆ M., VUKOJEVIĆ D., ZAVIŠIĆ N., SIMIĆ J. and VIDIĆ M. (2012): Utecaj mikrobiološkog gnojiva Slavol na sprečavanje opadanja plodova jabuke pred berbu. Proceedings. 47th Croatian and 7th International Symposium on Agriculture. Opatija. Croatia. 807-811.
- MISIRLI A. and AKSOY U. (1994). A study on the leaf and stomatal properties of Sarilop fig variety. J. Ege Univ. Agric. Fac. 31. 57-63.

ZEMLJIŠTE I BILJKA, Vol. 62, No. 1, 1-13, 2013.

MOHSENIN N.N. (1986): Physon adBrical Properties of Plant and Animal Materials. Gordon and Breach Science Publishers. New York. NY. USA.

MOLNAR. I. (1995): Opšte ratarstvo. Biblioteka matice Srpske. Novi Sad.

- NAJDENOVSKA O., GOVEDARICA M., LAZAREVSKA S. (1998): Ulogata na azotobakterot vo zaštitata na kompirot od telenite crvi (Elateridae). Združenje za zaštita na rastenijata na Republika Makedonija. MAK-ISSN 1409-5084, Skopje. 191-197.
- NAJDENOVSKA O., JARAK M., DORĐEVIĆ S. (2002a): The influence of Azotobacter on chemical characteristics of potato bulbs. Mikrobiologija, vol. 39, No 1-2. 27-34, Beograd.
- NAJDENOVSKA O., GOVEDARICA M., ĐORĐEVIĆ S. (2002b): The influence of bacterization of potatoes on the number of ammonifers, dehydrogenase, potease and urease activity on soil. Mikrobiologija, vol. 39, No 1-2. 35-42, Beograd.
- NAJDENOVSKA O., MITKOVA T. AND DJURIĆ S. (2002c): The Influence of Azotobacter chroococcum on the Acticity under Potato Plant, Zemljište i biljka, Vol.51, No 3: 161-166, Beograd.
- NAJDENOVSKA O., GOVEDARICA M., MITKOVA T. DORDEVIĆ S. (2004). The Possibility of Nitrogenbacter croococcum Use in Soil Sown with Potatoes. Zemljište i biljka, Vol. 53, No 2, 137-146, Beograd.
- NAJDENOVSKA O. (2011): The influence of the insecticides and microbiological fertilizer on the number of microorganisms in the soil planted with potato. Zemljište i biljka. Vol. 60, No. 1 7-14, Beograd.
- NAJDENOVSKA O. (2012): The Influence of the Application of Microorganisms in the Production of Quality Potatoes: Zemljište i biljka. Vol. 61, 15-21, Beograd.
- NESMITH D.S. (1991): Nondestructive Leaf Area Estimation of Rabbiteye Blueberries. HortScience 26(10): 1332.
- NESMITH D.S. (1992): Estimating Summer Squash Leaf Area Nondestructively, HortScience 27(1): 77.
- RAJENDRAN P.C., THAMBURAJ S. (1987): Estimation of Leaf Area in Watermelon by Linear Measurements, South Indian Hort., 35(4): 325-327.
- RYUGO K. (1988): Fruit Culture: Its Science and Art. John Wiley and Sons Inc, New York. p. 344. SARIĆ M., STANKOVIĆ Z. and KRSTIĆ B. (1989): Fiziologija biljaka. Naučna knjiga. Novi Sad.
- SMART R.E. (1974): photosynthesis by grapevine canopics. Journal of Applied Ecology. Vol. 11. 3: 997-1006.
 STOLILKOVIĆ B. (1988): Uticaj komplesnih dubaju
- STOLILJKOVIĆ B. (1988): Uticaj komplesnih dubriva na prinos i kvalitetet krompira. Zemljište i biljka. Vol. 37, No. 1. 63-71, Beograd.
 STOPPANI M L. WOLD D. T. STOPPANI M L. ST
- STOPPANI M.I., WOLF R., FRANCESANGELI N. and MARTL H. (2003). A Non Destructive and Rapid Method for Estimating Leaf area of Broccoli. Adv Hort Sci 17. 173-175.

UZUN S. and KAR H. (2004): Quantitative Effects of Planting Time on Vegetative Growth of Broccoli (*Brassica oleraceae* var. *italica*). Pak. J. Bot. 36 (4): 769-777.

VESSEY J.K. (2003): Plant Growth Promoting Rhizobacteria as Bio-fertilizers. Plant Soil 255. 571-586.

VOLKENBURGH E. Van. (1999). Leaf expansion – an integrating plan behavior. Plant. Cell and Environment. 22. 1463-1473.

- WILLIAMS L. E. (1987): Growth of "Thompson Seedless" grapevines: Leaf Area Development and Dry Weight Distribution". Journal of American Society and Horticultural Science. Vol. 112. 2: 325-330.
- YILDIRIM E., GUVENC I., TURAN M. and KARATAS A. (2007): Effects of Foliar Urea Application on Quality, Growth, Mineral Uptake and Yield of Broccoli (Brassica oleracea L. v.r. italica)., Acta Hort. (ISHS)579. 249-253.
- YOLDAS F., CEYLAN B. YAGMUR B. and MORDOGAN N. (2008): Effect of nitrogen fertilizer on yield quality and nutrient content in broccoli. J. Plant Nutr. 31: 1333-1343.

Olga NAJDENOVSKA et al .: BROCCOLI LEAF AND APPLICATION OF SLAVOL

OPREDELENJE LISNE POVRŠINE I BROJ STOMA U BROKULI KORISTEČI RAZLIČITE METODE APLIKACIE MIKROBIOLOŠKOG ĐUBRIVA – SLAVOL

Olga NAJDENOVSKA¹, Snezana DJORDJEVIC², Elizabeta MISKOSKA-MILEVSKA¹, Daniela DIMOVSKA³ and Zoran POPOVSKI¹

¹Facultet za zemjodelski nauki i hrana – Skopje Poljoprivredni fakultet, Beograd-Zemun, Srbija ³Centar za primenteti istrazuvanja i permanentno obrazovanie vo zemjodelstvoto -Skopje

Izvod

Primena mikrobioloških dubriva u poljoprivredi ima pozitivan uticaj na biljnu proizvodnju, a isto tako rešava probleme vezane za zaštitu životne sredine. Trenutna istraživanja fokusiraju se na efekte mikrobioloških dubriva na žive organizme i zemljišta. Cilj istraživanja je bio da se ispita uticaj različitih metoda aplikacije mikrobiološkog dubriva - Slavol na lisnu površinu i broj stoma u brokuli "Verdia F1" (Brassica oleracea L. var. italica Plenck.). U istraživanju korišćene su dve različite metode aplikacije đubriva: folijarna aplikacija i aplikacija kroz vodu za navodnjavanje (sistem kap po kap). Ogled je postavljen na aluvijalan tip zemljišta na privatnom posedu u selu Jurumleri, u blizini Skoplja, Makedonija. Dobijeni rezultate pokazuju da je maksimalna prosečna površina lista bila 361.39 cm² u biljaka koje su bile folijarno tretirana sa Slavolom, dok minimalna prosečna površina lista bila je 167.86 cm² u kontroli. Lisna površina biljaka tretirane sa Slavolom kroz sistem za navodnjavanje bila je 262.68 cm². Prema LSD-testu, ovi podatci su pokazale statističku razliku (p<0.01). Može se zaključiti da metoda folijarne aplikacije Slavola daje najveće vrednosti za površinu lista.

> Primljeno 7. februara 2013. Odobreno 14. decembra 2013.

12