

GOCE DELCEV UNIVERSITY - STIP
FACULTY OF AGRICULTURE



JOURNAL OF AGRICULTURE AND PLANT SCIENCES

YEAR 2019

VOLUME 17, Number 2

EDITOR IN CHIEF

Editor in Chief

Liljana Koleva Gudeva, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, liljana.gudeva@ugd.edu.mk

Editors

Ljupco Mihajlov, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, ljupco.mihajlov@ugd.edu.mk

Fidanka Trajkova, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, fidanka.trajkova@ugd.edu.mk

Administrator

Biljana Atanasova, Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, biljana.atanasova@ugd.edu.mk

Technical Editing

Slave Dimitrov, Publishing Department, Goce Delcev University, Stip,
Republic of North Macedonia, slave.dimitrov@ugd.edu.mk

Language Editors

Biljana Ivanova, MA, Senior lecturer, Faculty of Philology, Goce Delcev University, Stip,
Republic of North Macedonia Macedonia, biljana.petkovska@ugd.edu.mk – English language Editor

Danica Gavrilovska Atanasovska, Goce Delcev University, Stip, Republic of North Macedonia
Macedonia, danica.gavrilovska@ugd.edu.mk – Macedonian language Editor

Editorial Office

Faculty of Agriculture, Goce Delcev University, Stip, Krste Misirkov Str., No.10-A P.O. Box 201, 2000
Stip, Republic of North Macedonia
japs@ugd.edu.mk
<http://js.ugd.edu.mk/index.php/YFA>

**GOCE DELCEV UNIVERSITY - STIP, REPUBLIC OF NORTH MACEDONIA
FACULTY OF AGRICULTURE**

UDC 63(058)
ISSN 2545-4447 print
ISSN 2545-4455 on line



Journal of Agriculture and Plant Sciences, JAPS, Vol 17, No. 2

Successor of the Yearbook of Faculty of Agriculture of GDU, Vol 14

YEAR 2019

VOLUME XVII, Number 2

EDITORIAL BOARD

Aco Kuzelov,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, aco.kuzelov@ugd.edu.mk

Danijela Raičević,

Biotechnical Faculty, University of Montenegro, Mihaila Lalica bb., Podgorica,
Montenegro, nelar@mail.com

Dragan Skorić,

Serbian Academy of Sciences and Arts, Knez Mihajlova 35, 11000 Belgrade,
Serbia, draganskoric@sbb.rs

Dragomir Vlcev,

Institute of Agriculture – Karnobat, Bulgaria, vulchevd@abv.bg

Hatice Gülen,

Istanbul Bilgi University, Faculty of Engineering and Natural Sciences,
Department of Genetics and Bioengineering, Istanbul, Turkey,
hatice.gulen@bilgi.edu.tr

Jovica Vasin,

Institute of Field and Vegetable Crops, Novi Sad, Serbia, jovica.vasin@ifvcns.ns.ac.rs

Kiril Bahcevandziev,

Coimbra Agricultural School, 3045-601 Coimbra, Portugal, kiril@esac.pt

Klemen Lisjak,

Agricultural Institute of Slovenia, Hacquetova ulica 17, Ljubljana, Slovenia, Klemen.Lisjak@kis.si

Marijan Bubola,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, marijan@iptpo.hr

Maryna Mardar,

Odessa National Academy of Food Technologies, Odessa, 65039, Kanatnaya Str., Ukraine,
marinamardar2003@gmail.com

Rubin Gulaboski,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, rubin.gulaboski@ugd.edu.mk

Sanja Radeka,

Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, sanja@iptpo.hr

Sasa Mitrev,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, sasa.mitrev@ugd.edu.mk

Shuhe Wei,

Institute of Applied Ecology, Chinese Academy of Sciences, China, shuhewei@iae.ac.cn
Velichka Nikolova Rodeva,
Maritsa Vegetable Crops Research Institute, 32 Brezovsko shosse St., Plovdiv 4003, Bulgaria,
velirod@yahoo.com

Violeta Dimovska,

Faculty of Agriculture, Goce Delcev University, Stip,
Republic of North Macedonia, violeta.dimovska@ugd.edu.mk

Wolfram Schnäckel,

Anhalt University of Applied Sciences, Bernburger Straße 55, 06366 Köthen, Germany,
Wolfram.Schnaeckel@hs-anhalt.de

CONTENT

Darko Andronikov, Aco Kuzelov, Julija Sazdova, Kiro Mojsov, Aco Janevski, Sonja Jordeva, Saška Golomeova Longurova QUANTITATIVE CHARACTERISTICS OF RABBIT HYBRIDS	9
Katerina Bojkovska, Nikolche Jankulovski, Goran Mihajlovski, Gordana Dimitrovska, Elena Joshevska CONSUMER PREFERENCES AS DETERMINANT OF THE SUCCESSFUL QUALITY MANAGEMENT OF DAIRY PRODUCTS	15
Daniela Dimovska, Ljupco Mihajlov, Fidanka Trajkova, Liljana Koleva Gudeva, Igor Iljovski THE INFLUENCE OF ORGANIC FERTILIZERS ON THE GROWTH AND YIELD OF BROCCOLI (<i>Brassica oleracea</i> L. var. <i>italica</i>)	25
Elena Joshevska, Aco Kuzelov, Gordana Dimitrovska, Katerina Bojkovska IMPROVEMENT OF THE PRODUCTION TECHNOLOGY OF MEAT PRODUCTS WITH THE ADDITION OF FUNCTIONAL COMPONENTS	31
Karolina Kocoska, Ilija Risteski THE SIZE AND NUMBER OF MIDDLE BELT LEAVES IN SOME VARIETIES OF WILD TOBACCO	39
Liljana Koleva Gudeva, Valentina Butleska Gjoroska, Fidanka Trajkova, Ljupco Mihajlov ACTIVITY OF ENZYME CATALASE IN ALFALFA (<i>Medicago sativa</i> L.) AS AN INDICATOR FOR ABIOTIC STRESS	45
Daniela Nikolovska Nedelkoska, Tatjana Kalevska, Natalija Atanasova-Pancevska, Mitko Karadelev, Zora Uzunoska, Dzoko Kungulovski EVALUATION OF BACTERICIDAL ACTIVITY OF SELECTED WILD MACROFUNGI EXTRACTS AGAINST <i>Escherichia coli</i>	53
Silvana Pashovska, Trajko Miceski PERSPECTIVES AND POSSIBILITIES FOR DEVELOPMENT OF TOBACCO PRODUCTION IN THE REPUBLIC OF NORTH MACEDONIA	59

INTRODUCTION

In the past twelve years the educational, research and applicative activities of the Faculty of Agriculture – Stip, Goce Delcev University – Stip, contributed to the development of agriculture sector in the country and broader region.

The Faculty of Agriculture organized the 2nd International Meeting Agriscience & Practice (ASP 2019), giving an opportunity to the participants for presentation and discussion of original scientific and practical results in different fields of agriculture.

The 2nd International Meeting Agriscience & Practice (ASP 2019), held on 12 April 2019 at the Faculty of Agriculture - Stip, was organized with intention to bring together all agricultural stakeholders for sharing their knowledge, experience and obstacles. One of the main aims was to link research and field work in agricultural sector in the country and broader, giving it an international dimension. All oral presentations as well as poster presentations at ASP 2019 were organized in several scientific sessions:

- Agricultural economics,
- Plant biotechnology,
- Plant production,
- Plant protection,
- Quality control and food safety,
- Soil science and hydrology,
- Viticulture, enology and fruit production.

The main goal of the Meeting was linking and promoting scientific achievements and practical knowledge, presented in different thematic areas, which were achieved in the Republic of North Macedonia and wider in the region.

Journal of Agriculture and Plant Sciences Vol. 17, No. 2 mostly contains the presented papers from the 2nd International Meeting Agriscience & Practice (ASP 2019).

Editorial Board,

December, 2019

Editor in chief,

Prof. Liljana Koleva Gudeva, PhD



THE INFLUENCE OF ORGANIC FERTILIZERS ON THE GROWTH AND YIELD OF BROCCOLI (*Brassica oleracea* L. var. *italica*)

Daniela Dimovska^{1*}, Ljupco Mihajlov¹, Fidanka Trajkova¹,
Liljana Koleva Gudeva¹, Igor Iljovski²

¹Faculty of Agriculture, Goce Delcev University, Krste Misirkov 10-A, P.O. box 201, 2000 Stip,
Republic of North Macedonia

²University Ss. Cyril and Methodius, Faculty of Agricultural Sciences and Food - Skopje, Blvd. Aleksandar
Makedonski b.b. P.O. Box 297, 1000 Skopje, Republic of North Macedonia

*Corresponding author: daniela.dimovska@ugd.edu.mk

Abstract

The experiment was conducted in order to determine the influence of organic fertilizers on the growth dynamics and yield of broccoli grown in the open field. The variety Verdija F1 cultivated in Skopje region during one year (2018) was used in the experiment. The treatments were as follows: Ø control - no use of organic fertilizer, V-1 - foliar treatment with Biohumus, V-2 - foliar treatment with Orgalife, V-3 - foliar treatment with Bio – Vital. The treatments were conducted every 10 days, starting on 19.08.2018, three times during vegetation period. The following traits were examined: dynamics of growth and broccoli yield. According to the results, the V-2 foliar treatment with the organic fertilizer Orgalife resulted in considerable higher yield on the central flower (6.98 t/ha) as compared to the control treatment (5.41 t/ha), proven with statistically significant difference, at the level of 0.05 with the LSD test. The results obtained for the average height of the plants in the three treatments showed that there are highly statistically significant differences at the level of 0.01 between the V-2 treatment (55.6 cm) and the control (48.7 cm). According to the overall results of the experiment, it can be concluded that the V-2 foliar treatment has given the best results in both traits as the plant height and the yield obtained.

Key words: growth dynamics, Biohumus, Orgalife, Bio – Vital

INTRODUCTION

Broccolis belong to the group of leafy vegetables from the family Brassicaceae; they are an important vegetable crop with high nutritional and good commercial value (Yoldas et al., 2008). They are food with a low concentration of salts, without fats and calories, rich in vitamin C and a good source of vitamin A, vitamin B2 and calcium (Decoteau, 2000). According to Apahidean (2011), the broccoli inflorescences (flowers) have great therapeutic value in the battle against cancerous diseases. Today, broccolis attract more attention because of their variety of uses and excellent nutritional value (Salunkhe and Kadam, 1998; Talalay and Fahey, 2001; Rangkadilok et al., 2002;). The broccoli production, along with the application of organic fertilizers, could serve as an alternative to mineral fertilizers (Gupta

et al., 1988; Wong et al., 1999; Yoldas et al., 2008; Farahzety and Siti Aishah, 2013) as well as to improve the soil structure (Bin, 1983;) and microbiological biomass (Suresh et al., 2004). However, it has been shown that liquid organic fertilizer as compost tea contains nitrogen mainly in an inorganic form such as ammonia (Price and Duddles, 1984; Gross et al., 2007) and make the nutrients immediately available to the plants. Bio-fertilizers play a multifunctional role, not only in the soil improvement process with microorganisms, but also as nutrients, stabilizers, hormones and isolates (Mohapatra et al., 2013). The application of organic (bio) fertilizers in the production of vegetables leads to higher plant productivity, quality improvement, early fruit growth and greater fruit durability.

MATERIAL AND METHODS

The experiment was set up near Skopje, in the village of Jurumleri, on alluvium soil type, during the cultivation season in 2018. Broccoli (*Brassica oleracea* L. var. *italica*), hybrid Verdia F1 was used as research material. The broccoli was cultivated from seedlings produced in cold beds, transplanted on open field. During the vegetation, the plants were treated with organic fertilizers Biohumus - organic fertilizers made from California worms in liquid (V-1 treatment), Orgalife - organic fertilizers made from California worms in liquid (V-2 treatment) and Bio - Vital microbiological fertilizer which contains bacteria in its composition (nitrifying microorganisms, phosphate-solubilizing microorganisms and cellulolytic microorganisms) (V-3 treatment). All fertilizers were used in concentration of 1% solution.

The treatments were set in three replications in the following order:

1. Ø control – no use of organic fertilizer,
2. V-1 – foliar treatment with Biohumus,
3. V-2 – foliar treatment with Orgalife,
4. V-3 – foliar treatment with Bio – Vital.

Each treatment was applied every 10 days. The beginning of the treatments started on 19.08, the second application of fertilizers was on 29.08 and the third application was performed on 09.09.

During the vegetation, dynamics of plant growth and yield were monitored. The dynamics of plant growth were monitored by measuring the height of the plants before each treatment.

The plant height was determined using a metro and the height was measured from the base to the top of plant. The height measurement was performed before each application of the organic fertilizers.

The yield per plant was determined only by the central inflorescence, by measuring them on an analytical scale and determining the arithmetic mean.

An analysis of - variance (ANOVA) for the two factors, the application timing and the fertilizer type for plant height and a single yield factor, was performed to determine the data obtained, and the smallest significant data ratios were determined by the LSD test at the level of $p = 0.05$ and $p = 0.01$.

RESULTS AND DISCUSSION

Plant height

It is widely known that the broccoli height is usually from 50 to 100 cm with spirally arranged leaves with an elongated shape, in the base more or less cut off depending on the type. In general, the plant is permeable, and the inflorescence is formed at the tip of the elongated trunk. In the generational phase, the broccoli tree grows from 50 to 90 cm (Lešić et al., 2002).

The results for the average plant height after applied treatments are given in Table 1. According to the results, the average plant height ranged from 32.4 cm in V-3 treatment at the first measurement to 67.6 cm, in V-2 treatment at the third measurement. The difference between the control (no use of organic fertilizer) and the treatments ranged from 6.9 cm (V-2 treatment) to 0.7 cm (V-3 treatment). The absolute dispersion was the highest in the control treatment and V-3 treatment, while the lowest was found for the V-1 treatment. The relative

placement of treatment data was the highest for the V-2 treatment with the highest plants (8%) and the lowest for the control treatment.

Statistical significance at the level of $p=0.01$ was obtained only in the V-2 treatment. While insignificant differences between the treatments were obtained in V-1 in terms of control and the V-3 variant, while the V-2 secondary variant varies considerably with all variants (V-2 and V-1) and control. The V-3 treatment did not show any significance regarding the treated, and the data was closest to the control variant.

The average height of plants ranged from 48.7 cm in control, at a V-3 height of 49.4 cm, at a V-1 height of 49.7 cm, and the highest average height had plants in V-2 of 55.6 cm.

The results obtained for average plant height from all three treatments showed high statistically significant differences of 0.01 between control and V-2.

Table 1. Average plant height (cm) after three applications of different treatments.

Treatments		∅	V-1	V-2	V-3
Application	I	32.8	33.7	39.7	32.4
	II	49.6	53.3	59.5	52.6
	III	63.7	62.2	67.6	63.3
±		48.7a	49.7ab	55.6c**	49.4a
+ - од ∅		∅	1.0	6.9	0.7
SD		15.5	13.6	14.3	15.7
CV (%)		7.5	6.9	8.0	7.7
LSD 0.05		2.82			
0.01		4.27			

SD – standard deviation; CV – coeficinet of variation

Significant differences at level * $p=0.05$, ** $p=0.01$, for treatment and a, b, c for applications.

The differences on plant height between the treatments and the control are shown in Figure 1. The plants in treatment V-2 were averagely 14% higher as compared to the plants in the control, 11% of treatment V-1 and 13% of

the V-3 treatments. Treatments V-1 showed 3% higher plants than the control and 2% of the treatments V-3, while plants of treatments V-3 were 1% higher than the control.




Figure 1. Average plant height (cm) and percentage of difference among different treatments.

Yield of central inflorescence in broccolis

Adding organic matter enhanced the soil structure conditions, creates conducive conditions for good root development (Arisha et al., 2003; Togun- and Akanbi, 2003) and mineralization by microorganisms. Hence, plants are able to get nutrients for higher yield (Radwan et al., 1993; El-Mansi et al., 1999; Wong et al., 1999; Abdelrazzag 2002; Al-Nasir 2002; Togun & Akanbi, 2003) upon application of organic fertilizers.

The harvesting of the inflorescences of the broccoli should be done before the inflorescences begin to open. Harvesting can

be one-off or multiple. According to Đurovka et al. (2008), the yield can range from 15 to 30 t/ha, including main and lateral inflorescence.

According to the results from our research regarding the central inflorescence yield, the highest average yield was obtained in V-2 treatment (6.98 t/ha) which was 1.57 t/ha higher yield as compared to the control. The lowest yield was recorded in V3 treatment (5.18 t/ha) and it was 0.23 t/ha lower than control. The standard deviation in all treatments was low, while the relative distribution was the highest in control (13.82%) and the lowest in V-3 treatment (7.54%).

Table 2. Broccoli yield (t/ha) after three applications of different treatments.

Variants		∅	V-1	V-2	V-3
Repetitions	I	6.25	7.05	6.96	4.64
	II	5.54	5.36	7.77	5.56
	III	4.43	6.84	6.19	5.34
±		5.41a	6.42b	6.98c*	5.18a
+ - ∅		∅	1.01	1.57	-0.23
SD		0.75	0.75	0.65	0.39
CV (%)		13.82	11.75	9.25	7.54
LSD 0.05		1.55			
0.01		2.25			

SD – standard deviation; CV – coeficinet of variation

Significant differences of level * $p=0.05$, ** $p=0.01$, for treatment and a, b, c for applications.

Significant difference ($p = 0.05$) between the organic fertilizer treatments and control was noted only for V-2 treatment. A significant difference was showed for V-1 treatment in relation to V-3 treatment and the control.

The average yield from all treatments and the control in our research was 6.01 t/ha, higher results showed V-2 (0.97 t/ha) and V-1 (0.41 t/ha), while the V-3 and control showed lower results than the average yield of all treatments (Figure 2).




Figure 2. Differences between the average yields in different treatments.

CONCLUSIONS

The application of organic fertilizers in the production of broccoli gave positive results, both for average plant height and yield per hectare. The application of the liquid organic fertilizer based on California worms, Orgalife, gave the best results based on the two examined parameters.

Organic fertilizer treatments gave averagely higher plants up to 14% in V-2, 3% in V-1 and 1% in V-3 as compared to the control plants.

According to the results, it can be concluded that the use of Orgalife contributed to a significantly higher yield on the central flower in treatment V-2 (6.98 t/ha) as compared to control (5.41 t/ha), treatment V-1 (6.42 t/ha) and treatment V-3 (5.18 t/ha). gave the best results based on the two examined parameters.

REFERENCES

- Abdelrazzag, A. (2002). Effect of chicken manure, sheep manure and inorganic fertilizers on yield and nutrient uptake by onion. *Pakistan Journal of Biological Science.*, 5, 266 – 268.
- Al-Nasir, F. (2002). Effect of organic fertilizers on yield and nutrients concentration of cauliflower plant. *Arch. Acker-Pfl. Boden.*, 48, 37 – 47.
- Apahidean, I.A. (2011). Research regarding cauliflower and broccoli cultivation in the environmental conditions of Transilvanyan tableland. Summary of PhD. Thesis. Cluj – Napoca.
- Arisha, H.M.E., Gad, A.A., & Younes, S.E. (2003). Response of some pepper cultivars to organic and mineral nitrogen fertilizer under sandy soil conditions. *Zagazig J. Agric. Res.*, 30, 1875 – 1899.
- Bin, J. (1983). Utilization of green manure for raising soil fertility in China. *Soil Sci.* 135, 65–9.
- Gross, A., Arusi, R., & Nejidat, A. (2007). Assessment of extraction methods with fowl manure for the production of liquid organic fertilizers. *Bioresource Technology*, 99, 327–334.
- Gupta, A.P., S.R. Antil, & Narwal, P.R. (1988). Effect of farmyard manure on organic carbon, available N and P contents of soil during different periods of wheat growth. *J. Indian Soil Sci.*, 36, 269–73.
- Decoteau, D.R. (2000). Vegetable Crops. Upper Rever Company. New Jersey, U.S.A.
- El-Mansi, A.A.A., Bardisi, A., Arisha, H.M.E., & Nour, E.M. (1999). Studies on some factors affecting growth and yield of pea under sandy soil conditions using drip irrigation system: 2. Effect of farmyard manure and irrigation water quality. *Zagazig J. Agric. Res.*, 26(5), 1409 – 1428.
- Yoldas, F., S. Ceylan, B. Yagmur, & Mordogan, N. (2008). Effect of nitrogen fertilizer on yield quality and nutrient content in broccoli. *J. Plant Nutr.*, 31, 1333–43.
- Lešić, R., Borošić, J., Buturac, I., Čustiċ, M., Poljak, M., & Romić, D. (2002). Povrċarstvo. Ćakovec.
- Mohapatra, S.K., Munsu, P. S., & Mahapatra, P.N. (2013). Effect of integrated nutrient management on growth, yield and economics of broccoli (*Brassica oleracea* L. Var. *italica* plenck.). *Vegetable Science*, 40(1), 69-72.
- Price, M.L., & Duddles, N. (1984). Chicken Manure Tea: Research Report. *Echo community*.
- Radwan, S.A., Aboel-fadl, M., & Abo-Hussien, E.A. (1993). Effect of salinity and organic material (farmyard manure) addition on growth and mineral contents of tomato in sandy calcareous soil of Egypt. *Menofiya J. Agric. Res.*, 18, 1929 – 1946.
- Rangkadilok, N., Nicolas M.E., Bennett R.N., Premier. R., Eagling D.R., & Taylor, P.W.J. (2002). Determination of sinigrin and glucoraphanin in Brassica species using a simple extraction method combined with ion-pair HPLC analysis. *Scientia Horticulturae*, 96, 27–41.
- Salunkhe, D.K., & Kadam, S.S. (1998). *Handbook of Vegetable Science and Technology: Production, Storage and Processing*, 1st edition. Marcel Dekker, Inc. Madison Avenue, New York, USA.
- Suresh, K.D., G. Sneha, K.K. Krishn, & Mool, C.M. (2004). Microbial biomass carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Archives Agron. Soil Sci.*, 50, 641–7.
- Talalay, P., & Fahey, J.W. (2001). Phytochemicals from Cruciferous plants protect against cancer by modulating carcinogen metabolism. *American Soc. Nutr. Sci.*, 23, 3027–33.
- Togun, A.O., & Akanbi, W.B. (2003). Comparative effectiveness of organic-based fertilizers to mineral fertilizers on tomato growth and fruit yield. *Compost Sci. and Utilization*, 11, 337 – 342.
- Farahzety, A.M., & Siti Aishah, H. (2013). Effects of organic fertilizers on performance of cauliflower (*Brassica oleracea* var. *botrytis*) grown under protected structure. *J. Trop. Agric. and Fd. Sc.*, 41 (1), 15– 25.
- Đurovka M., Laziċ B., Bajkin A., Potkonjak A., Markoviċ V., Ilin Ź., & Todoroviċ, V. (2008). Proizvodnja povrċa i cveċa u zaštićenom prostoru. Univerzitet u Novom Sadu, Poljoprivredni fakultet.
- Wong, J.W.C., Ma, K.K., Fang, K.M., & Cheung, C. (1999). Utilization of manure compost for organic farming in Hong Kong. *Bio-resource Technol.*, 67, 43 – 46.

ВЛИЈАНИЕТО НА ОРГАНСКИТЕ ЃУБРИВА НА РАСТОТ И ПРИНОСОТ НА БРОКУЛА (*Brassica oleracea* L. var. *italica*)

Даниела Димовска^{1*}, Љупчо Михајлов¹, Фиданка Трајкова¹,
Лилјана Колева-Гудева¹, Игор Иљовски²

¹Земјоделски факултет, Универзитет „Гоце Делчев“ - Штип, ул. „Крсте Мисирков“ 10-А, п. фах 201, 2000
Штип, Република Северна Македонија

²Универзитет „Св. Кирил и Методиј“, Факултет за земјоделски науки и храна - Скопје, бул. „Александар
Македонски“ бб, п. фах 297, Република Северна Македонија

*Контакт автор: daniela.dimovska@ugd.edu.mk

Резиме

Експериментот беше спроведен со цел да се утврди влијанието на органските ѓубрива врз динамиката на растот и приносот на брокула одгледувана на отворено. Беа користена сортата *Verdija F1*, која е одгледувана во Скопскиот регион во текот на 2018 година. Беа користени следниве третмани со органски ѓубрива: Ø контрола - без употреба на органско ѓубриво, V-1 - фолијарен третман со Biohumus, V-2 - фолијарен третман со Orgalife, V-3 - фолијарен третман со Bio - Vital. Третирањето е изведувано на секои 10 дена, почнувајќи од 19.8.2018 г. и тоа три пати во текот на вегетацијата. Целта на истражувањата беше да се испита влијанието на соодветните третмани врз динамиката на растот и приносот кај брокулата. Според добиените резултати може да се утврди дека со користење на органски ѓубрива Orgalife е добиен значително повисок принос на централниот цвет со третманот V-2 (6.98 t/ha) во споредба со контролниот третман (5.41 t/ha). Статистички значајна разлика, на ниво од 0,05 со LSD-тест, беше добиена помеѓу V-2 и контролата. Резултатите добиени за просечната висина на растенијата, за време на трите третмани, покажаа дека постои високостатистички значајна разлика на ниво од 0,01 помеѓу V-2 (55.6 cm) и контролата. Според добиените резултати може да се заклучи дека V-2 даде најдобри резултати и во зголемувањето на растот на брокулата и во приносот.

Клучни зборови: динамика на раст, Biohumus, Orgalife, Bio – Vital.