

UDK 37

ISSN 2545 - 4439
ISSN 1857 - 923X

INTERNATIONAL JOURNAL

Institute of Knowledge Management

KNOWLEDGE



Scientific Papers

MEDICAL SCIENCES AND HEALTH

Vol. 38.4

KIJ

Vol. 38

No. 4

pp. 639 - 1010

Skopje 2020

Global Impact & Quality Factor 1.822 (2017) <http://globalimpactfactor.com/knowledge-international-journal/>

KNOWLEDGE



INTERNATIONAL JOURNAL

**SCIENTIFIC PAPERS
VOL. 38.4**

March, 2020

**INSTITUTE OF KNOWLEDGE MANAGEMENT
SKOPJE**



KNOWLEDGE
International Journal Scientific Papers Vol. 38.4

ADVISORY BOARD

Vlado Kambovski PhD, Robert Dimitrovski PhD, Siniša Zarić PhD, Maria Kavdanska PhD, Venelin Terziev PhD, Mirjana Borota – Popovska PhD, Cezar Birzea PhD, Veselin Videv PhD, Ivo Zupanovic, PhD, Savo Ashtalkoski PhD, Zivota Radosavljević PhD, Laste Spasovski PhD, Mersad Mujevic PhD, Milka Zdravkovska PhD, Drago Cvijanovic PhD, Predrag Trajković PhD, Lazar Stosic PhD, Krasimira Staneva PhD, Nebojsa Pavlović PhD, Daniela Todorova PhD, Baki Koleci PhD, Lisen Bashkurti PhD, Zoran Srzentić PhD, Sinisa Opic PhD, Marija Kostic PhD

Print: GRAFOPROM – Bitola

Editor: IKM – Skopje

Editor in chief

Robert Dimitrovski, PhD

KNOWLEDGE - International Journal Scientific Papers Vol. 38.4

ISSN 1857-923X (for e-version)

ISSN 2545 – 4439 (for printed version)

INTERNATIONAL EDITORIAL BOARD

President: Academic, Prof. Vlado Kambovski PhD, Skopje (N. Macedonia)

Vice presidents:

Prof. Robert Dimitrovski PhD, Institute of Knowledge Management, Skopje (N. Macedonia)

Prof. Sinisa Zanic, PhD, Faculty of Economics, University of Belgrade, Belgrade (Serbia)

Prof. Venelin Terziev PhD, University of Rousse, Rousse (Bulgaria)

Prof. Mersad Mujevic PhD, Public Procurement Administration of Montenegro (Montenegro)

Prof. Tihomir Domazet PhD, President of the Croatian Institute for Finance and Accounting, Zagreb (Croatia)

Members:

- Prof. Aleksandar Korablev PhD, Faculty of economy and management, Saint Petersburg State Forest Technical University, Saint Petersburg (Russian Federation)
- Prof. Azra Adajalic – Dedovic PhD, Faculty of criminology and security, Sarajevo (Bosnia & Herzegovina)
- Prof. Anita Trajkovska PhD, Rochester University (USA)
- Prof. Anka Trajkovska-Petkoska PhD, UKLO, Faculty of technology and technical sciences, Bitola (N. Macedonia)
- Prof. Alisabri Sabani PhD, Faculty of criminology and security, Sarajevo (Bosnia & Herzegovina)
- Prof. Aneta Mijoska PhD, Faculty of Dentistry, University “St. Cyril and Methodius”, Skopje (N. Macedonia)
- Prof. Ahmad Zakeri PhD, University of Wolverhampton, (United Kingdom)
- Prof. Ana Dzumalieva PhD, South-West University “Neofit Rilski”, Blagoevgrad (Bulgaria)
- Prof. Aziz Pollozhani PhD, Rector, University Mother Teresa, Skopje (N.Macedonia)
- Prof. Artan Nimani PhD, Rector, University of Gjakova “Fehmi Agani” (Kosovo)
- Prof. Branko Sotirov PhD, University of Rousse, Rousse (Bulgaria)
- Prof. Branko Boshkovic, PhD, College of Sports and Health, Belgrade (Serbia)
- Prof. Branimir Kampl PhD, Institute SANO, Zagreb (Croatia)
- Prof. Baki Koleci PhD, University Hadzi Zeka, Peja (Kosovo)
- Prof. Branislav Simonovic PhD, Faculty of Law, Kragujevac (Serbia)
- Prof. Bistra Angelovska, Faculty of Medicine, University “Goce Delcev”, Shtip (N.Macedonia)
- Prof. Cezar Birzea, PhD, National School for Political and Administrative Studies, Bucharest (Romania)
- Prof. Cvetko Andreevski, Faculty of Tourism, UKLO, Bitola (N.Macedonia)
- Prof. Drago Cvijanovic, PhD, Faculty of Hotel Management and Tourism, University of Kragujevac, Vrnjacka Banja (Serbia)
- Prof. Dusan Ristic, PhD Emeritus, College of professional studies in Management and Business Communication, Novi Sad (Serbia)
- Prof. Darijo Jerkovic PhD, Faculty of Business Economy, University “Vitez”, (Bosnia & Herzegovina)
- Prof. Daniela Todorova PhD, “Todor Kableshev” University of Transport, Sofia (Bulgaria)
- Prof. Dragan Kokovic PhD, University of Novi Sad, Novi Sad (Serbia)
- Prof. Dragan Marinkovic PhD, High health – sanitary school for professional studies, Belgrade (Serbia)
- Prof. Daniela Ivanova Popova PhD, Faculty of Public Health and Sport, SWU Neofit Rilski, Blagoevgrad (Bulgaria)

-
- Prof. Dzulijana Tomovska, PhD, Faculty of Biotechnical sciences, Bitola (N.Macedonia)
 - Prof. Evgenia Penkova-Pantaleeva PhD, UNWE -Sofia (Bulgaria)
 - Prof. Fadil Millaku, PhD, Rector, University “Hadzi Zeka”, Peja (Kosovo)
 - Prof. Fatos Ukaj, University “Hasan Prishtina”, Prishtina (Kosovo)
 - Prof. Georgi Georgiev PhD, National Military University “Vasil Levski”, Veliko Trnovo (Bulgaria)
 - Prof. Halit Shabani, PhD, University “Hadzi Zeka”, Peja (Kosovo)
 - Prof. Halima Sofradzija, PhD, University of Sarajevo, Sarajevo (Bosnia and Herzegovina)
 - Prof. Haris Halilovic, Faculty of criminology and security, University of Sarajevo, Sarajevo (Bosnia and Herzegovina)
 - Prof. Helmut Shramke PhD, former Head of the University of Vienna Reform Group (Austria)
 - Prof. Hristina Georgieva Yancheva, PhD, Rector, Agricultural University, Plovdiv (Bulgaria)
 - Prof. Hristo Beloev PhD, Bulgarian Academy of Science, Rector of the University of Rousse (Bulgaria)
 - Prof. Hristina Milcheva, Medical college, Trakia University, Stara Zagora (Bulgaria)
 - Prof. Izet Zeqiri, PhD, Academic, SEEU, Tetovo (N.Macedonia)
 - Prof. Ivan Marchevski, PhD, D.A. Tsenov Academy of Economics, Svishtov (Bulgaria)
 - Prof. Ibrahim Obhodjas PhD, Faculty of Business Economy, University “Vitez”, (Bosnia & Herzegovina)
 - Doc. Igor Stubelj, PhD, PhD, Faculty of Management, Primorska University, Koper (Slovenia)
 - Prof. Ivo Zupanovic, PhD, Faculty of Business and Tourism, Budva (Montenegro)
 - Prof. Ivan Blazhevski, PhD, Institute for Sociological, Political and Juridical Research, Skopje (N.Macedonia)
 - Prof. Ivan Petkov PhD, European Polytechnic University, Pernik (Bulgaria)
 - Prof. Isa Spahiu PhD, AAB University, Prishtina (Kosovo)
 - Prof. Ivana Jelik PhD, University of Podgorica, Faculty of Law, Podgorica (Montenegro)
 - Prof. Islam Hasani PhD, Kingston University (Bahrein)
 - Prof. Jamila Jaganjac PhD, Faculty of Business Economy, University “Vitez”, (Bosnia & Herzegovina)
 - Prof. Jova Ateljevic PhD, Faculty of Economy, University of Banja Luka, (Bosnia & Herzegovina)
 - Prof. Jove Kekenovski PhD, Faculty of Tourism, UKLO , Bitola (N.Macedonia)
 - Prof. Jonko Kunchev PhD, University „Cernorizec Hrabar“ - Varna (Bulgaria)
 - Prof Karl Schopf, PhD, Akademie fur wissenschaftliche forchung und studium, Wien (Austria)
 - Prof. Katerina Belichovska, PhD, Faculty of Agricultural Sciences, UKIM, Skopje (N. Macedonia)
 - Prof. Krasimir Petkov, PhD, National Sports Academy “Vassil Levski”, Sofia (Bulgaria)
 - Prof. Kamal Al-Nakib PhD, College of Business Administration Department, Kingdom University (Bahrain)
 - Prof. Kiril Lisichkov, Faculty of Technology and Metallurgy, UKIM, Skopje (N.Macedonia)
 - Prof. Krasimira Staneva PhD, University of Forestry, Sofia (Bulgaria)
 - Prof. Lidija Tozi PhD, Faculty of Pharmacy, Ss. Cyril and Methodius University, Skopje (N.Macedonia)
 - Prof. Laste Spasovski PhD, Vocational and educational centre, Skopje (N.Macedonia)
 - Prof. Larisa Velic, PhD, Faculty of Law, University of Zenica, Zenica (Bosnia and Herzegovina)

-
- Prof. Łukasz Tomczyk PhD, Pedagogical University of Cracow (Poland)
 - Prof. Lujza Grueva, PhD, Faculty of Medical Sciences, UKIM, Skopje (N.Macedonia)
 - Prof. Lazar Stosic, PhD, Association for development of science, engineering and education, Vranje (Serbia)
 - Prof. Lulzim Zeneli PhD, University of Gjakova “Fehmi Agani” (Kosovo)
 - Prof. Lisen Bashkurti PhD, Global Vice President of Sun Moon University (Albania)
 - Prof. Lence Mircevska PhD, High Medicine School, Bitola, (N.Macedonia)
 - Prof. Ljupce Kocovski PhD, Faculty of Biotechnical sciences, Bitola (N.Macedonia)
 - Prof. Marusya Lyubcheva PhD, University “Prof. Asen Zlatarov”, Member of the European Parliament, Burgas (Bulgaria)
 - Prof. Marija Magdinceva – Shopova PhD, Faculty of tourism and business logistics, University “Goce Delchev”, Shtip (N. Macedonia)
 - Prof. Maria Kavdanska PhD, Faculty of Pedagogy, South-West University Neofit Rilski, Blagoevgrad (Bulgaria)
 - Prof. Maja Lubenova Cholakova PhD, Faculty of Public Health and Sport, SWU Neofit Rilski, Blagoevgrad (Bulgaria)
 - Prof. Mirjana Borota-Popovska, PhD, Centre for Management and Human Resource Development, Institute for Sociological, Political and Juridical Research, Skopje (N.Macedonia)
 - Prof. Mihail Garevski, PhD, Institute of Earthquake Engineering and Engineering Seismology, Skopje (N.Macedonia)
 - Prof. Misho Hristovski PhD, Faculty of Veterinary Medicine, Ss. Cyril and Methodius University, Skopje (N.Macedonia)
 - Prof. Mitko Kotovchevski, PhD, Faculty of Philosophy, UKIM, Skopje (N.Macedonia)
 - Prof. Milan Radosavljevic PhD, Dean, Faculty of strategic and operational management, Union University, Belgrade (Serbia)
 - Prof. Marija Topuzovska-Latkovikj, PhD, Centre for Management and Human Resource Development, Institute for Sociological, Political and Juridical Research, Skopje (N.Macedonia)
 - Prof. Marija Knezevic PhD, Academic, Banja Luka, (Bosnia and Herzegovina)
 - Prof. Margarita Bogdanova PhD, D.A.Tsenov Academy of Economics, Svishtov (Bulgaria)
 - Prof. Mahmut Chelik PhD, Faculty of Philology, University “Goce Delchev”, Shtip (N.Macedonia)
 - Prof. Marija Mandaric PhD, Faculty of Hotel Management and Tourism, University of Kragujevac, Vrnjacka Banja (Serbia)
 - Prof. Marina Simin PhD, College of professional studies in Management and Business Communication, Sremski Karlovci (Serbia)
 - Prof. Miladin Kalinic, College of professional studies in Management and Business Communication, Sremski Karlovci (Serbia)
 - Prof. Marijan Tanushevski PhD, Macedonian Scientific Society, Bitola (N. Macedonia)
 - Prof. Mitre Stojanovski PhD, Faculty of Biotechnical sciences, Bitola (N.Macedonia)
 - Prof. Miodrag Smelcerovic PhD, High Technological and Artistic Vocational School, Leskovac (Serbia)
 - Prof. Nadka Kostadinova, Faculty of Economics, Trakia University, Stara Zagora (Bulgaria)
 - Prof. Natalija Kirejenko PhD, Faculty For economic and Business, Institute of Entrepreneurial Activity, Minsk (Belarus)
 - Prof. Nenad Taneski PhD, Military Academy “Mihailo Apostolski”, Skopje (N.Macedonia)
 - Prof. Nevenka Tatkovic PhD, Juraj Dobrila University of Pula, Pula (Croatia)
 - Prof. Nedžad Korajlic PhD, Dean, Faculty of criminal justice and security, University of Sarajevo (Bosnia and Herzegovina)
 - Prof. Nonka Mateva PhD, Medical University, Plovdiv (Bulgaria)

- Prof. Nikolay Georgiev PhD, “Todor Kableshkov” University of Transport, Sofia (Bulgaria)
- Prof. Nishad M. Navaz PhD, Kingdom University (India)
- Prof. Nano Ruzhin PhD, Faculty of Law, FON University, Skopje (N.Macedonia)
- Prof. Oliver Dimitrijevic PhD, High medicine school for professional studies “Hipokrat”, Bujanovac (Serbia)
- Prof. Paul Sergius Koku, PhD, Florida State University, Florida (USA)
- Prof. Primoz Dolenc, PhD, Faculty of Management, Primorska University, Koper (Slovenia)
- Prof. Predrag Trajkovic PhD, JMPNT, Vranje (Serbia)
- Prof. Petar Kolev PhD, “Todor Kableshkov” University of Transport, Sofia (Bulgaria)
- Prof. Pere Tumbas PhD, Faculty of Economics, University of Novi Sad, Subotica (Serbia)
- Prof. Rade Ratkovic PhD, Faculty of Business and Tourism, Budva (Montenegro)
- Prof. Rositsa Chobanova PhD, University of Telecommunications and Posts, Sofia (Bulgaria)
- Prof. Rossana Piccolo PhD, Università degli studi della Campania - Luigi Vanvitelli (Italy)
- Prof. Rumen Valcovski PhD, Imunolab Sofia (Bulgaria)
- Prof. Rumen Stefanov PhD, Faculty of public health, Medical University of Plovdiv (Bulgaria)
- Prof. Rumen Tomov PhD, Rector, University of Forestry, Sofia (Bulgaria)
- Prof. Sasho Korunoski PhD, UKLO, Bitola (N.Macedonia)
- Prof. Snezhana Lazarevic, PhD, College of Sports and Health, Belgrade (Serbia)
- Prof. Stojan Ivanov Ivanov PhD, Faculty of Public Health and Sport, SWU Neofit Rilski, Blagoevgrad (Bulgaria)
- Prof. Stojna Ristevska PhD, High Medicine School, Bitola, (N. Macedonia)
- Prof. Suzana Pavlovic PhD, High health – sanitary school for professional studies, Belgrade (Serbia)
- Prof. Sandra Zivanovic, PhD, Faculty of Hotel Management and Tourism, University of Kragujevac, Vrnjacka Banja (Serbia)
- Prof. Shyqeri Kabashi, College “Biznesi”, Prishtina (Kosovo)
- Prof. Temelko Risteski PhD, Faculty of Law, FON University, Skopje (N. Macedonia)
- Prof. Todor Krystevich, D.A. Tsenov Academy of Economics, Svishtov (Bulgaria)
- Prof. Todorcka Atanasova, Faculty of Economics, Trakia University, Stara Zagora (Bulgaria)
- Doc. Tatyana Sobolieva PhD, State Higher Education Establishment Vadiym Getman Kiyev National Economic University, Kiyev (Ukraine)
- Prof. Tzako Pantaleev PhD, NBUniversity, Sofia (Bulgaria)
- Prof. Violeta Dimova PhD, Faculty of Philology, University “Goce Delchev”, Shtip (N. Macedonia)
- Prof. Vojislav Babic PhD, Institute of Sociology, University of Belgrade (Serbia)
- Prof. Volodymyr Denysyuk, PhD, Dobrov Center for Scientific and Technological Potential and History studies at the National Academy of Sciences of Ukraine (Ukraine)
- Prof. Valentina Staneva PhD, “Todor Kableshkov” University of Transport, Sofia (Bulgaria)
- Prof. Venus Del Rosario PhD, Arab Open University (Philippines)
- Prof. Vjollca Dibra PhD, University of Gjakova “Fehmi Agani” (Kosovo)
- Prof. Yuri Doroshenko PhD, Dean, Faculty of Economics and Management, Belgorod (Russian Federation)
- Prof. Zlatko Pejkovski, PhD, Faculty of Agricultural Sciences, UKIM, Skopje (N.Macedonia)
- Prof. Zivota Radosavljevic PhD, Faculty FORCUP, Union University, Belgrade (Serbia)
- Prof. Zorka Jugovic PhD, High health – sanitary school for professional studies, Belgrade (Serbia)

REVIEW PROCEDURE AND REVIEW BOARD

Each paper is reviewed by the editor and, if it is judged suitable for this publication, it is then sent to two referees for double blind peer review.

The editorial review board is consisted of 63 members, full professors in the fields 1) Natural and mathematical sciences, 2) Technical and technological sciences, 3) Medical sciences and Health, 4) Biotechnical sciences, 5) Social sciences, and 6) Humanities from all the Balkan countries and the region.

CONTENTS

XANTHOGRANULOMATOUS EPIDIDYMO-ORCHITIS AS A UNCOMMON AND RARE PSEUDONEOPLASTIC CONDITION	657
Ilbert Ademi.....	657
Adnan Vrajnko	657
Majlinda Ademi.....	657
CORRELATION BETWEEN THE POST 5ARI SYNDROME AND DEHIDROEPIANDROSTERONE LEVELS IN PATIENTS WITH BPH	663
Maja Sofronievska Glavinov	663
IMPACT OF COMPLEX PHARMACOLOGICAL AND NON-PHARMACOLOGICAL TREATMENT IN PATIENTS WITH CHRONIC PROSTATITIS/CHRONIC PELVIC PAIN SYNDROME	671
Galina Mratskova	671
Rumen Deliev	671
PREVENTION OF THROMBOPHILIA AND THE FOLLOWING PREGNANCY COMPLICATIONS	679
Anna Mihaylova	679
Petar Uchikov	679
Ekaterina Uchikova	679
Nikoleta Parahuleva.....	679
THE CORELATION BETWEEN FETO-PLACENTAL UNIT, FETAL BIOMETRY AND AMNIOTIC FLUID IN PREGNANT WOMEN WITH PREECLAMPSIA IN SECOND TRIMESTER	685
Pranvera Izairi.....	685
Kristina Skeparovska.....	685
Nevenka Velickova.....	685
LABORATORY DIAGNOSIS OF INFECTIONS CAUSED BY CHLAMYDIA TRACHOMATIS, EXPERIENCES FROM THE PHF CLINICAL HOSPITAL “M.D. TRIFUN PANOVSKI” – BITOLA FOR THE PERIOD FROM 2013 TO 2019	691
Blagica Krsteska	691
Dino Karpicarov	691
Biljana Gjorgjeska	691
PROLONGED NEONATAL JAUNDICE – FREQUENCY AND CAUSE OF ITS OCCURRENCE... 697	697
Tatyana Itova	697
Victorya Atanasova	697
ANATOMICAL CHARACTERISTICS OF THE VERTEBRAL ARTERY USING CT ANGIOGRAPHY	701
Svetlana Jovevska.....	701
Merima Idrizovikj.....	701
APPLICATION OF SPECIALIZED METHODS OF PFYSIOTHERAPY ON THE FLEXIBLE MUSCLE GROUP IN PATIENTS WITH COXARTROSIS.....	705
Zhulieta Gerenova	705
ANKYLOSING SPONDYLITIS (MORBUS BECHTEREW).....	711
Merima Idrizovikj.....	711
Svetlana Jovevska.....	711
METHODS FOR NEUROMUSCULAR REEDUCATION FOR REDUCING MOTOR DEFICIT AND MUSCLE SPASM IN PEOPLE WITH CEREBRAL PALSY.....	717
Lence Nikolovska	717
Zana Markova.....	717
KINESITHERAPY EFFECT IN PATIENTS WITH SHOULDER BURSITIS	723
Danche Vasileva	723
Suat Berisha.....	723

EPIDEMIOLOGICAL CHARACTERISTICS OF ACUTE VIRAL HEPATITIS TYPE C IN MACEDONIA	729
Velo Markovski	729
Vlatko Cvetanovski	729
EPIDEMIOLOGICAL CHARACTERISTICS OF MEASLES IN MACEDONIA DURING THE PERIOD 2010-2019	735
Kolevska Natasha	735
Bosevska Golubinka	735
Zdravkovska Milka	735
Markovski Velo	735
NURSING PROCESS IN TRAUMATIC CHEST INJURY OF THE HEART - A CLINICAL CASE..	741
Lilyana Stancheva.....	741
Ivanka Stambolova	741
Tsvetelina Evtimova	741
HEALTH-RELATED QUALITY OF LIFE TO PEOPLE WITH CHRONIC RESPIRATORY AND CARDIOVASCULAR DISEASES	747
Katya Popova.....	747
Galina Terzieva.....	747
Monika Obreykova	747
SELF-CONTROL OF DISEASE IN PATIENTS WITH ARTERIAL HYPERTENSION	753
Lyubomir Marinov	753
Pavlinka Dobrilova	753
PATIENTS WITH CHRONIC DISEASE AWARENESS ABOUT THE OPPORTUNITIES FOR PREVENTIVE INTERVENTIONS	759
Pavlinka Dobrilova	759
PHARMACOTHERAPY APPROACH IN KIDNEY CANCER THERAPY	767
Sanela Andova.....	767
Tijana Serafimovska	767
Milka Zdravkovska.....	767
Trajan Balkanov	767
Marija Darkovska Serafimovska	767
ANTHROPOLOGY IN ORTHODONTICS	773
Natasa Toseska-Spasova.....	773
Natasha Stavreva	773
Biljana Dzipunova	773
DISCOLORATION OF NANO COMPOSITE RESIN BY EVERYDAY DRINKS: IN VITRO STUDY	779
Aleksandar Andreevski.....	779
Ivona Kovacevska.....	779
METHODS AND TECHNIQUES FOR BLEACHING VITAL TEETH	785
Natasa Longurova.....	785
Ivona Kovacevska.....	785
PROSTHODONTIC SOLUTION OF FLABBY RIDGES USING WINDOW IMPRESSION TECHNIQUE.....	791
Natasha Stavreva	791
Natasha Tosheska Spasova.....	791
THE DIGITAL IMPRESSION AS AN INITIAL UNIT OF MODERN DENTAL MEDICINE - LITERATURE REVIEW	797
Dobromira Shopova.....	797
Miroslava Yordanova	797
Svetlana Yordanova.....	797

PROSTHETIC TREATMENT WITH PLATE IN PATIENT WITH MAXILLARY RESECTION	803
Ivan Gerdzhikov	803
RISKS DURING ORTHODONTIC TREATMENT.....	807
Ana Radeska-Panovska	807
DRUG SUPPLY CHAIN SAFETY - FIGHT AGAINST COUNTERFEIT DRUGS	813
Elena Drakalska Sersemova	813
Bistra Angelovska.....	813
Dijana Miceva.....	813
Elmedina Sakir	813
MONITORING OF THE RENAL FUNCTION IN PIROXICAM TREATED PATIENTS WITH HEADACHE.....	819
Drita Yzeiri Havziu	819
Biljana Gjorgjeska	819
Edita Alili Idrizi.....	819
Gjylaj Alija	819
Sihana Lika.....	819
Merita Dauti.....	819
SESAME ALLERGY: A GROWING FOOD ALLERGY OF GLOBAL PROPORTIONS?	825
Verica Jakjimoska.....	825
Biljana Gjorgjeska	825
REHABILITATION IN PATIENTS WITH TYPE 2 DIABETES	831
Alexandra Yankova Maleva	831
DEPRESSION AND DIABETES	837
Aneta Spasovska Trajanovska	837
Danijela Janicevic Ivanovska	837
Zora Mitic.....	837
FINAL ANALYSIS OF THE RESULTST IN MAN WITH MULTIPLE SCULPROSIS AFTER A PHYSIOTHERAPEUTIC PROGRAM.....	843
Inna Ivanova	843
PSYCHOSOMATIC DISORDERS IN ADOLESCENTS	849
Lubomira Dimitrova.....	849
CHEMICAL CORROSIVE DISEASES OF THE EYE.....	853
Gazepov Strahil	853
Viktorija Micevska	853
Alen Gorgiev	853
Slavena Stojkova	853
HEALTH BEHAVIOR AND PARENTS' AWARENESS HOW TO PROTECT THE CHILDREN'S EYE HEALTH.....	859
Krassimira Dimitrova	859
Christina Grupcheva.....	859
CONTROL MATERIALS-MEDICAL DEVICES FOR INCREASING THE QUALITY OF CLINICAL LABORATORY ANALYSES	865
Denitsa Trancheva	865
LABORATORY QUALITY MANAGEMENT.....	871
Katerina Tosheska-Trajkovska	871
MEDICAL SUPPLIES FOR CHEMICAL ACCIDENTS AND TERRORIST ATTACKS WITH CHLORINE IN THE PHARMACY NETWORK IN VARNA	877
Maria Panteleeva	877
Nicolina Radeva	877
Rostislav Kostadinov.....	877

AWARENESS AND ATTITUDES OF NURSING STAFF REGARDING TO EVALUATION IN HOSPITAL	883
Daniela Stankeva	883
Galina Tchaneva	883
Milena Stoycheva	883
HEALTH CARE CENTERS IN BULGARIA – NECESSITY AND CHALLENGES	889
Ivanka Stambolova	889
Stefan Stambolov	889
NURSING CARE ASSESSMENT AND NURSES' PROFESSIONAL ATTITUDES	895
Galya Yankova	895
CREATION, DEVELOPMENT AND PERSPECTIVES OF NURSING EDUCATION IN BULGARIA	903
Silviya Mladenova	903
THE ROLE OF THE FAMILY NURSE IN THE IMPLEMENTATION OF IMMUNIZATION AND EDUCATION OF THE POPULATION IN THE MUNICIPALITY OF KOCANI FOR THE PERIOD 2015-2019	909
Gordana Panova	909
Christina Janeva	909
STRESS AND CRISIS IN THE FAMILY IN CHILDREN WITH AUTISM AND DOWN SYNDROME	915
Plamen Tzokov	915
TRAINING AND ADDITIONAL QUALIFICATIONS OF SPECIALISTS WORKING IN THE FIELD OF CHILD REHABILITATION IN ORDER TO IMPROVE THE QUALITY OF THE SERVICES OFFERED	923
Stela Marinkova	923
Antoaneta Terzieva	923
ROLE OF THE PHYSICAL ACTIVITY AND REHABILITATION IN THE PROCESS OF AGING	929
Vanina Mihaylova	929
Ina Ivanova	929
Adolf Alakidi	929
Kristina Kilova	929
Mariana Liochkova	929
TREATMENT OF PATIENTS WITH FIBROMYALGIA WITH METHODS OF TRADITIONAL CHINESE MEDICINE	937
Lence Nikolovska	937
Bojan Radinski	937
THE ROLE OF THE PATRONAGE NURSE IN CONDUCTING AND EDUCATING THE ELDERLY	943
Hristina Trajkovska	943
Gordana Panova	943
THE CHALLENGES OF THE ELECTROMAGNETIC RADIATION- ARE WE READY FOR THEM - A MINI REVIEW	949
Vassilka Ilieva	949
TELEMEDICINE, TELEHEALTHCARE AND TELEHEALTH – LITERATURE REVIEW	957
Stanka Uzunova	957
Kristina Kilova	957
MEDICAL DEVICES AND ADAPTED MEANS IN OCCUPATIONAL THERAPY PROFESSION	965
Yoana Negrova	965
SOCIAL SIGNIFICANT ASPECTS OF COMBUSTION INJURY	971
Anushka Dimitrova	971

CORRELATION BETWEEN THE POST 5ARI SYNDROME AND DEHYDROEPIANDROSTERONE LEVELS IN PATIENTS WITH BPH

Maja Sofronievska Glavinov

University Surgical Clinic “ST.Naum Ohridski”, Skopje, Republic of North Macedonia
msofronievska@yahoo.com

Abstract: Conservative or non-surgical treatment of benign prostatic hyperplasia relies on the use of mainly two groups of medicines: selective alpha blockers and 5 alpha reductase inhibitors (5ARI). The most efficient non-surgical treatment is combination therapy of these medicines but occasionally after the introduction of 5ARI in the treatment, certain adverse effects that include sexual disturbances and depression may appear that are known as "post-5ARI syndrome". The purpose of this study is to show the significance and role of the dehydroepiandrosterone as an important predictor of post-5ARI syndrome in patients with BPH who are treated with 5ARI. In this randomized controlled prospective pharmaco epidemiological study, we investigated 120 patients with BPH combination therapy (tamsulosin and dutasteride) versus control group that consisted of 130 patients with monotherapy for BPH (tamsulosin). The correlation between both erectile dysfunction and depression on one side and serum dehydroepiandrosterone levels on the other was investigated at the time of active use of therapy in period of 6 months. We evaluated two groups of patients through 3 questionnaires: IPSS (International Prostate Score System) to evaluate the symptoms of benign prostatic hyperplasia (BPH), IIEF (International Index for Erectile Function) questionnaire for erectile dysfunction (ED) and a Beck depression inventory (BDI). We concluded that there is a law-proportionate correlation between appearance of post 5ARI syndrome and dehydroepiandrosterone levels in patients with BPH.

Keywords: BPH, erectile dysfunction, depression, dehydroepiandrosterone, post 5ARI

1. INTRODUCTION

Post 5 ARI syndrome initially known as “postfinasteride syndrome” since 2012 relates to all adverse effects of 5 alpha reductase inhibitors (5ARI), mainly erectile dysfunction and depressive disorders¹. The basis of its occurrence is the inhibition of 5 alpha reductases enzymes thus blocking the conversion of testosterone to dihydrotestosterone (DHT) as well as catalyzing the biosynthesis of powerful neurosteroids from their precursor dehydroepiandrosterone. Analysis of the literature indicates that after 2 years of treatment with 5ARI, approximately 12% of patients must discontinue treatment with these medicines due to adverse effects, particularly erectile dysfunction². Research also suggests that 5ARI not only inhibits the process of conversion of testosterone to dihydrotestosterone, but they also cause a reduction in nitric oxide synthetase (NOS) activity, and therefore the erection of cavernous bodies in these individuals is compromised³. The 5 alpha reductase inhibitors are a group of drugs that not only enable conversion of testosterone to dihydrotestosterone but also enable the conversion of other steroids and neurosteroids that act as antiepileptic, antidepressants and anxiolytics to brain tissue. In that way 5 ARI are responsible for the production of several neuroactive steroids⁴.

From the physiological point of view both testosterone and dihydrotestosterone also modulate the neuroendocrine stress response and are inversely proportional to depressive disorders⁵. The neurosteroids such as alopregnanolones are produced in the brain with mediation of 5-alpha reductase enzyme and shows lower values in men with

¹ Foundation P-FS. *Overview*. (2013). <http://www.pfsfoundation.org/post-finasteride-syndrome-overview/> on 23.12.2019

² Byung Hoon Chi, Sae Chul Kim. *Changes in Sexual Function in Benign Prostatic Hyperplasia Patients Taking Dutasteride: 1-Year Follow-Up Results* www.kjurology.org <http://dx.doi.org/10.4111/kju.2011.52.9.632> on 23.12.2019

³ Seo SI, Kim SW, Paick JS. *The effects of androgen on penile reflex,erectile response to electrical stimulation and penile NOS activity in the rat*. *Asian J Androl*,1:169-74 (1999)

⁴ Celec P, Ostatnikova D, Hodosy J. *On the effects of testosterone on brain behavioral functions*. *Front Neurosci*.,9(12):1-17 (2015)

⁵ Barrett-Connor E, Von Muhlen DG, Kritz-Silverstein D. *Bioavailable testosterone and depressed mood in older men: the Rancho Bernardo Study*. *J Clin Endocrinol Metab*,84 (2):573-577 (1999).

depressive symptoms⁶. Patients with clinical depression have also been shown to have lower values of 5-alpha reductase enzymes in the prefrontal cortex⁷.


Dehydroepiandrosterone (DHEA) is steroid precursor and its sulfated depot in the human body is DHEA-S, excreted by zona reticularis of the adrenal gland. It also has been proven both as a direct stimulator to NOS activity in the endothelium and as neuromodulator in the brain. Genetic studies have shown that there is a precisely specific receptor (G-protein) on the plasma membrane to which DHEA binds. This DHEA receptor is functionally linked to all three subtypes of G-protein, which mediate the activation of nitric oxide synthetase (NOS). The released NO afterwards, causes a series of intracellular reactions, the ultimate goal of which is to decrease intracellular calcium which results in relaxation of the smooth muscle of the blood vessels and establishing erections of cavernous bodies⁸. Several studies have reported a reduced serum level of DHEA in patients with erectile dysfunction (ED), which suggests that decreased DHEA and DHEA-S secretion are important risk factors for ED in aging men⁹.

These lower levels of serum DHEA in addition with decreased levels of DHT in BPH patients, raises the possibility of appearance of post 5ARI syndrome in those patients who are treated with 5ARI.

2. MATERIAL AND METHODS

In this randomized controlled prospective pharmaco-epidemiological study, a total of 250 outpatients with diagnosis of BPH were followed for 6 months in a specialist urological clinic. The research procedures were in line with ethical standards and the 1975 Helsinki Declaration, which was revised in Seoul in 2008. Patients were divided in two groups: control group that consisted of 130 patients treated with monotherapy for BPH (tamsulosin) and investigated group of 120 patients treated with BPH combination therapy (tamsulosin and dutasteride). The correlation of post 5ARI syndrome on one side and serum dehydroepiandrosterone levels on the other, were investigated at the time of active use of medicinal anti BPH therapy, in period of 6 months after starting the therapy.

Graph 1. Bivariant histogram of patients with BPH by age and therapy they receive; a control group (M) and an investigated group (D)



We evaluated two groups of patients through 3 questionnaires: IPSS (International Prostate Score System) in order to evaluate the symptoms of benign prostatic hyperplasia (light, medium or severe), IIEF-5 (International Index for erectile function-5) to determine erectile dysfunction (ED) and a Beck depression inventory to evaluate patients' psychological status.

Inclusion criteria for the study were men with diagnosis of BPH aged 45 to 70 years, prostate volume equal to or greater than 40cm³, prostate specific antigen (PSA) less than or equal to 4ng/ml, IPSS from 12 to 26, IIEF-5 equal to or greater than 21 and Beck depression index (BDI) equal to or greater than 11.

⁶ Melcangi RC, Caruso D, Abbiati F, et al. *Neuroactive steroid levels are modified in cerebrospinal fluid and plasma of post-finasteride patients showing persistent sexual side effects and anxious/depressive symptomatology.* J Sex Med.10(10):2598-2603(2013)

⁷ Hammond GL, Hirvonen J, Vihko R. *Progesterone, androstenedione, testosterone, 5 alpha-dihydrotestosterone and androsterone concentrations in specific regions of the human brain.* J Steroid Biochem, 18(2):185-189 (1983).

⁸ Liu, D., Dillon JS. *Dehydroepiandrosterone activates endothelial cell nitric-oxide synthase by a specific plasma membrane receptor coupled to Galpha(i2,3).* J Biol Chem, 14, 277(24), 1379-1388 (2012).

⁹ Morales, A. & Heaton, JP. *Andropause: a misnomer for a true clinical entity.* J Urol, 163, 705-712 (2002).

Exclusion criteria were dementia, deafness, medical history of mental disorders, vascular diseases, and myocardial infarction and prostate cancer.

Diagnosis of BPH was established by routine analysis of prostate volume by transabdominal echosonography, prostate specific antigen (PSA) determination, and lower urinary tract symptoms (LUTS) evaluated by IPSS. At the same time serum levels of dehydroepiandrosterone were measured in the same laboratory between 8 and 9 a.m in all 250 patients at the point of introduction of anti BPH medicinal therapy.

After the patients signed consent to use the data for the study, the urologist determined the IPSS score and the degree of ED through a questionnaire and gave them a self-assessment questionnaire for depression i.e BDI. Self-evaluation is considered to provide a more objective insight into the degree of depression, as the patient is not ashamed to give an objective response when self-evaluating, which would be the case if confronted with a physician. The urologist notes and interprets the final score of the questionnaires.

3. RESULTS

The data obtained during the survey were entered into a database and processed with the Statistica 7 program. The score obtained from the IPSS questionnaire classifies all patients with BPH into three groups: patients with mild (1-7), moderate (8-19) and severe (20-35) symptoms as showed in table 1.

Table 1: Number of patients classified according the lower urinary tract symptoms and IPSS

IPSS	N° of patients	Percentage of patients%	Description of symptom score
(7-15)	117	46,8	mild
(16-20)	106	42,4	moderate
(21-35)	27	10,8	severe
Total	250		


The IIEF-5 erectile function questionnaire contains 5 self-evaluation questions interpreted by an urologist. According to the obtained score, patients were classified in five groups: no ED (22-25), minor ED (17-21), mild ED (12-16), moderate ED (8-11), severe ED (5-7) as shown in table 3.

Table 2: Number of patients classified according to IIEF-5 for erectile dysfunction at the beginning of the study

IIEF-5	N° of patients	Percentage of patients%	Description of symptom score
20-25	249	99,6	NO ED
17-21	1	0,4	Minor ED
12-16	0		Mild ED
8-11	0	0	Moderate ED
5-7	0	0	Severe ED

As shown in graph 1.erectile function at the point of starting BPH medicinal therapy ranged from 17 to 25 in the control group (1a) which means that none of these patients suffered from impaired erectile function. As for the examined groip (D) at the beginning of the therapy patients had satisfying erectile function as well ranging from 17 to 25. After 6 months of BPH therapy (1b), patients in the control group (M) remain with satisfying erectile function, while there was a decreasing of erectile function index in some patients of the examined group (D). Total of 26 patients in the examined group had IIEF score from 4 to 12 points and experienced ED.

Graph 2: Categorized histogram of IIEF-5 in control group (M) and examined group (D) at the beginning of the study (a) and after 6 months of receiving anti BPH therapy (b)




We evaluated the symptoms of depression by administering a Beck Depression Inventory containing 21 items and self-evaluating patients. Scoring and interpretation of the results were done by an urologist, according to whom patients were classified as: normal oscillations (1-10), moderate oscillations (11-16), borderline clinical depression (17-20), moderate depression (21-30) and severe depression (> 40) as shown in table 2.

Table 3: Number of patients classified according to IEFF-5 for erectile dysfunction at the beginning of the study

BDI	N° of patients	Percentage of patients%	Description of symptom score
1-10	227	90,8	Normal oscillations
11-16	23	9,2	Physiologic oscillations
17-20	0	0	Borderline clinical depression
21-30	0	0	Moderate clinical depression
31-40	0	0	Severe depression
> 40	0	0	Extreme depression

As shown in graph 3. BDI index at the point of starting BPH medicinal therapy ranged from 4 to 18 in the control group (1a) which means that none of these patients suffered from depressive disorder. As for the examined group (D) at the beginning of the therapy only few patients had mild mood disorders due to impaired sleep because of the symptoms of BPH. After 6 months of BPH therapy (1b), patients in the control group (M) had an improve of the psychological status due to improvement of BPH symptoms and overnight sleep, while there was a worsening of BDI index in some patients of the examined group (D). Total of 26 patients in the examined group had BDI score from 20 to 30 points and expressed clinical signs of depression.


Graph 3: Categorized histogram of BDI in control group (M) and examined group (D) at the beginning of the study (a) and after 6 months of receiving anti BPH therapy (b)



From the analysis of the obtained data, we made a brake throw the insights of the impact of anti BPH therapy on the sexual function and psychological changes in patients with BPH and correlation of their appearance and the serum levels of DHEA. Processing consists of tabular and graphical representation of statistical series according to defined variables.

The serum level of DHEA at the beginning of the study ranged from 15 to 290 in both control and examined group as seen from Graph 4.

Graph 4: Serum DHEA levels at the beginning of the study (a) and after 6 months of the study(b) in control group (M) and in examined group (D)



With regard to DHEA values in both groups of patients, as shown in graph 2, it is evident that there were no significant differences in the DHEA serum values of the two groups at the beginning of the study.

After 6 months of BPH therapy serum DHEA measurements showed no significant oscillations compared to the original taken at the beginning of the study (graph 4).

In order to determine the possible correlation of erectile dysfunction with DHEA levels, a Spearman rank correlation was performed. It showed a significance ($p < 0.05$) in the correlation of dehydroepiandrosterone levels with erectile dysfunction and mood disorders.

Patients who had low levels of DHEA (below physiological limits) in the study showed lower IIEF-5 score and expressed ED. The results of this correlation are shown in Table 4.

Table 4: Correlation of DHEA levels with erectile function

	IIEF_0 m.	IIEF_6 m.
DHEA_0	-,0126	,4400
	p=,843	p=,000
DHEA_6	,0049	,4450
	p=,939	p=,000

DHEA levels throughout the study showed a significant correlation ($p < 0.05$) with a rise in the Beck Depression Index and mood disorders. The values of this correlation are shown in Table 5.

Table 5: Correlation of testosterone and DHEA levels with Beck's depression index

	BDI_0m.	BDI_6m.
DHEA_0	,0628	-,3954
	p=,323	p=,000
DHEA_6	,0599	-,4011
	p=,345	p=,000

The above results indicate that serum DHEA levels are significantly correlated with erectile function and psychological status in the patients included in this study.

4. DISCUSSION

Summarized data from a large number of clinical trials that were processed during the development of this study clearly showed that some patients with BPH, after pharmacological treatment with 5ARI, experienced adverse reactions such as erectile dysfunction and mood disorders.

The role of DHEA/ DHEA-S is still unclear but several studies have reported a reduced serum level of DHEA / DHEA-S in patients with ED, meaning that reduced synthesis of these pro hormones is a significant risk factor in the development of ED in older men. Reduced levels of DHEA-S, especially in young men with ED, may be an etiological factor for the occurrence of ED. However, patients treated with DHEA had a statistically significant increase in all domains of IIEF-5 compared to the placebo group⁹.

The prevalence of depressive symptoms in the Polish cohort study of patients with BPH, PolSenior (2007-2012), revealed that only 13.6% of the elderly included in the study were initially diagnosed with depression. Following the introduction of 5ARI in the treatment of BPH, the incidence of mood disorder and DS in BPH patients in their study was approximately 26%, excluding patients who previously had a depressive condition¹⁰. This is in support of the hypothesis in this study that after the introduction of 5ARI in the treatment of BPH, there is a possibility of post-5ARI syndrome, and its symptoms correlate with each other and change the quality of life of those patients in whom they occur.

The most relevant finding is the association between BPH pharmacotherapy and DSs occurrence is demonstrated in a study that shows that the use of a 5ARI is associated with a 1.52-fold higher prevalence of depressive symptoms¹¹. It is in line with a study showing that the use of finasteride, a 5ARI might induce development of depression¹².

¹⁰ Szybalska, A., Broczek, K., Slusarczyk, P., et al *Utilization of medical rehabilitation services among older Poles: results of the PolSenior study*. European geriatric medicine, 9(5), 669–677 (2018).

¹¹ Pietrzyk, B., Olszanecka-Glinianowicz, M., Owczarek, A., et al. *Depressive symptoms in patients diagnosed with benign prostatic hyperplasia*. Int Urol Nephrol, 47(3), 431-440(2015).

¹² Rahimi-Ardabili, B., Pourandarjani, R., Habibollahi, P., Mualeki, A. *Finasteride induced depression: a prospective study*. BMC clinical pharmacology 6, 7 (2006).

There is experimental evidence for the direct effects of DHEA on the brain. The elucidation of the role of DHEA in sexual function is confirmed by the administration of DHEA which results in the biosynthesis of active androgens that act on tissue target targets without repercussions at their level in the peripheral circulation¹³.

5. CONCLUSION

In this study it is concluded that 5ARI therapy has a negative effect on erectile function and psychological health only in a limited number of patients with BPH, (2 out of 10 BPH patients), receiving combination therapy of alpha 1 blocker and dutasteride.

The analysis in this study provided conclusion that patients with negative effects of 5ARI (dutasteride) i.e the expression of post-5 ARI syndrome have lower serum levels of DHEA-S that deviate from normal, physiological age ranges. In a situation where conversion of testosterone to its more potent androgen DHT is inhibited by the action of 5ARI, DHEA/DHEA-S acts as a secondary conditional androgen and is the only steroid that can express its positive effect on erectile and psychological function. The reason lies in the fact that 5 alfa reductase enzymes does not have any effect in the biochemical changes of DHEA and its sulfurated depot, DHEA-S. In a condition where DHEA/DHEA-S serum values are within physiological limits they have a direct impact in both erection of cavernose bodies and act as neurosteroids in the brain. This study showed that when DHEA-S values were lower than normal, physiologically ranges for the appropriate age, in patients treated with 5ARI, BPH patients expressed a post-5ARI syndrome.

REFERENCES

- Agis-Balboa, R.C., Guidotti, A., & Pinna, G., (2014). 5 α -reductase type I expression is downregulated in the prefrontal cortex/Brodman's area 9 (BA9) of depressed patients. *Psychopharmacology (Berl)*, 231(17), 3569-3580
- Barrett-Connor, E., Von Muhlen, D.G., & Kritz-Silverstein, D., (1999). Bioavailable testosterone and depressed mood in older men: the Rancho Bernardo Study. *J Clin Endocrinol Metab.*,84 (2), 573-577.
- Celec, P., Ostatnikova, D., Hodosy, J., (2015). On the effects of testosterone on brain behavioral functions. *Front Neurosci.*, 9(12), 1-17.
- Dimitropoulos, K., & Gravas, S. (2016). Fixed-dose combination therapy with dutasteride and tamsulosin in the management of benign prostatic hyperplasia. *Therapeutic advances in urology*, 8(1), 19–28.
- Edwards, J.E., & Moore, R.A., (2002). Finasteride in the treatment of clinical benign prostatic hyperplasia: a systematic review of randomised trials. *BMC Urol*, 2, 14.
- Dunphy, C., Laor L., Te, A., Kaplan, S., & Chughtai, B. (2015). Relationship Between Depression and Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. *Reviews in urology*, 17(2), 51-57.
- Hammond, G.L., Hirvonen, J., & Vihko, R., (1983). Progesterone, androstenedione, testosterone, 5 alpha-dihydrotestosterone and androsterone concentrations in specific regions of the human brain. *J Steroid Biochem*, 18(2), 185-189.
- Labrie, F., Belanger, A., Cusan, L., & Candas, B., (1997). Physiological changes in dehydroepiandrosterone are not reflected by serum levels of active androgens and estrogens but of their metabolites: intracrinology. *J Clin Endocrinol Metab*, 82, 2403-2409.
- Liu, D., & Dillon JS. (2002). Dehydroepiandrosterone activates endothelial cell nitric-oxide synthase by a specific plasma membrane receptor coupled to Galpha(i2,3). *J Biol Chem*, 14, 277(24), 1379-1388.
- Melcangi, R.C., Caruso, D., Abbiati, F., et al. (2013). Neuroactive steroid levels are modified in cerebrospinal fluid and plasma of post-finasteride patients showing persistent sexual side effects and anxious/depressive symptomatology. *J Sex Med*, 10(10), 2598-2603
- Morales, A. & Heaton, JP. (2000). Andropause: a misnomer for a true clinical entity. *J Urol*, 163, 705-712.
- Pietrzyk, B., Olszanecka-Glinianowicz, M., Owczarek, A., et al. (2015). Depressive symptoms in patients diagnosed with benign prostatic hyperplasia. *Int Urol Nephrol*, 47(3), 431-440
- Rahimi-Ardabili, B., Pourandarjani, R., Habibollahi, P., Mualeki, A. (2006). Finasteride induced depression: a prospective study. *BMC clinical pharmacology* 6, 7. doi:10.1186/1472-6904-6-7 on 16.12.2019

¹³ Labrie, F., Belanger, A., Cusan, L., Candas, B. *Physiological changes in dehydroepiandrosterone are not reflected by serum levels of active androgens and estrogens but of their metabolites: intracrinology.* *J Clin Endocrinol Metab*, 82, 2403-2409 (1997).