

ZBORNIK SAŽETAKA 2022 OKTOBAR 20-21



# 21. KONGRES STOMATOLOGA SRBIJE

STOMATOLOŠKA KOMORA SRBIJE





# 20. Kongres stomatologa Srbije

30. septembar – 3. oktobar 2021




## **SPONZORI**





































# 21. KONGRES STOMATOLOGA SRBIJE



20-21. oktobar 2022.

EP35	FRACTURE RESISTANCE AT ALL-CERRAMIC RESTORATIONS IN RELATION TO THE TYPE OF PREPARATION
	Natasha Stavreva, Natasha Toseska Spasova, Biljana Dzipunova, Efka Zabokova Bilbilova, Olga Kokocheva Ivanovska, Tanja Stefanoska, Muhamed Bajrami, Kristina Mitik96
EP36	PREVENTIVE METHODS FOR OCCLUSAL SURFACE OF THE FIRST MOLAR Tanja Stefanoska, Efka Zabokova-Bilbilova, Slavica Najdenoska Bojchinoska, Natasha Stavreva, Danica Stojanovska, Vasilka Trendafilova Strezova, Julijana Stevanandjija98
EP37	PREPARACIJA INSTRUMENTACIONE PUTANJE PRIMENOM RUČNIH I MAŠINSKIH INSTRUMENATA
	Dr Miljan Stošić, Ass. Dr Milica Jovanović-Medojević100
EP38	ORALNO ZDRAVLJE KOD DECE PREDŠKOLSKOG UZRASTA dr Tanja Stefanović, dr Branka Virijević, dr Sanja Radojević, Milena Stanišić Rašić101
EP39	PRESENCE OF PERIODONTOPATHOGENS IN THE ORAL MUCOSAL LESIONS Uroš Tomić DDM, Đorđe Mihailović DDM, Sanja Petrović DDM PHD, Drago Jelovac MD DDM MSc PhD, Nađa Nikolić PhD, Ana Pucar DDS MSc PhD102
EP40	OCCLUSION: UNDERSTANDING OR MISUNDERSTANDING Vesna Trpevska, Ivan Tanatarec, Daniela Srbinoska, Aneta Mijoska, Olga Kokoceva- Ivanovska104
EP41	THE TIME DURATION OF INSTRUMENTATION WITH PROTAPER UNIVERSAL NITI HAND SYSTEM, PROTAPER NEXT NITI ROTARY SYSTEM AND PROTAPER GOLD ROTARY SYSTEM  Verica Toneva Stojmenova, Natasha Longurova, Ivona Kovacevska, Sonja Rogoleva
	Gjurovski, Julija Zarkova Atanasova105
EP42	USE OF CHLORHEXIDINE IN RESTORATIVE DENTISTRY AND ENDODONTICS Longurova Natasha, Toneva Stojmenova Verica, Kovachevska Ivona



# 21. KONGRES STOMATOLOGA SRBIJE

PREDAVANJA PO POZIVU

# 21. KONGRES STOMATOLOGA SRBIJE



20-21. oktobar 2022.

# EP42 USE OF CHLORHEXIDINE IN RESTORATIVE DENTISTRY AND ENDODONTICS

### Longurova Natasha, Toneva Stojmenova Verica, Kovachevska Ivona

 $\label{thm:condition} \mbox{Univerzity Goce Delchev Shtip} \ , \mbox{R.N.Macedonia Faculty of medical sciences}, \mbox{Dental medicine} \ \ \,$ 

Nowadays, caries and periodontal diseases are the two dominant diseases of the oral cavity, and dental plaque is the main factor leading to their development and progression. The control of dental plaque, its accumulation and development is the basis for the fight against caries.

The aim of this research is to define the mechanism of the antimicrobial effect of chlorhexidine, its use in preventive and restorative dentistry, the advantages and disadvantages during endodontic therapy, as well as to review its negative properties and toxicity. For the realization of this paper, was made a research of scientific and professional literature, clinical studies and presentation of cases obtained by searching and analyzing Pubmed and Research Gate.

**Results**: The antimicrobial spectrum of activity of CHX includes a large number of microorganisms such as Gram+ and Gram bacteria, bacterial spores, lipophilic viruses (including HBV), fungi, especially dermatophytes. Preventing the accumulation of plaque, it becomes an anti plaque and anti gingivitis agent. Regarding its use in endodontics as an irrigant, the most commonly used protocol today involves irrigation with a 2.5% 6% sodium hypochlorite (NaOCI) solution supplemented with 2% CHX In addition to its antibacterial effect, chlorhexidine is also used during the placement of composite restorations in order to inactivate matrix adhesive systems used to place composite restorations. Chlorhexidine is used in such situations for complete inhibition of the enzyme MMP. After etching with orthophosphoric acid, the cavity is washed with water and dried, and then is applied a 2% solution of chlorhexidine. That concentration is sufficient to inhibit the enzyme.

**Conclusion**: When it comes to modern research on chlorhexidine, it is widely used and can act as a bacteriostatic (0.02% 0.06%) or bactericidal (0.12%), depending on the dose and the bacterial species, where in the same dose in some species it shows a bacteriostatic effect, and in others, a bactericidal effect.

