



**fdi**  
FDI World Dental Federation

**DORU**

**DENTAL Plus+**  
IMPLANT KA



**TİKA**  
TURKISH COOPERATION AND COORDINATION AGENCY



# 27<sup>th</sup> Congress of the **Balkan Stomatological Society (BaSS)**

## “ Past, Present and Future of Dental Implants and Gerodontology ”

November 9-11, 2023  
Istanbul Atlas University  
Istanbul, Türkiye



[www.e-bass.org/27thcongress](http://www.e-bass.org/27thcongress)



**ISTANBUL ATLAS UNIVERSITY**  
Atlas Vadi Kampüsü,  
Anadolu Caddesi  
No:40 Kağıthane  
Istanbul - TÜRKİYE  
[www.atlas.edu.tr](http://www.atlas.edu.tr)



**ORGANIZATION SECRETARIAT**  
**Topkon Congress & Event Management (PCO)**  
Zuhtupasa Mah. Rifatbey Sok. No:24  
34724 Kalamis-Kadıköy-Istanbul / TÜRKİYE  
Phone: +90 216 330 90 20 (pbx)  
Fax: +90 216 330 90 05  
E-mail: [bass2023@topkon.com](mailto:bass2023@topkon.com)



fdi  
FDI World Dental Federation

DORUK

DENTAL Plus+  
IMPLANT KA



TİKA  
TÜRKİYE İÇİŞİLERİ BAKANLIĞI



# 27<sup>th</sup> Congress of the Balkan Stomatological Society (BaSS)

“ Past, Present and Future of Dental Implants and Gerodontology ”

November 9-11, 2023 Istanbul Atlas University / Istanbul, Türkiye

## Committee

### President of the 27th BaSS Congress

Prof. Dr. A. Bülent KATİBOĞLU

### Scientific Committee

#### President of the Scientific Committee of the 27th BaSS Congress

Prof. Dr. Tamer Lütfi ERDEM

#### 27. BaSS Congress Scientific Committee Vice President

Prof. Dr. Nuriye Emel DERVİŞ

#### 27. BaSS Congress Members of the Scientific Committee

Prof. Dr. Mehmet Kamil GÖKER

Prof. Dr. Nurgül KÖMERİK

Prof. Dr. Ruzhdie QAFMOLLA

Prof. Dr. Dejan MARKOVIC

Prof. Dr. Athanasios POULOPOULOS

Prof. Dr. Ana MINOVSKA

Prof. Dr. Aleksa MARKOVIC

Prof. Dr. Paula PERLEA

Prof. Dr. Norina FORNA

Assoc. Prof. Konstantinos ARAPOSTATHIS

Assoc Prof. Edit XHAJANKA

Dr. Mihael STANOJEVIĆ

Assoc. Prof. Smiljka ČIČMIĆ

Assoc. Prof. Seda AYDEMİR MUTLU

Assoc. Prof. Muammer Çağrı BURDURLU

Assoc. Prof. Şirin HATİPOĞLU

Dr. Marijan DENKOVSKI

Dr. Jale TUNÇER

Dr. Canan DUMAN

Dr. Oktay DÜLGER

Dr. Ayşe KARKAÇ

Dr. Nilay BUDAK

Dr. Tuğçe BÖREKÇİ

Dr. Gamze KAVUNCU

Dr. Berna İZMİRLİ EVRENOL

Dr. Selim ÇÖMELEKOĞLU

Dr. Zeynep ÖZTÜRKMEN

Dr. Alev Eda OKUTAN SARİBEYLİLER

Dr. Begüm GÖK ÇOBAN

Dr. Erkin ÖZCAN

Dr. Beril ŞAHİNER

Dr. Mustafa Orkun ERTUĞRUL

Dr. Elif ÖZÇELİK



fdi  
FDI World Dental Federation

DORUK

DENTAL Plus+  
IMPLANT KA



TİKA



## 27<sup>th</sup> Congress of the Balkan Stomatological Society (BaSS)

“ Past, Present and Future of Dental Implants and Gerodontology ”

November 9-11, 2023 Istanbul Atlas University / Istanbul, Türkiye



[P-42]

### Facial convexity in macedonian subjects

Biljana Dzipunova<sup>1</sup>, Natasha Tosheska Spasova<sup>1</sup>, Vera Radojkova Nikolovska<sup>1</sup>, Bruno Nikolovski<sup>2</sup>, Matea Dzipunova<sup>2</sup>

<sup>1</sup>Faculty of dentistry, Ss. Cyril and Methodius University, Skopje, North Macedonia

<sup>2</sup>Faculty of medical sciences, Goce Delcev University, Stip, North Macedonia

**Objective:** The assessment of the patients' soft tissue profile is a critical step in orthodontic diagnosis and treatment planning. Achieving a pleasing esthetic profile is an important goal of orthodontic therapy, and can influence the treatment plan and mechanotherapy. The aim of the present study were to evaluate the convexity angles in Macedonian participants with different sagittal irregularities. The examination was performed on 90 profile cephalograms on Macedonian subjects with permanent dentition, aged 16-21 years, divided in Class I, II/1 and III malocclusions, with symmetrical gender distribution and no previous orthodontic treatment. Examined parameters were angle of facial convexity ( $N'-Sn'-Pg'$ ) and angle of total facial convexity ( $N'-Pr'-Pg'$ ). **Results:** The analysis of the size of the angle of facial convexity and angle of total facial convexity showed that the highest average value was in the subjects of class III, followed by class I and the lowest average values were in the subjects with II/1. For  $p < 0.05$ , the analysis of variance indicated a significant difference between the three groups of malocclusion, in both sexes. There is significant linear positive weak correlation between age and the parameter  $N'-Sn'-Pg'$  (with increasing age the value of this parameter also increased). **Conclusion:** Our study has shown that the biological range of facial soft tissues values needs to be determined according to age, sex and orthodontic anomalies for each ethnic group.

**Keywords:** esthetic profile, orthodontic therapy, cephalograms