

Authors: Marija Atanasova Lazareva^{1*}, Katerina Kolevska¹, Maja Chochevska¹
Co-authors: Maja Velichkovska¹, Filip Jolevski¹, Ana Ugrinska¹, Emilija Janevik-Ivanovska²

¹University Institute of Positron Emission Tomography, Bledski dogovor 10, Skopje, Macedonia

²Faculty of Medical Sciences, Goce Delcev University, Stip, North Macedonia

*marija.atanasova@ugd.edu.mk, marija.atanasova6@gmail.com




Background and goal of the present work

Sodium [¹⁸F]fluoride is PET radiopharmaceutical containing fluorine-18 in the form of sodium fluoride, for visualization of the skeletal system and microcalcification.

Aim: Development of an in-house method for [¹⁸F]NaF production using only a dispensing module for both, radiopharmaceutical synthesis and dose dispensing.

Materials and methods

For the production of [¹⁸F]NaF, the modification of the kit was done. On the dispensing kit, the Y connector and anion-exchange cartridge and sterile vial with 0.9% sodium chloride solution were installed. The preconditioned QMA cartridge was connected to the distal part of the Y connector and attached to the first valve of the kit. One proximal ending of the Y connector was connected to the [¹⁸F]F⁻ cyclotron transfer line and the other proximal ending to the vial with 5 mL 0.9% NaCl solution for elution.




Modified kit for production of [¹⁸F]NaF


Results

After development of our in-house method for synthesis, the process optimization was carried out, 10 production batches were performed, with yield higher than 98%, decay-corrected. All obtained QC results confirmed that the quality of [¹⁸F]NaF meets the acceptance criteria defined in the specification.

Test	Method	Acceptance criteria	Results
Pre-release tests			
Appearance	Visual inspection	Clear, colorless solution	Clear colorless solution
Identification	Half-life determination	Radioactivity measurements	1,832 ± 0,014
	Difference in retention time	HPLC	≤ 40 s
Approximate pH value	pH strips	5.5 - 8.0	6.5-7.0
Chemical purity: fluoride (F ⁻)	HPLC	≤ 0.452 mg/mL	≤ 0.452
Radiochemical purity: [¹⁸ F]fluoride	HPLC/gamma detector	min 98.5 % of the total radioactivity	100%
Post-release tests			
Bacterial endotoxins	Chromogenic LAL method	≤ 17.5 IU/mL	< 5.00
Sterility	Test for sterility (Ph. Eur)	Sterile	Sterile
Radionuclidic purity Fluorine-18	Gamma-ray spectrometry	min 99.9 % of the total radioactivity	1,34 x 10 ⁻⁰⁵ ± 7,63 x 10 ⁻⁰⁶



Radiochemical purity



Chemical purity

Conclusion

The in-house method for [¹⁸F]NaF synthesis was successfully designed and developed. The produced [¹⁸F]NaF radiopharmaceutical meets the acceptance criteria defined in the specification, based on Ph. Eur. monograph.