









18th World Congress in Fetal Medicine, Alicante, Spain Abstracts

- Aneuploidies
- cfDNA test
- Defects
- Fetal therapy
- Impaired placentation
- GDM / Obesity
- Preterm birth
- Congenital infection
- Labour / Delivery
- Maternal conditions
- Multiple pregnancies
- Other

Labour / Delivery

- 1. A case of cauda equina syndrome after caesarean section
- 2. A case of conservative surgical management of caesarean scar pregnancy at 14 weeks 📆
- 3. A case of conservative treatment of cesarean scar ectopic pregnancy
- 4. A case of cornual uterine rupture with exit of amniotic sac with umbilical cord and fetal lower limb by left uterine horn 📆
- 5. A case of labor and delivery with deep vein thrombosis
- 6. A case of spontaneous uterine laceration of an unscarred uterus 📆
- 7. Angle of progression and head-to-perineum distance during labor: correlation between midwives and gynecologists 📆
- 8. Cefazolin for neonatal sepsis prophylaxis: analysis of effectiveness and comparative study with penicillin and ampicillin 📆
- 9. Current status of the optimisation of corticosteroids for fetal lung maturity in a tertiary level hospital in Latin America 📆
- 10. Development and validation of a machine learning model for prediction of shoulder dystocia 📆
- 11. Elastography of the uterine cervix: implications for success of induction of labor 🔁
- 12. Elective induction of labour versus spontaneous birth at or after 41 weeks 📆
- 13. Fetal extraction manoeuvers during second stage of labor cesarean delivery
- 14. How should IVF babies be delivered? 📆
- 15. Hypogastric artery ligation in the management of postpartum haemorrhage risk stratification 📆
- 16. Induction of labor for suspected intrauterine growth restriction: extra-amniotic balloon versus prostaglandin E2 📆
- 17. Induction of labor methods in isolated term oligohydramnios 📆