







- Aneuploidies
- cfDNA test
- Defects

C

- Fetal therapy
- Impaired placentation
- GDM / Obesity
- · Preferm birth
- Congenital infection
- Labour / Delivery
- Maternal conditions
- Multiple pregnancies
- Other

Labour / Delivery

- 1. A case of cauda equina syndrome after caesarean section 📆
- 2. A case of conservative surgical management of caesarean scar pregnancy at 14 weeks
- 3. A case of conservative treatment of cesarean scar ectopic pregnancy 🚮
- 4. A case of cornual uterine rupture with exit of amniotic sac with umbilical cord and fetal lower limb by left uterine horn 📆
- A case of labor and delivery with deep vein thrombosis
- 6. A case of spontaneous uterine laceration of an unscarred uterus 51
- 7. Angle of progression and head-to-perineum distance during labor: correlation between midwives and gynecologists 📆
- 8. Cefazolin for neonatal sepsis prophylaxis: analysis of effectiveness and comparative study with penicillin and ampicillin 📆
- 9. Current status of the optimisation of corticosteroids for fetal lung maturity in a tertiary level hospital in Latin America 📆
- 10. Development and validation of a machine learning model for prediction of shoulder dystocia 📆
- 11. Elastography of the uterine cervix: implications for success of induction of labor 📆
- 12. Elective induction of labour versus spontaneous birth at or after 41 weeks 📆
- 13. Fetal extraction manoeuvers during second stage of labor cesarean delivery 📆
- 14. How should IVF babies be delivered? 🐔
- 15. Hypogastric artery ligation in the management of postparturn haemorrhage risk stratification 📆
- 16. Induction of labor for suspected intrauterine growth restriction, extra-amniotic balloon versus prostaglandin E2 📆
- 17. Induction of labor methods in isolated term oligohydramnios
- 18. Intrapartum prediction of birth weight with simplified algorithmic approach derived from maternal characteristics 📆
- 19. Labor induction in twin pregnancies. Does the perintal outcome differ according to chorionicity?
- 20. Levator ani muscle co-activation and active second stage of labor in women undergoing induction of labor 📆
- 21. Obstetric anal sphincter injuries (OASIS): study of frequency after the Parto Adequado program implementation 📆
- 22. Prediction of successful induction of labour in nulliparous at term 📆
- 23. Pregnancy after uterine rupture 📆
- 24. Process of fetal head descent as depicted by ultrasonography How does it compare to the conventional first stage of labor?
- 25. Severe postpartum hemorrhage due to rupture of the uterine artery treated by uterine artery embolisation 📆
- 26. The pink code: obstetric quick team response in obstetrics for safe management of maternal-felal emergencies 📆
- 27. The predictive value of cervical shear wave elastography in the outcome of labour induction 📆
- 28. To insist or not to insist on continuing labour induction after failed cervical ripening? 📆
- 29. To show the outcome of the delivery of patients after previously performed caesarean section 📆
- 30. Ultrasound techniques in predicting a successful vaginal delivery in patients undergoing induction of labour 📆
- 31. Utility of transperineal ultrasound parameters in predicting the mode of delivery during second stage of labour 📆