

ISSN 2671-3497 (Online)

INTERNATIONAL JOURNAL


Institute of Knowledge Management

M E D I S



Vol. 3 Issue 3

MEDICAL SCIENCES & RESEARCH



ISSN 2671-3497 (Online)

**MEDIS – International Journal of Medical
Sciences and Research**

Volume 3, Issue 3, September 2024.

<https://medisij.com>

IMPRESSUM

**International Journal of Medical Sciences and Research
(MEDIS IJMSR)
(Volume 3, Issue 3 September 2024.)**

**Editor in chief:
Prof. d-r Robert Dimitrovski**

**Publisher:
Institute of Management and Knowledge
Address: Gjuro Gjonovikj, 11/4 , Skopje 1000, Macedonia
Phone: +389 70 207 370, + 381 63 700 4281**

**<http://medisij.com>
E-mail: editor@medisij.com**

**For publisher:
Prof. d-r Robert Dimitrovski**

**Print:
GRAFOPROM Bitola**

**Circulation:
50 copies**

CONTENTS

CORRELATION OF HYPOTHALAMIC-PITUITARY-ADRENAL AXIS ACTIVITY WITH CHRONIC COMPLICATIONS IN PATIENTS WITH TYPE 2 DIABETES

Selma Jusufović, Alma Halilčević, Šefkija Balić, Enra Đešević, Alma Horozić, Vedad Herenda.....1-6

THE CRITICAL ROLE OF DETECTION IN EFFECTIVE MELANOMA TREATMENT

Branislav Sančanin, Aleksandra Penjišević.....7-12

COMPARASION OF KINESITHERAPY AND POWERPLATE IN PATIENTS WITH CERVICOARTHROSIS

Tamara Adjiska, Tose Krstev, Lence Nikolovska.....13-17

INFLUENCE OF STRESS COPING BEHAVIOR ON THE EMOTIONAL LOAD OF TEACHERS

Tsvetan Petkov.....19-23

FUNCTIONAL RECOVERY IN PATIENTS WITH ULNAR NERVE INJURY

Danche Vasileva, Tatjana Andonovska.....25-27

MEASURED - REAL VALUES OF INTRAOCULAR TENSION, 24 H AFTER THE APPLICATION OF BROLUCIZUMAB INTRAVITREAL INJECTION IN PATIENTS WITH wAMD

Arsim Hajdari, Nevenka Velickova.....29-32

CHILD TOXOPLASMOSIS AND LIMPHADENOPATHY - CASE REPORT

Valentina Risteska Nejasnikj.....33-36


UTERINE ARTERIOVENOUS MALFORMATIONS – DIAGNOSIS AND TREATMENT IN SERIES OF CASES

Mariana Tsankova, Ivan Kostov, Ivo Petrov.....37-42

FUNCTIONAL RECOVERY IN PATIENTS WITH ULNAR NERVE INJURY

Danche Vasileva^{1*}, Tatjana Andonovska¹

¹Faculty of Medical Sciences, Goce Delcev University, Stip, North Macedonia,
e-mail: dance.vasileva@ugd.edu.mk, tatjana.211546@student.ugd.edu.mk



Abstract: Painful sensations that occur after peripheral nerve injury are a major problem for patients. They occur most often in the acute phase of muscle denervation (denervation pain) or in the early phase of the regeneration process (regenerative pain). The pathophysiology of these painful conditions is complex and results in difficult differentiation of different types of painful conditions that are of crucial importance for the selection of an appropriate therapeutic approach. The place, the degree of the injury, the age of the patient, as well as the delay in starting the treatment significantly affect the outcome of the rehabilitation of injuries of the ulnar nerve. Ulnar nerve rehabilitation in elderly patients, high-level injuries, and delayed cases may result in a poor prognosis. For high-level lesions and lesions that are close to the elbow, the nerve transposition treatment is used. Immobilization in wrist flexion and/or elbow extension with carpal tunnel and release of Guyon's canal are necessary for lesions closer to the wrist. Cubital tunnel syndrome is the second most common compression syndrome of the upper extremities, after carpal tunnel syndrome. The type and severity of nerve injury determines the extent of pathological changes, the capacity for regeneration and the prognosis for recovery. The aim of the study is to shorten the duration of the rehabilitation process and increase the functionality of the affected limb by timely recognizing the symptoms and starting the rehabilitation treatment of the patients, so that the patient would return completely to the activities of everyday life. Material and methods: The study included 10 subjects (5 male subjects and 5 female subjects) in whom the ulnar nerve was diagnosed during the examination. They are treated at the Public Health Institution University Clinic for physical medicine and rehabilitation in Skopje, for a period of 20 working days. The effect of the conducted physical and kinesiotherapy procedures is evaluated. The following were used during the treatment: paraffin therapy, electrical stimulation, Transcutaneous Electrical Nerve Stimulation, galvanic current, diadynamic currents, kinesiotherapeutic exercises (pastoral and actively assisted). Due to the number of studies, it is challenging to make a definitive recommendation before conducting research with a larger number of respondents.

Keywords: ulnar nerve paresis, cubital syndrome

Field: Public Health and Medicine

1. INTRODUCTION

There are several mechanisms of injury to the ulnar nerve: traumatic (fracture, cut) and mechanical, which occurs when n. ulnaris suffers prolonged pressure, usually in the back of the elbow area where it is placed most superficially under the skin. N. ulnaris emerges from the medial bundle of the plexus brachialis (C8 and Th1) and in the upper part of the upper arm lies medial to a. brachialis. Then it goes to the sulcus nervus ulnaris on the outside of the epicondyle of the humerus, and from there it goes to the forearm in the sulcus antebrachii medialis. On the forearm it goes together with the ulnar artery, and with final branches it divides on the medial part of the palm. N. ulnaris is a mixed nerve. Motor innervates m. flexor carpi ulnaris, hypothenar muscles, part of the thenar muscles not innervated by n. medianus, all interosseous muscles and the two medial lumbrical muscles. It sensory innervates the skin of the hypothenar and partially the skin of the fingers.

2. MATERIAL AND METHODS

The study included 10 subjects (5 male subjects and 5 female subjects) who were diagnosed with a rupture of the ulnar nerve during the examination. They are treated at the Public Health Institution University Clinic for physical medicine and rehabilitation in Skopje, for a period of 20 working days.

- Method of kinesiotherapy:

During the treatment, physical procedures (heat procedures and electrical procedures) and kinesiotherapeutic exercises were used in order to reduce pain, improve trophism and allow a faster return of the complete function of the affected limb. During the dosing of all the above-mentioned procedures, attention was paid to the patient's condition in terms of fatigue and load. The following methods were used during the treatment: paraffin therapy, electrical stimulation, TENS, galvanic current, diadynamic currents, kinesiotherapeutic exercises (pastoral and actively assisted).

- Examination methods:

*Corresponding author: dance.vasileva@ugd.edu.mk



In the course of the research, functional diagnostic methods were applied, summarizing the results obtained at the beginning of the treatment, on the tenth day and on the twentieth day of the treatment. The results and condition of the muscles (m. flexor carpi ulnaris and m. oponents digiti minimi) are registered and evaluated with the MMT and the mobility of the wrist with a goniometer.

Tab.1. Tabular presentation of subjects with injury of n. ulnaris

Group of respondents	Age	Duration of the problem
10 patients (5 men and 5 women)	36,5 ± 9,4	From 2 months to 6 months

Source: Authors' own research Vasileva, D. and Andonovska, T.

- Statistical methods:

For the purposes of the research, a program for qualitative data processing was used. They are processed with variation (Student-Fisher t-test) and alternative analysis, which summarize the changes during the therapy.

3. RESULTS

After finishing the treatment and summarizing the obtained results, a comparison was made of the achieved changes after two ie. four weeks. The obtained data are shown in tab.2, tab.3, figure 1. and figure 2. through which a significant improvement in the condition of the majority of patients was observed.

Tab.2. Tabular data display of changes in m.flexor carpi ulnaris and m. oponents digiti minimi

Musculus	Day 1, X ±SD	Day 10, X ±SD	Day 20, X ±SD
m. flexor carpi ulnaris	1.4 ± 0.62 **	2.27 ± 0.77 **	3.84 ± 0.93 **
m. oponents digiti minimi	0.63 ± 0.54 **	1.83 ± 0.67 **	2.82 ± 0.73 **

** p<0,05 – significant changes in movements in relation to the beginning and end of the treatments

Source: Authors' own research Vasileva, D. and Andonovska, T.


Tab.3. Tabular data display of changes in movements of the wrist

ROM	Day 1, X ±SD	Day 20, X ±SD
Dorsiflexion	42 ± 12.48 **	59 ± 10.44 **
Palmar flexion	48 ± 10.53 **	65.5 ± 12.93 **
Radial deviation	10 ± 3.87 **	17.5 ± 4.6 **
Ulnar deviation	1.5 ± 5.5 **	13.5 ± 6.72 **

** p<0,05 – significant changes in movements in relation to the beginning and end of the treatments


Source: Authors' own research Vasileva, D. and Andonovska, T.

Figure 1. Results obtained by manual muscle test of m. flexor carpi ulnaris and m. oponents digiti minimi



Source: Authors' own research Vasileva, D. and Andonovska, T.

Figure2. Changes obtained by measuring movements of the wrist



Source: Authors' own research Vasileva, D. and Andonovska, T.

4. DISCUSSION AND CONCLUSION

Ulnar nerve impingement is a common occurrence in forearm bone injuries. Patients face difficulties in performing daily life activities and are dysfunctional in society. With timely recognition of the symptoms and starting the rehabilitation treatment of the patients, the duration of the rehabilitation process is shortened and the functionality of the affected limb is increased.

REFERENCES

- Атанасовска, С. (2024) Физиотерапија кај пациенти со повреда на нервус улнарис (кубитален синдром). Физиотерапија Македоница, 5, 20-22 <https://zpft.org/spisanie/5.pdf>
- Василева, Д. (2020). Кинезиологија практикум. Универзитет „Гоце Делчев“ - Штип, ISBN 978-608-244-741-4
- Василева, Д. (2022). Основи на кинезитерапија Универзитет „Гоце Делчев“ - Штип
- Василева, Д. (2023). Кинезиологија. Универзитет „Гоце Делчев“, Факултет за медицински науки, Штип. ISBN 978-608-244-951-7
- Димитрова, Е. (2003). Мобилизација на нервната система - НСА Прес.
- Крстев, Т (2020) Проприоцептивно нервно-мускулно олеснување - практикум.
- Крстев, Т (2020) Проприоцептивно нервно-мускулно олеснување - скрипта. ISBN 978-608-244-751-3.
- Крстев, Т. (2022). Функционална дијагноза на локомоторен апарат „Гоце Делчев“ – Штип
- Крстев, Т., Николовска, Л., Василева, Д., Страторска, Т. (2020) Медицинска масажа. ISBN 978-608-244-753-7.
- Abaskhron, M., Ezzat, M., Boulis, A., Safoury, Y. (2024) Supercharged end-to-side anterior interosseous nerve transfer to restore intrinsic function in high ulnar nerve injury: a prospective cohort study. *BMC Musculoskelet Disord.* 25(1):566. doi: 10.1186/s12891-024-07650-4.
- Agarwal, A., Chandra, A., Jaipal, U., et al. (2019) Imaging in the diagnosis of ulnar nerve pathologies—a neoteric approach. *Insights Imaging.* 10(1),1–12.
- Bašcarević, V., Samardžić, M., Ivanović, S., Mičović, M., Cvrkota, I. (2004). Terapijski pristup bolnim sindromima posle povreda perifernih nerava—Institut za neurohirurgija KCS, Beograd. 51(4), 65-68.
- Bordes, S., Jenkins, S., Bang, K., et al. (2021) Ulnar nerve subluxation and dislocation: a review of the literature. *Neurosurg Rev.* 44(2),793–798.
- Chaudhary, R., Karkala, N., Nepal, P., Gupta, E., Kaur, N., Batchala, P., Sapire, J., Alam, S. (2024) Multimodality imaging review of ulnar nerve pathologies. *Neuroradiol J.* 37(2):137-151. doi: 10.1177/19714009231166087. Epub 2023 Mar 24. PMID: 36961518; PMCID: PMC10973834.
- Natroshevili, T., van de Warenburg, M., Heine, E., Slater, N., Walbeehm, E., Bartels, R. (2023) Conservative Treatment of Ulnar Nerve Compression at the Elbow: A Systematic Review and Meta-Analysis. *Arch. Plast. Surg.* 50:70–81. doi: 10.1055/s-0042-1757571
- Nyman, E., Dahlin, L. (2024) The Unpredictable Ulnar Nerve-Ulnar Nerve Entrapment from Anatomical, Pathophysiological, and Biopsychosocial Aspects. *Diagnostics (Basel).* 14(5),489. doi: 10.3390/diagnostics14050489
- Power, H., Sharma, K., El-Haj, M., Moore, A., Patterson, M., Mackinnon, S. (2019) Compound Muscle Action Potential Amplitude Predicts the Severity of Cubital Tunnel Syndrome. *J Bone Joint Surg Am.* 101(8),730-738. doi: 10.2106/JBJS.18.00554.
- Vij, N., Traube, B., Bisht, R., Singleton, I., Cornett, E., Kaye, A., Imani, F., Mohammadian, Erdi A., Varrassi, G., Viswanath, O., et al. (2020) An Update on Treatment Modalities for Ulnar Nerve Entrapment: A Literature Review. *Anesth. Pain Med.* 10:e112070. doi: 10.5812/aapm.112070.
- Zhang, D., Earp, B., Homer, S., Blazar, P. (2023) Factors Associated With Severity of Cubital Tunnel Syndrome at Presentation. *Hand (N Y).* 18(3):401-406. doi: 10.1177/15589447211058821.

