



PLANT DERIVED ESSENTIAL OILS AS ECO-FRIENDLY ALTERNATIVES OF SYNTHETIC PESTICIDES

Biljana Kovacevik, Sasa Mitrev, Emilija Arsov, Natalija Markova Ruzdik
Faculty of Agriculture, Goce Delcev University, Stip, North Macedonia

biljana.kovacevik@ugd.edu.mk




Introduction

- According to the European Pharmacopoeia, essential oils (EOs) are complex mixtures of volatile, aromatic, lipophilic compounds obtained from plant material through methods such as steam or water distillation, dry distillation, or mechanical processes, without the application of heat.
- These compounds are predominantly localized in the leaves and flowers of higher plants and are primarily biosynthesized via enzymatic mevalonate (MVA) and methylerythritol phosphate (MEP) pathways.



Distillation of Essential oils



ORGANIC
serum



Introduction

- The majority of studies emphasize the insecticidal properties of lemongrass, citronella, clove, and eucalyptus EOs.
- Additional investigations have identified EOs with herbicidal, acaricidal, nematocidal, antibacterial, and antifungal activities.
- However, only a limited number have been approved as commercial biopesticides, underscoring the need for further research into their mechanisms of action, regulatory compliance, and integration into the pest management systems.





Introduction

Some EOs based commercial biopesticides for plant protection

EO source	Commercial name	Producer	Effective against
<i>Allium sativum</i>	GC-3	JH Biotech, Inc., California	Powdery Mildew
Clove oil Garlic oil	GC-Mite	JH Biotech, Inc., California	various mites and insects
Clove oil	BIOXEDA, Mat-ratec, WeedZap	Xeda International Brandt, JH Biotech, Inc.	Fungicide, bioherbicide
Thyme oil	Tymox	Laboratoire M2	powdery mildew, gray mold and fire blight
<i>Rosmarinus officinalis</i>	EcoTrol Plus (10%) Hexacide (5%) Sporan (17.6%)	EcoSMART Technologies Inc.	Mites, aphids, beetles, thrips, whiteflies, caterpillars, mealybugs
Orange oil	GreenMatch	Pomerix	Aphids, whiteflies, soft-bodied insects
<i>Neem essential oil combined with Beauveria bassiana (a fungal biopesticide)</i>	Naturalis-L	AgriGrem Ltd.	Whiteflies, aphids, and thrips
<i>Beauveria bassiana and rosemary EO</i>	BotaniGard MAXX	Certis Biologicals	aphids, thrips, and spider mites
<i>Rosemary, geraniol and peppermint EOs</i>	Ecotec	Brandt Consolidated, Inc.	Mites, thrips, and aphids



- The commercialization of EO-based biopesticides began in the 1990s with the introduction of EcoSMART, a formulation containing oils such as peppermint, clove, rosemary, and thyme, demonstrating efficacy against a range of phytopathogens and insect pests.



Chemical composition of EOs


- EOs are complex mixtures of VOCs, lacking fatty substances typically found in oils.
- Their chemical composition is influenced by factors such as plant physiology, climate, soil conditions, plant health, harvest timing, and extraction methods.
- The most abundant and bioactive compounds: terpenes, terpenoids, and phenylpropanoids.
- Terpene synthesis occurs in plastids via the MEP pathway, with isopentenyl diphosphate and dimethylallyl diphosphate as precursors.

EO compounds	Chemical class	Source/Essential Oil	Effectiveness
<i>Azadirachtin</i>	Tetranortriterpenoid	Neem	Broad-spectrum insecticidal and antifungal; disrupts feeding, molting, reproduction
<i>Citronellal</i>	Monoterpenoid aldehyde	Citronella, geranium, lemongrass	Repellent/insecticidal (mosquitoes, aphids, mealybugs, thrips); nematocidal and fungicidal activity
<i>Geraniol</i>	Monoterpenoid alcohol	Citronella, geranium, lemongrass	
<i>Carvacrol</i>	Monoterpenoid phenol	Oregano, thyme	Potent antifungal, anti- <i>Rhizoctonia</i> activity, insect/acarid toxicity;
<i>Camphor</i>	Monoterpenoid ketone	Eucalyptus, rosemary	Antifungal, insecticidal, acaricidal; inhibits nematode egg hatching
<i>1,8-Cineole (Eucalyptol)</i>	Monoterpenoid ether	Eucalyptus, rosemary	
<i>Eugenol</i>	Phenylpropanoid	Clove, cinnamon	Antifungal (e.g. <i>Colletotrichum</i> , <i>Fusarium</i>), insecticidal, acaricidal
<i>Trans-Anethole</i>	Phenylpropanoid	Fennel, anise, star anise	Insecticidal and nematocidal activity
<i>Linalool</i>	Monoterpenoid alcohol	Lavender, basil, coriander	insecticidal/fumigant (stored-grain pests, thrips)
<i>Thymol</i>	Monoterpenoid phenol	Thyme, oregano, savory	Strong antifungal (<i>Fusarium</i> , <i>Verticillium</i> , <i>Phytophthora</i>) and mitocidal activity



Antimicrobial activity

- EOs exhibit both single and multi-target bioactivity against microorganisms, with their efficacy largely attributed to their chemical composition.
- While major EO compounds are often studied, bioactivity is typically enhanced by the unique chemical profiles of EOs, which depend on the plant species, plant part, and extraction method.
- Lipophilic nature allows them to penetrate microorganism cell walls, increasing permeability, and contributing to higher bioactivity, especially against gram-positive bacteria with an outer peptidoglycan layer.
- For example, *Psoralea glandulosa* EO was found effective against *Clavibacter michiganensis*, a seed-borne pathogen.



Antifungal activity of *Calamintha nepeta* EO against *Penicillium digitatum* (a) and *Botrytis cinerea* (b) (Ambrocio et al., 2019)




***Calamintha nepeta* (Lesser calamint)**




Antimicrobial activity

- Terpene alcohols with varying carbon chain lengths, such as **farnesol** and **nerolidol**, have shown strong antibacterial properties.
- Monoterpenes, commonly found in **thyme** and **oregano** EOs, are particularly effective against plant pathogens by disrupting cell membranes and interfering with metabolic processes.
- **Thymol**, for instance, inhibits *Botrytis cinerea*, *Fusarium oxysporum*, and *Alternaria alternata*, while **carvacrol** targets *Rhizoctonia solani*.
- Other monoterpenes like **linalool** and **menthol** demonstrate efficacy against root rot pathogens and postharvest molds.
- Additionally, compounds like **eucalyptol** (1,8-cineole) from *Eucalyptus globulus* show effectiveness against *Verticillium dahliae* and *Xanthomonas* spp.
- Terpenes such as **α -pinene** and **β -pinene**, found in conifer oils, also display strong antimicrobial activity against fungi like *Fusarium* spp. and *Botrytis cinerea*.



Morphological abnormalities in larvae after treatment with EO (Khater, 2013)



(A) Normal *C. pipiens* larva (B) Abnormal larvae produced after treatment with oil of *S. terebinthifolius* unripe fruits at 100 mg/L, showing a destroyed digestive system, especially midgut (arrows).



Limitations and challenges

Pros and Cons	
Availability of adequate plant biomass	Valuable in integrated pest management
Composition can vary between batches or brands	Valuable in organic agriculture
Lower Efficacy than synthetic pesticides	Low mammalian toxicity
Frequent reapplication	Target-Specific
Often more expensive	Low environmental persistence

- Despite increasing evidence of the pesticidal and repellent efficacy of essential oils (EOs), commercial products remain scarce due to several challenges.
- Key limitations include the availability of adequate plant biomass, variability in EO chemical composition, difficulties in formulation standardization, and regulatory barriers.
- While some EOs are produced at scale for aromatherapy or food applications, those from rare or difficult-to-cultivate species face supply constraints, limiting their commercial viability.
- Chemical variability within species necessitates strict quality control and standardization to maintain consistent bioactivity.



Seed priming tests in lab

Limitations and challenges

- Advanced formulations could also mitigate volatility and enhance chemical stability, optimizing EOs for practical use.
- Recent studies have explored the indirect effects of EOs, such as priming, which enhances plant defense mechanisms.
- Research on EO-induced priming, particularly in response to fungal pathogens, has shown promising results.
- Notably, seed coating with EOs has demonstrated priming effects in *Solanum lycopersicum* seedlings against *Fusarium oxysporum*, triggering metabolic and epigenetic changes with minimal EO usage and long-lasting effects.





Conclusion

- Essential oils (EOs) have demonstrated efficacy against a range of plant pests and pathogens, including fungi, insects, and nematodes.
- Their synergistic effects lower the risk of resistance development, positioning them as a promising biopesticides.
- However, challenges such as biomass availability, chemical instability, formulation difficulties, and potential phytotoxicity hinder their widespread use.
- Addressing these issues could be facilitated through plant domestication and cultivation, which may improve chemical stability (chemotype consistency) and biomass production.
- These findings underscore the potential of EOs in biopesticide development, but further research is necessary to elucidate the influence of EO composition on plant responses to biotic stress.
- This knowledge will be critical for optimizing EO applications and maximizing their effectiveness in agriculture.



**THANK YOU
FOR
YOUR ATTENTION**

