

Volume 10, Number 2, July 2026

AJTES

ALBANIAN JOURNAL OF TRAUMA AND EMERGENCY SURGERY

Focus on
“Trauma and Emergency Surgery”
and not only...

Official publication of

ISSN: 2521-8778 (print version) ISSN:
2616-4922 (electronic version)



INDEXING

The Albanian Journal of Trauma and Emergency Surgery (AJTES) is a peer-reviewed, open-access journal committed to high editorial standards, transparency, and international best practices in scholarly publishing.

AJTES is currently indexed and registered in the following recognized scholarly infrastructures:

DOAJ DIRECTORY OF
OPEN ACCESS
JOURNALS

Directory of Open Access Journals (DOAJ)

<https://doaj.org/toc/2616-4922>



Digital Object Identifier (DOI) via Crossref

<https://doi.org/10.32391/ajtes.v10>

ISSN INTERNATIONAL
STANDARD
SERIAL
NUMBER
INTERNATIONAL CENTRE

ISSN International Centre

Print ISSN: 2521-8778

Online ISSN: 2616-4922

<https://portal.issn.org/resource/ISSN/2616-4922>

VISIBILITY & SCHOLARLY DISSEMINATION

Articles published in AJTES are discoverable through major academic search engines, bibliographic platforms, and library networks, which enhance international visibility and citation potential:

- Google Scholar
- Semantic Scholar
- Scilit
- WorldCat
- ResearchGate
- Mendeley
- ROAD (Directory of Open Access Scholarly Resources)
- Global Open Access Portal (GOAP)
- ORCID
- Publons
- COBISS.AL
- Joint Information Systems Committee (JISC)
- Developing Library Network (DELNET)
- ...

EDITORIAL & ETHICAL STANDARDS

AJTES follows the recommendations of the International Committee of Medical Journal Editors (ICMJE) and complies with international standards for editorial independence, peer review integrity, publication ethics, and open-access transparency.

OPEN ACCESS & LICENSING

All articles published in AJTES are made immediately and permanently available under the Creative Commons Attribution–NonCommercial (CC BY-NC) License.



This license permits non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.





Volume 10, Number 2, July 2026

Focus on “Trauma and Emergency Surgery” and not only...

Official publication of





Albanian Journal of Trauma and Emergency Surgery - AJTES
Official Publication of the
Albanian Society for Trauma and Emergency Surgery - ASTES

CHAIRMAN OF THE EDITORIAL BOARD

Prof. Kastriot HAXHIREXHA, MD, Ph.D.
State University of Tetova, Republic of North MACEDONIA

EDITOR IN CHIEF

Prof. Agron DOGJANI, MD, Ph.D., FACS, FISS, FICS
University of Medicine of Tirana, ALBANIA

DEPUTY EDITOR

Asc. Prof. Alfred IBRAHIMI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Altin VESHTI, MD, Ph.D.
University Hospital Centre "Mother Teresa", Tirana, ALBANIA

Asc. Prof. Basri LENJANI MD, Ph.D.
University Clinical Center of Kosovo, Pristina, KOSOVO

Asc. Prof. Edvin SELMANI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Francesk MULITA, MD, PhD
Department of Surgery, Medical School, Democritus University
of Thrace, Alexandroupolis, GREECE

INTERNATIONAL EDITORIAL BOARD

Prof. Juan Carlos PUYANA, MD, FRCSC, FACS, FACCP
Director of Global Health - Surgery, University of Pittsburgh; USA;
Royal College of Surgeons in IRELAND

Prof. Fausto CATENA, MD, Ph.D., FACS.
Department of Emergency and General Surgery, Parma University
Hospital, ITALY

Prof. Ada IŞIK, MD, FTBS, FEBS, FACS, FSIS
Istanbul Medeniyet University, Istanbul, TURKEY

Prof. Dietrich DOLL, MD, Ph.D., FACS.
St. Marien Hospital Vechta, GERMANY

Prof. Chrysanthos GEORGIU, MD, Ph.D., FACS, F.E.B.S
Professor of Surgery, University of Nicosia, CYPRUS

Prof. Peep TALVING, MD., Ph.D., FACS
Faculty of Medicine, University of Tartu, ESTONIA

Prof. Bellal A. JOSEPH MD, FACS
The University of Arizona, College of Medicine - Tucson, USA

Joakim JORGENSEN, MD, Ph.D., FACS.
Oslo University Hospital, Oslo, NORWAY

Prof. Hayato KURIHARA, MD, FEBS, FACS.
Emergency Surgery Unit Fondazione IRCCS Ca' Granda Ospedale
Maggiore Policlinico, Milan, ITALY

Prof. Pantelis VASSILIU, MD, Ph.D., FACS.
National and Kapodistrian University of Athens, GREECE

Ass. Prof. Faton T. HOXHA, MD, Ph.D.
University Clinical Center of Pristina, Republic of KOSOVO

Prof. Boris KESSEL MD,
Professor of Surgery, Rappaport Faculty of Medicine, Haifa, ISRAEL

Prof. Mauro ZAGO, MD, FEBS, FACS,
A. Manzoni Hospital, ASST Lecco, ITALY

Prof. Boris SAKAKUSHEV, MD, Ph.D.
"Saint George" University Hospital, Plovdiv, BULGARIA

Asc. Prof. Bogdan DIACONESKU MD, Ph.D.
Carol Davila University of Medicine, Bucharest, ROMANIA

Ass. Prof. Ilir HASANI MD, Ph.D.
PHI UC TOARICEC Skopje, Republic of North MACEDONIA

NATIONAL EDITORIAL BOARD

Prof. Xheladin DRAÇINI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Arben GJATA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Arben BEQIRI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Arvin DIBRA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Rudin DOMI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Sokol XHEPA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Artur XHUMARI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Ridvan ALIMEHMETI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Arsen SEFERI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Monika FIDA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Prof. Myzafer KAÇI, MD, Ph.D.
The University of Medicine of Tirana, ALBANIA

Asc. Prof. Zamira SHABANI, MD, Ph.D.
The University "Luigj Gurakuqi", Shkodra, ALBANIA

Asc. Prof. Saimir KUÇI, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Asc. Prof. Leart BERDICA, MD, Ph.D.
University of Medicine of Tirana, ALBANIA

Shkëlzen OSMANAJ, MD, Ph.D.
University Hospital of Trauma, Tirana, ALBANIA

Zamir DEMIRAJ, MD, Ph.D.
University Hospital of Trauma, Tirana, ALBANIA

ADVISORY EXPERT

Hysni BENDO, MD.
SRH Clinic Karlsbad, Karlsruhe GERMANY

Amarildo BLOSHMI, MD
The University of Medicine of Tirana, ALBANIA.

Kledia POLLO, MD, Ph.D.c
The University Hospital of Trauma, Tirana, ALBANIA.

ASSISTANT EDITORS

Besmira ZAMA, MD, Ph.D.
The University of Medicine of Tirana, ALBANIA.

Sabina DOGJANI, MD
The University of Medicine of Tirana, ALBANIA.

Erjona ZOGAJ MSN, Ph.Dc
The University Hospital of Trauma, Tirana, ALBANIA.

Besa ISUFAJ, MSN
The University of Medicine of Tirana, ALBANIA.

Web developer: Klevis DOÇI MD - The University of Medicine of Tirana, ALBANIA.

E-mail: ajtesjournal@gmail.com ; ajtes.editor@gmail.com

Website: <https://journal.astes.org.al>

Mobile: +355 69 346 5944 | +355 69 205 6123

Address: Street "Lord Bajron" No.40, PC 1000 AL, Tirana, Albania

Design & Layout: Bujar KAROSHI **Printed:** M&B BOTIME & Graphic Studio

The **Albanian Journal of Trauma and Emergency Surgery (AJTES)** is the official scientific publication of the **Albanian Society for Trauma and Emergency Surgery (ASTES)**.

Structure and format

AJTES is an international, peer-reviewed, open-access journal that publishes a broad range of scholarly contributions, including: Letters to the Editor, Editorials, Perspectives, Systematic Reviews and Narrative Reviews, original articles, Case-reports, Short reports, Clinical Practice Guidelines, Health Policy and Management Articles, Book Reviews, Ethics and Medical Humanities, Symposium and Mini-Symposium Papers, health care policy and management, and practice guides. The content and structure of each issue are determined by the Editor-in-Chief in consultation with the Editorial Board and reflect the journal's editorial policy and scientific priorities.

Editorial expertise

The Editorial Board comprises internationally recognized experts representing diverse disciplines within trauma, emergency surgery, and related medical sciences.

All submitted manuscripts undergo an independent peer-review process conducted by reviewers selected according to their scientific expertise and experience in the relevant subject area.

Important notice for authors

All manuscripts must be submitted electronically through the AJTES Online Submission System:

<https://journal.astes.org.al>

Authors are advised to carefully consult the **Instructions for Authors** before submission.

Submitted manuscripts must be original, must not have been published previously, and must not be under consideration for publication elsewhere.

Following publication, copyright and licensing conditions are governed by the journals official Copyright and Open Access Policy.

No part of any published article may be reproduced, stored in a retrieval system, transmitted, translated, or distributed in any form or by any means without prior written permission from the copyright holder, except as permitted under the applicable Creative Commons license.

Permissions

Requests to reproduce, reuse, translate, or distribute material published in AJTES should be addressed to the Editorial Office.

Advertising policies

AJTES accepts both display and classified advertising that is consistent with the journal's mission and ethical standards.

The Editorial Office reserves the right to reject any advertisement deemed inappropriate or inconsistent with the journal's advertising policies.

Publication of advertisements or commercial material does not imply endorsement or approval by AJTES, the Editorial Board, ASTES, or the Publisher of any product, service, manufacturer, or claim.

Disclaimer

The opinions and conclusions expressed in published articles are solely those of the authors and do not necessarily reflect the views of the Editorial Board, the Albanian Society for Trauma and Emergency Surgery (ASTES), or the Publisher.

Publication of an article does not constitute endorsement of its content by the Journal, its Editorial Board, ASTES, or the Publisher.

Volume 10
Number 2
July 2026

LETTER TO EDITOR

2056 Advancing Pilonidal Sinus Care Through Structured International Collaboration: The PiloNERDs Project

Dietrich Doll

DOI: <https://doi.org/10.32391/ajtes.v10i2.525>

ORIGINAL ARTICLES

2058 An Assessment of the Quality of YouTube Videos on the Treatment of Nosebleeds

Resmiye Nur Okudan, Cihan Bedel, Fatih Selvi

DOI: <https://doi.org/10.32391/ajtes.v10i2.540>

2063 The Foreign Body Aspiration in Children in Albania during 2006-2025

Melpomeni Bizhga, Spiro Sila, Anita Pollozhani, Klevis Thomai

DOI: <https://doi.org/10.32391/ajtes.v10i2.539>

2067 Optimizing Varicocele Treatment: A Comparative Review of Surgical and Minimally Invasive Approaches

Erzen Elezi, Florin Deari, Vjosa Elezi, Aferdita Ademi

DOI: <https://doi.org/10.32391/ajtes.v10i2.461>

2075 Non-Conventional Risk Factors in Acute Coronary Syndrome

Taulant Gishto, Edlira Rruci, Naltin Shuka, Silvia Methoxha

DOI: <https://doi.org/10.32391/ajtes.v10i2.523>

2080 Gestational Age and Pregnancy Outcomes in SARS-CoV-2 - Positive Women: A Retrospective Cohort Study with External Comparison

Vera Beca, Mirela Rista, Daniela Nakuçi, Eliona Demaliaj

DOI: <https://doi.org/10.32391/ajtes.v10i2.535>

2084 Beyond Stone Removal: Modern Challenges and Future Directions in the Management of Urinary Tract Calculi

Florin Deari, Erzen Elezi, Hana Deari, Aferdita Ademi,

DOI: <https://doi.org/10.32391/ajtes.v10i2.421>

2093 Anatomical Variations of the Vertebral Arteries, and their Impact on Cerebral Circulation

Gazmend Zylbeari, Zamira Bexheti, Koco Cakalaroski, Sadi Bexheti, Tose Krstev

DOI: <https://doi.org/10.32391/ajtes.v10i2.545>

2097 Management of Acute Myocardial Infarction in the Emergency Department of Elbasan Hospital

Ergys Mici, Evisa Mici

DOI: <https://doi.org/10.32391/ajtes.v10i2.524>

ORIGINAL ARTICLES

Anatomical Variations of the Vertebral Arteries, and their Impact on Cerebral Circulation

Gazmend Zylbeari^{1,2*}, Zamira Bexheti³, Koco Cakalaroski⁴, Sadi Bexheti¹, Tose Krstev²

¹Faculty of Medical Sciences, University of Tetovo, North Macedonia

²Faculty of Medical Sciences, Goce Delcev University, Stip, North Macedonia

³South East European University-Tetovo, North Macedonia

⁴Faculty of Medical Sciences, Ss. Cyril and Methodius University in Skopje, North Macedonia

Corresponding author: Gazmend Zylbeari, MD

Faculty of Medical Sciences, University of Tetovo, Republic of North Macedonia

Email: gazmend.zylbeari7@gmail.com

Received: 04 May 2026 • Accepted: 20 June 2026 • Published online: 20 July 2026

ABSTRACT

Background: The vertebral arteries are integral to the vertebrobasilar system, contributing approximately 20% of cerebral blood flow and ensuring perfusion of the posterior brain and spinal cord. Anatomical variations such as hypoplasia and dolichoectasia may compromise posterior circulation hemodynamics and influence cerebrovascular risk.

Objective: To investigate the prevalence, diameter, and gender distribution of vertebral artery variations and assess their potential clinical implications for posterior cerebral circulation.

Materials and Methods: A retrospective observational study was conducted on 60 patients (25 females, 41.7%; 35 males, 58.3%; mean age 45.0 ± 12.0 years). Radiological evaluation included Magnetic Resonance Angiography, Doppler ultrasonography, and Computed Tomography. Vertebral artery morphology was classified as normal, hypoplastic, or dolichoectatic.

Results: Hypoplasia was the most frequent variation. In females, hypoplasia was observed in 28.0% of left and 40.0% of right vertebral arteries. In males, it was present in 20.0% of left and 22.9% of right arteries. Dolichoectasia was more prevalent in males. These findings highlight gender-related differences and underscore the importance of vascular imaging in patients with posterior circulation symptoms.

Conclusion: Vertebral artery variations are relatively common and clinically significant. Recognition of these variations is essential for accurate diagnosis, risk stratification, and management of cerebrovascular disorders involving posterior circulation.

Keywords: Vertebral artery; anatomical variations; hypoplasia; dolichoectasia; posterior cerebral circulation

How to cite this article:

Zylbeari, G., Bexheti, Z., Cakalaroski, K., Bexheti, S., & Krstev, T. (2026). Anatomical variations of the vertebral arteries and their impact on cerebral circulation. *Albanian Journal of Trauma and Emergency Surgery*, 10(2), 2093-2096. <https://doi.org/10.32391/ajtes.v10i2.545>

OPEN ACCESS

This article is published with Open Access at journal.astes.org.al.

© The Author(s) 2026.

The Albanian Journal of Trauma and Emergency Surgery (AJTES) is an Open Access journal. All articles are distributed under the terms of the Creative Commons Attribution–NonCommercial License (CC BY-NC), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

License: <https://creativecommons.org/licenses/by-nc/4.0/>

INTRODUCTION

Anatomical variations of the vertebrobasilar system are relatively common, and many of them are considered benign. Numerous studies have reported unilateral hypoplasia of the vertebral arteries, variations in the posterior and anterior inferior cerebellar arteries, as well as differences in the origin and branching patterns of the vertebral and basilar arteries.

These variations are believed to arise due to developmental errors during embryogenesis (1,2). Hemodynamic assessment of the vertebrobasilar system (VBS) plays an important role in identifying factors that influence cerebral circulation abnormalities and anatomical-physiological deviations. Such evaluations may help predict potential sites of pathological changes, enable early detection of cerebrovascular disease, and contribute to the assessment of treatment options and therapeutic efficacy (3,4).

Clinically, the vertebral arteries are divided into four segments (V1–V4), each with a distinct anatomical course from their origin in the subclavian artery to their formation of the basilar artery at the vertebrobasilar junction (5). Variations in this system are mainly related to vessel origin, diameter asymmetry, and branching patterns (6,7).

The most frequently reported anatomical variations include vertebral artery asymmetry, hypoplasia, fenestration of the basilar artery, aplasia of a vertebral artery, trifurcation of the basilar artery, and agenesis of the posterior cerebral arteries (8–10). These anomalies may have significant clinical implications, including an increased risk of ischemia, posterior circulation infarctions, aneurysm formation, and other cerebrovascular disorders (11,12).

Vertebrobasilar insufficiency (VBI) represents a clinically important condition characterized by reduced blood flow to the posterior part of the brain. It may result from atherosclerotic changes, vascular occlusion, or embolic events, and can lead to transient or permanent neurological deficits (13,14).

The basilar artery, as a key vessel supplying the brainstem and posterior cerebral hemispheres, plays a crucial role in maintaining adequate cerebral perfusion through its contribution to the Circle of Willis (CW) (15,16).

Given the potential clinical significance of these anatomical variations, their early identification is essential for understanding cerebrovascular risk and improving diagnostic and therapeutic strategies.

Purpose of the work:

The purpose of this study is to evaluate the frequency and diameter of anatomical variations of the vertebral arteries (left and right vertebral artery), analyze their distribution according to gender, and assess their potential clinical significance in posterior cerebral circulation and vertebrobasilar hemodynamics.

MATERIALS AND METHODS

This study was designed as a retrospective observational analysis. A total of 60 patients were included, comprising 25 females (41.67%) and 35 males (58.33%), with a mean age of 45.0 ± 12.0 years. The patients were examined over a one-year period in the Departments of Neurosurgery and Traumatology at Klinička Bolnica Tetovo.

Clinical indications for imaging included head trauma, dizziness, cervical complaints, neck trauma, and other neurological symptoms. Radiological evaluation was performed using Magnetic Resonance Angiography (MRA), color Doppler ultrasonography, and Computed Tomography (CT). The diameters of the left and right vertebral arteries were systematically analyzed.

Vertebral artery hypoplasia was defined as a vessel diameter of less than 2.0 mm, while dolichoectasia was defined as a diameter greater than 3.7–4.0 mm, in accordance with previously established criteria in the literature (15,16).

Due to the absence of a healthy control group, the obtained results were interpreted and compared with data reported in the international literature.

Statistical processing of the material examined

Statistical analysis was performed using descriptive statistical methods. Categorical variables were presented as absolute numbers (n) and percentages (%), while continuous variables were expressed as arithmetic mean (mean) and standard deviation (SD). Due to the absence of a control group, comparisons with international studies were carried out using a descriptive approach.

RESULTS

In the female group (n = 25), the left vertebral artery showed normal morphology in 17 cases (68.0%), hypoplasia in 7 cases (28.0%), and dolichoectasia in 1 case (4.0%). The right vertebral artery demonstrated normal morphology in 14 cases (56.0%), hypoplasia in 10 cases (40.0%), and dolichoectasia in 1 case (4.0%).

In the male group (n = 35), the left vertebral artery was normal in 25 cases (71.4%), while hypoplasia and dolichoectasia were observed in 7 (20.0%) and 3 (8.6%) cases, respectively.

The right vertebral artery showed normal morphology in 23 cases (65.7%), hypoplasia in 8 cases (22.9%), and dolichoectasia in 4 cases (11.4%).

Overall, hypoplasia was the most frequent anatomical variation, with a higher prevalence observed in females, particularly affecting the right vertebral artery.

Table 1. Left and right vertebral artery distribution, female**Female (n = 25)**

Vertebral artery	Normal morphology	hypoplasia	dolichoectasia
Left	17 (68.0%)	7 (28.0%)	1 (4.0%)
Right	14 (56.0%)	10 (40.0%)	1 (4.0%)

Table 2. Left and right vertebral artery distribution, male**Male (n=35)**

Vertebral artery	Normal morphology	hypoplasia	dolichoectasia
Left	25 (71.4%)	7 (20.0%)	3 (8.6%)
Right	23 (65.7%)	8 (22.9%)	4 (11.4%)

DISCUSSION

In our study, the prevalence of vertebral artery hypoplasia was notably higher compared to most international reports. *Valenzuela-Fuenzalida et al.* (2024) (15) reported an average prevalence of approximately 11% in the general population. Similarly, *Park et al.* (2007) (16) demonstrated that vertebral artery hypoplasia is associated with ischemic stroke territory involvement and is more frequently observed in patients with posterior circulation ischemia.

In our material, hypoplasia was identified in 28.0% of females on the left side and 40.0% on the right side, while in males it was present in 20.0% on the left side and 22.9% on the right side. These values are considerably higher than those reported in the international literature.

This discrepancy may be explained by the fact that our study population consisted of symptomatic patients rather than a healthy general population. *Katsanos et al.* (2017) (17) also demonstrated that vertebral artery hypoplasia is significantly more common in patients with posterior circulation ischemia compared to those with anterior circulation ischemia, supporting the clinical relevance of our findings.

The findings of this study suggest that anatomical variations of the vertebral arteries (VA) are relatively common in the examined group of patients. Among these variations, hypoplasia was the most frequent, especially in females, with the highest rate observed in the right vertebral artery (40.0%).

This result aligns with the work of *Park et al.* (2007), (16) who highlighted the relationship between vertebral artery hypoplasia and changes in posterior cerebral circulation.

Bruneau et al. (2006) (18) also pointed out that understanding these vascular variations is important in clinical practice, particularly during surgical procedures in the cervical region, where unexpected anatomical differences may increase procedural risk.

The higher frequency (f) of hypoplasia in our study compared to the meta-analysis by *Valenzuela-Fuenzalida et al.* (2024) (15) may be explained by the fact that our study included symptomatic patients, who are more likely to present vascular abnormalities.

Dolichoectasia was more commonly observed in

males. Although the number of cases was limited, this finding is still clinically important, as vessel dilation can influence blood flow within the vertebrobasilar system.

One limitation of this study is the lack of a healthy control group. However, comparison with previously published international studies provides useful context and helps support the interpretation of the results.

CONCLUSION:

Anatomical variations of the vertebral arteries were relatively common in the examined population. Hypoplasia was the most frequent variation, particularly among females, whereas dolichoectasia was observed more often in males.

The prevalence of hypoplasia in our study was higher than that reported in the international literature, which is likely related to the symptomatic nature of the study sample. These morphological changes may have important clinical implications for the hemodynamics of posterior cerebral circulation and should be taken into account during radiological and neurological evaluation of symptomatic patients.

Editorial Statements

Ethics Approval and Consent to Participate

This study involved the analysis of data from human participants. The research protocol was reviewed and approved by the Institutional Ethics Committee and was conducted in accordance with institutional regulations, the principles of the Declaration of Helsinki, and internationally accepted standards for biomedical research ethics.

Data Availability Statement

The datasets generated and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Author Contributions (CRediT): Conceptualization & Methodology: GZ, SB; Data Curation, Investigation and Writing – Original Draft: TK; Formal Analysis: SB, ZB; Writing – Review & Editing: KC, SB; Supervision: GZ. All authors have read and approved the final version of the manuscript.

Artificial Intelligence (AI) Statement

The authors used a generative artificial intelligence tool solely for language editing and grammar improvement. All scientific content, data analysis, interpretation, and conclusions were independently developed, verified, and approved by the authors.

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of Interest: The authors declare that they have no competing interests or conflicts of interest related to this study.

Acknowledgements: The authors thank all the healthcare professionals at our institution whose work contributed to the development of evidence-based data, regarding our study

REFERENCES

1. Gray's Anatomy, 39th Edition: The Anatomical Basis of Clinical Practice. (2005). *AJNR: American Journal of Neuroradiology*, 26(10), 2703–2704.
2. Campero, A., Rubino, P.A., Rhoton, A.L. (2011). Anatomy of the vertebral artery. In: Pathology and surgery around the vertebral artery. Springer, Paris. https://doi.org/10.1007/978-2-287-89787-0_4
3. Caplan L. (2000). Posterior circulation ischemia: then, now, and tomorrow. The Thomas Willis Lecture-2000. *Stroke*, 31(8), 2011–2023. <https://doi.org/10.1161/01.str.31.8.2011>
4. Savitz, S. I., & Caplan, L. R. (2005). Vertebrobasilar disease. *The New England journal of medicine*, 352(25), 2618–2626. <https://doi.org/10.1056/NEJMr041544>
5. Lippincott Williams & Wilkins, Philadelphia, PA: 1999. (1999). Osborn AG, Diagnostic Cerebral Angiography. *AJNR: American Journal of Neuroradiology*, 20(9), 1767–1769.
6. Satti, S. R., Cerniglia, C. A., & Koenigsberg, R. A. (2007). Cervical vertebral artery variations: an anatomic study. *AJNR. American journal of neuroradiology*, 28(5), 976–980.
7. Hong JT, et al. Anatomical variations of vertebral artery. *J Korean Neurosurg Soc*. 2009.
8. Park JH, Kim JM, Roh JK, Hypoplastic vertebral artery: frequency and associations with ischaemic stroke territory, *J Neurol Neurosurg Psychiatry*, 2007, 78(9):954–958.
9. El Otmani, H., Fotso, V., El Moutawakil, B., & Rafai, M. A. (2020). Basilar artery fenestration and ischemic stroke: An unclear causal relationship. *Journal de medecine vasculaire*, 45(2), 62–66. <https://doi.org/10.1016/j.jdmv.2020.01.147>
10. Hirano, Y., Miyawaki, S., Yamada, K., Shinya, Y., Kin, T., Amemiya, S., Asakawa, Y., Nakatomi, H., & Saito, N. (2020). Vascular anomaly of the posterior circulation associated with intracranial lipoma-like lesion in the cerebral peduncle manifesting as oculomotor nerve palsy. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*, 72, 463–466. <https://doi.org/10.1016/j.jocn.2019.12.020>
11. Katsanos, A. H., Kosmidou, M., Kyritsis, A. P., & Giannopoulos, S. (2013). Is vertebral artery hypoplasia a predisposing factor for posterior circulation cerebral ischemic events? A comprehensive review. *European neurology*, 70(1-2), 78–83. <https://doi.org/10.1159/000351786>
12. Piccinin MA, Munakomi S. Neuroanatomy, Vertebrobasilar System. [Updated 2023 Jul 24]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2026 Jan. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK540995/>
13. Caplan L. R. (1981). Vertebrobasilar disease. Time for a new strategy. *Stroke*, 12(1), 111–114. <https://doi.org/10.1161/01.str.12.1.111>
14. Doss, A., & Phatouros, C. C. (2006). Vertebrobasilar insufficiency. *Current treatment options in cardiovascular medicine*, 8(2), 111–119. <https://doi.org/10.1007/s11936-006-0003-0>
15. Rhoton AL. The cerebral arteries. Neurosurgery. 2002. van der Grond J, et al. Circle of Willis collateral flow. *Stroke*. 2004.
16. Valenzuela-Fuenzalida J, et al. Vertebral artery hypoplasia: prevalence and clinical significance in cerebrovascular disease. *J Neuroimaging*. 2024.
17. Park JH, et al. Vertebral artery hypoplasia and its association with posterior circulation ischemia. *AJNR Am J Neuroradiol*. 2007;28(8):1530–1534.
18. Katsanos AH, et al. Vertebral artery hypoplasia and posterior circulation stroke: a systematic review. *J Neurol Sci*. 2017; 375:139–144.
19. Bruneau, M., Cornelius, J. F., Marneffe, V., Triffaux, M., & George, B. (2006). Anatomical variations of the V2 segment of the vertebral artery. *Neurosurgery*, 59(1 Suppl 1), ONS20–ONS24. <https://doi.org/10.1227/01.NEU.0000219931.64378.B5>