

Status of the geochemical determination of the solar pp-neutrino flux by LOREX

M. K. Pavićević¹, G. Amthauer¹, I. V. Aničin², F. Bosch³, B. Boev⁴,
W. Bröchle³, V. Cvetković⁵, Ž. Djurčić⁶, W. F. Henning⁶,
R. Jelenković⁵, V. Pejović², A. Weiss⁷

April 15, 2012

Abstract

Allchar is a Sb-As-Tl-Au deposit located in FYR Macedonia at the north-western margins of the Kožuf Mountains, close to the border between FYR Macedonia and Greece. The mine contains the world's largest known concentrations of thallium-bearing minerals and, in particular, the mineral lorandite (TlAsS₂). LOREX, the acronym for LORandite EXperiment, is the only geochemical solar neutrino experiment still actively pursued. It addresses the determination of the long-time average of the solar neutrino flux using the lorandite from Allchar via the neutrino-capture reaction $^{205}\text{Tl} + \nu_e \rightarrow ^{205}\text{Pb} + e^-$. The final step of LOREX would be extraction of lorandite and quantitative determination of the ratio of $^{205}\text{Pb}/^{205}\text{Tl}$ atoms, thus providing the product of solar neutrino flux and neutrino-capture cross section, integrated over the age of the lorandite of 4.31 My. In contrast to the production of ^{205}Pb by solar neutrinos, which is independent of depth, cosmogenic ^{205}Pb produced from fast muons of the secondary cosmic-ray cascade is strongly depth-dependent and very sensitive to the long-term erosion history of the field area. To determine the erosion rate, we applied three different methods: geomorphologic analysis, accelerator mass spectrometry of long-lived radioactive cosmogenic nuclides (^{10}Be and ^{26}Al) and finally noble gas mass spectrometry of the stable cosmogenic nuclides ^3He and ^{21}Ne . For the two important Allchar locations with ore bodies containing lorandite, we thus propose erosion rates of 70 m/My at Crven Dol and 130 m/My at Centralni Deo. With these erosion rates the paleo-depths of lorandite bearing ore-bodies at the time of mineralization range from 450 to 1070 [mwe] what, contrary to some earlier discouraging estimates, leads to satisfactory signal/background ratios. Finally, it is discussed how to obtain the still unknown capture probability of solar pp-neutrinos from ^{205}Tl into ^{205}Pb via the measurement of the lifetime for the bound beta decay of the completely ionised ^{205}Tl , as well as how to count by Schottky mass spectrometry the extremely small number of ^{205}Pb atoms i.e. it is estimated that, depending on paleo-depth, 10kg of lorandite contains about $(3.5 - 11.6) \times 10^5$ atoms of ^{205}Pb , what is needed to attain the accuracy of the final result of the order of about 30%. Both crucial measurements are planned to be carried out at the GSI.

- ¹ University of Salzburg, Division of Material Sciences and Physics,
Department of Mineralogy, Hellbrunnerstr. 34 A-5020 Salzburg, Austria
- ² University of Belgrade, Institute of Physics, Pregrevica 118, 11000
Belgrade, Serbia
- ³ Gesellschaft für Schwerionenforschung GSI, Planckstr. 1, D-64291
Darmstadt, Germany
- ⁴ University of Štip, Faculty of Mining and Geology, Goce Delčev 89, 92000
Štip, FYR Macedonia
- ⁵ University of Belgrade-Faculty of Mining and Geology, Djušina 7/II,
11000 Belgrade, Serbia
- ⁶ Argonne National Laboratory, Physics Division, 9700 South Cass Avenue,
Argonne, Illinois 60439, USA
- ⁷ Max-Planck Institute for Astrophysics, Karl-Schwarzschild-Str.1, D-85741
Garching, Germany