



СЕЛСКОСТОПАНСКА АКАДЕМИЯ – БЪЛГАРИЯ
AGRICULTURAL ACADEMY – BULGARIA



РАСТЕНИЕВЪДНИ
АУКИ®

PLANT SCIENCE

СОФИЯ 2010
ГОД. XLVII, № 1

SOFIA 2010
VOL. XLVII, № 1

**СП. „РАСТЕНИЕВЪДНИ НАУКИ“
СЕ РАЗПРОСТРАНЯВА В НАД 50 СТРАНИ, В Т.Ч.**

Австралия	Дания	Литва	Татарстан
Англия	Индия	Мароко	Унгария
Азербайджан	Испания	Молдова	Узбекистан
Аржентина	Италия	Нова Зеландия	Украина
Армения	Ирландия	Норвегия	Филипините
Белгия	Естония	Полша	Финландия
Беларус	Куба	Португалия	Франция
Бразилия	Канада	Румъния	Холандия
Венецуела	Кения	Русия	Чехия
Гърция	Китай	САЩ	Швейцария
Грузия	Казахстан	Словакия	Швеция
Германия	Киргизстан	Сърбия	ЮАР
Гвинея	Латвия	Турция	Япония

**„PLANT SCIENCE“ JOURNAL (RASTENIEVUDNI NAUKI)
IS CIRCULATED IN MORE THAN 50 COUNTRIES OF THE WORLD**

Australia	Denmark	Lithuania	Hungary
England	India	Morocco	Uzbekistan
Azerbaijan	Spain	Moldova	Ukraina
Argentina	Italy	New Zealand	Philippines
Armenia	Ireland	Norwegia	Finland
Belgium	Estonia	Poland	France
Belorus	Cuba	Portugal	Netherlands
Brazil	Canada	Roumania	Czechia
Venezuela	Kenya	Russia	Switzerland
Greece	China	USA	Sweden
Georgia	Kazakhstan	Slovakia	Republic of South Africa
Germany	Latvia	Serbia	
Guinea	Kirghizia	Turkey	
		Tataria	Japan

РЕДАКЦИОННА КОЛЕГИЯ

Ст.н.с. | ст. дsn Лилия Кръстева; – отговорен редактор;
Ст.н.с. | ст. дsn Ivanka Vitanova; Проф. дsn Ivanka Lecheva;
Ст.н.с. д-р Mariyana Vlahova; Ст.н.с. д-р Neli Valkova;
Ст.н.с д-р Argir Zhivondov; Ст.н.с д-р Dimitrina Ilchovska;
Ст.н.с. д-р Ivan Kiryakov; Ст.н.с. д-р Stoika Masheva;
Ст.н.с. д-р Nikolai Tsenov

EDITORIAL BOARD

Prof. Lilia Krasteva, D. Sc. – Editor-in-Charge;
Prof. Ivanka Vitanova, D. Sc.; Prof. Ivanka Lecheva, D. Sc.;
Assoc. Prof. Mariana Vlahova, Ph.D.; Assoc. Prof. Neli Valkova, Ph.D.;
Assoc. Prof. Argir Zhivondov, Ph.D.; Assoc. Prof. Dimitrina Ilchovska, Ph.D.;
Assoc. Prof. Ivan Kiryakov, Ph.D.; Assoc. Prof. Stoika Masheva, Ph.D.;
Assoc. Prof. Nikolai Tsenov, Ph.D.

РАСТЕНИЕВ. НАУКИ/RASTENIEV. NAUKI

WEB: www.rastnauki.org
E-mail: plants@abv.bg

Списание „Растениевъдни науки“ е издание на Селскостопанска академия и излиза в 6 броя годишно. Списанието публикува на български и английски език оригинални изследователски статии, кратки съобщения и обзори, покриващи практически всички области на растениевъдството: растителна генетика, клетъчни и тъканни култури, генетично инженерство, генетични ресурси, селекция, растителна защита и опазване на екосистемите, агротехника, технологии за отглеждане и производство на семена, посадъчен материал и растителна продукция, почвознание и други сродни области, които могат да допринесат за развитието на този отрасъл.

„Plant Science“ (Rasteniev'dni nauki) journal is an edition of the Agricultural Academy and comes out in 6 issues annually. The journal publishes both in Bulgarian and English languages original research papers, brief reports and review articles covering practically all spheres of the plant growing industry: plant genetics, cell and tissue cultures, gene engineering, genetic resources, plant breeding, plant protection, field management, technologies of growing, production of seeds and planting material, soil science and other related spheres that may contribute to the development of this industry.

Списание „Растениевъдни науки“ е включено в следните реферативни издания: Agri Eng. Abstr., Agroforest. Abstr., Apic. Abstr., BSL, Biol., Bio-Contr. News & Info., Biol. Abstr., Chem. Abstr., Cott. & Trop Fibr. Abstr., Crop Physiol. Abstr., Curr. Adv. Ecol. Sci., Field Crop Abstr., Food Sci & Tech. Abstr., Forest. Abstr., Helminthol. Abstr., Herb. Abstr., Hort. Abstr., INIS Atomind., Irr. & Drain. Abstr., Maize Abstr., Nutr. Abstr., Ornarn Hort., Plant Breed. Abstr., Plant Gen. Res. Abstr., Plant Grow. Reg. Abstr., Postharvest, Potato Abstr., Ref. Zh., Rev. Appl. Entomol., Rev. Plant Path., Rice Abstr., Seed Abstr., Soils & Fert., Sorghum & Millets Abstr., Soyabean Abstr., Triticale Abstr., Trop. Oil Seeds Abstr., VITIS, Weed Abstr., Word Ag. Econ. & Rur. Soc. Abstr., Zoo. Rec.; Slovak Agricultural Library; Систематический указатель иностранной литературы на ЦНСХБ – Россия; Электронный каталог журналов, Реферативный журнал.

Списанието се обработва и в следните бази данни: AGRIS; AGRICOLA, BIOSIS CAB ABSTRACTS, CAPLUS SCOPUS.

Списание „Растениевъдни науки“ е включено и в SCIENCE CITATION INDEX (SCI).

Публикуваните в този брой на списанието материали са докладвани на проведената на 1-2. X. 2009 г. научна конференция, организирана от Земеделският институт в Стара Загора

РЕДАКЦИЯ „НАУЧНИ ИЗДАНИЯ“ НА СЕЛСКОСТОПАНСКА АКАДЕМИЯ

1113 София, бул. „Цариградско шосе“, 125, бл. 1
Снежана Дакева – редактор (тел.: 0878 976 248)

PUBLISHING DEPARTMENT OF SCIENTIFIC ISSUES OF AGRICULTURAL ACADEMY

1113 Sofia, 125 Tsarigradsko shosse Bul., Block 1
Snezhana Dakeva – Editor (tel.: +359 878 976 248)

Селскостопанска академия, 2010

C/o Jusautor, Sofia

Коректор – С. Дакева

Предпечатна подготовка – ФОТОНИКА
Печат: БПС ООД

формат	60x84/8	Годишен абонамент – 90,00 лв.	Печатни коли – 12,00
		Отделна книжка – 15,00 лв.	Авторски коли – 24,00

СЕЛСКОСТОПАНСКА АКАДЕМИЯ
AGRICULTURAL ACADEMY
РАСТЕНИЕВЪДНИ НАУКИ, 47, 33—35
PLANT SCIENCE, 47, 33—35
София, 2010, Sofia

DEPENDENCE ON THE FLORAL CONTENT AND THE WEED NUMBER FROM GROWING SYSTEM IN OAT IN STRUMICA REGION

DRAGICA SPASOVA*, DUSAN SPASOV, MITE ILIEVSKI*, BILJANA ATANASOVA
Goce Delcev University — Stip, Faculty of Agriculture, Goce Delcev b.b. 2400 Strumica, R. Macedonia

G. VASILEVSKI
St. Cyril and Methodius University — Skopje, Faculty of agricultural sciences and food, Aleksandar
Makedonski b.b. 1000 Skopje, R. Macedonia

Abstract: The main aim of this examination was to determine the dependence of the floral content and weed number from the growing system and to establish appropriate agro technical methods for reaching stable and high grain yields in conditions of conventional and organic production of oat. The examinations started in 2005 and lasted until 2007. In the experiment with conventional oat production ten (10) weed species were present, and in organic experiment nine (9) weed species were present. Considerate that plants in organic production, in all years of examination, in the initial growing stages, had faster vegetative growing, we could note that weed vegetation had more unfavorable growing conditions, and is less represented than the weed vegetation in the experiment with conventional growing system, without use of herbicides.

Key words: weeds, herbicides, oat, organic production, conventional production

D. SPASOVA, G. VASILEVSKI*, D. SPASOV, M. ILIEVSKI, B. ATANASOVA, University „Goce Delcev“ — Stip, Faculty of Agriculture, Goce Delcev b.b. 2400 Strumica, R. Macedonia, *St. Cyril and Methodius University — Skopje, Faculty of agricultural sciences and food, Aleksandar Makedonski b.b. 1000 Skopje, R. Macedonia. ЗАВИСИМОСТ НА БОТАНИЧНИЯ СЪСТАВ И БРОЯ ПЛЕВЕЛИ ОТ СИСТЕМАТА НА ОТГЛЕЖДАНЕ НА ОВЕС В РАЙОНА НА СТРУМИЦА

Резюме: Главната цел на това изследование беше да се определят зависимостта на ботаничния състав и броя плевели от системата на отглеждане и да се установят подходящи агротехнически методи за достигане на стабилен и висок зърнодобив в условията на обикновено и органично производство на овес. Изпитванията започнаха през 2005 г. и завършиха през 2007 г. В експеримента с обикновеното производство на овес бяха изследвани десет плевелни вида, а в органичния експеримент — девет плевелни вида. Растенията при органичното производство са имали през всички години на производство по-бърз вегетационен растеж в първоначалните фази на вегетация, като може да се забележи, че развитието на плевелите е било по-неблагоприятно и те са по-малко представени на брой в сравнение с експеримента, проведен при обикновени условия без използването на хербициди.

Ключови думи: плевели, хербициди, овес, органично производство, традиционно производство

INTRODUCTION

In Republic of Macedonia, the average grain yield from oat, in the period after the Second World War, is in continually decadence.

The main reasons for this problem are not productive varieties and inappropriate agrotechnical methods.

From the other side, today the oat is counted in the very important cultures in the human diet, with

increased call in the modern day culinary and the foodstuff technology. Gradually, the oat takes notably place in the human diet because it is the most nutritious cereal with big quantity of proteins in the grain. Besides the proteins the oat grain contain also soluble diet fibers as β -glucans, whose content varies from 2,5—6,5% (Pržulj et al. 1998). The content of β -glucans decreases the cholesterol in the blood, so today

*E-mail: dragica.spasova@ugd.edu.mk

diet food recommends food on the base of oat flakes (Welch, 1991, cit: Mlinar 1996).

To increase and stabilize the yields, it is necessary to make systematical approach in the choice of high yield varieties, that will reach theirs genetic potential in some soil and climatic conditions in the region, using appropriate agrotechnical measures.

Today, in Macedonia, the oat is grown on 2 162 ha (Statistical review: Agriculture 5.4.7.01/564, July 2007) and the average yield is 1746 kg/ha.

Because of the favorable climatic conditions that predominate in Strumica micro region, the favorable politics of our country in the last year for this culture, as well as the tradition that exist in this location, there are real chances to increase the surfaces with that culture in the next period, so the needs of the country would be satisfied, and the deficiency of that culture will be decreased.

One of the main factors in growing oat is weeds.

Besides the other analyses, in the same time we made an observation of the weeds in both growing systems, separately, by variants.

The main aim was to determine the dependence of the floral content and weed number from the growing system and to establish appropriate agro technical methods for reaching stable and high grain yields in conditions of conventional and organic production of oat.

MATERIAL AND METHODS

The examinations started in 2005 and lasted until 2007. The field experiments were set on the field of UGD, Faculty of agriculture, Strumica.

Five oat population, of which four are home population and one is introduced from Bulgaria (*population Krivogastani, population Trebenista, population Radolista, population from Bulgaria and population Kuceviste*), and three oat varieties (*Rejac, Slavuj and Lovken*) introduced from Serbia were analyzed.

In the three year examination two experiments were set (Fig. 1), one in condition of organic and one in condition of conventional production, in which all oat genotypes were maintained. The experiments



Fig. 1. Left, conventional — treated,
right, organic - untreated

consisted of 8 variants in four repetitions, arranged by randomized block system, with dimension of the basic parcel of 5 m². The distance between variants is 50 cm, and between repetitions 100 cm.

The distance between lines at conventional production was 20 cm, and at organic production was 10 cm. In all years of examination culture before oat was potato.

In all years of examination the soil was prepared in the same way. During the autumn the surface was plowed in deepness of 35 cm, than the surface was separated and fertilized by methodological principle, so the surface predicted for conventional production was fertilized with 300 kg/ha NPK fertilizer in combination of 15:15:15, while the surface predicted for organic production, 20 t/ha organic fertilizer was inserted. After fertilizing the surface was additionally cultivated and was leveled.

The seed in all years of examination was done in March, 17.03.2005, 28.03.2006 and 06.03. 2007, when the conditions for seeding were optimal. The seeding was made by hand in rows, on 5—6 cm in deepness.

During the vegetation of oat, we made control of the weeds in organic and conventional production, for determining the number and species of the weeds. The number of the weeds was determined by the method of squares, i.e. by counting the weeds on 1m² of each repetition of each variant, and than the average of all repetitions is taken. After estimation of the weeds, the oat in stage tillering grown in conditions of conventional production was treated with herbicide 2,4 — D, against wide leaves weeds in the

RESULTS AND DISCUSSION

The floral content of the weed vegetation is brought out in table 1. From the made measurements and analyses it could be seen that in the organic production, in all years of examination 9 weed species are represented: *Papaver rhoes*, *Veronica chederifolia*, *Stellaria media*, *Raphanus raphanistrum*, *Polygonum aviculare*, *Vicia Vilosa*, *Cirsium arvense*, *Equisetum arvense* and *Apera spica venti*. In the conventional production, in all years of examination 10 weed species are represented: *Papaver rhoes*, *Veronica chederifolia*, *Stellaria media*, *Raphanus raphanistrum*, *Sinapis arvensis*, *Polygonum aviculare*, *Vicia Vilosa*, *Cirsium arvense*, *Equisetum arvense* and *Apera spica venti*.

Apart the varieties and populations, the average weediness of the experiment grown in conditions of organic production is 15, 15 weeds/m² in 2007, to 19,20 weeds/m² in 2005. The average weediness of the experiment grown in conditions of organic production of oat is 17, 20 weeds/m².

Apart the varieties and populations, the average weediness of the experiment grown in conditions of conventional production is 36,45 weeds/m² in 2007, to 42,95 weeds/m² in 2006. The average weediness of

Table 1/Таблица 1

Floral content of the weed vegetation in the experiment (No/m^2)

Species of weeds	Organic production			Average 2005/07	Conventional production			Average 2005/07
	2005	2006	2007		2005	2006	2007	
<i>Papaver rhoeas</i>	3,50	3,25	3,25	3,33	6,25	6,50	5,25	6,0
<i>Veronica chederifolia</i>	3,55	1,80	1,20	2,18	4,75	4,80	4,50	4,68
<i>Stellaria media</i>	2,75	2,80	1,80	2,45	4,50	5,25	4,50	4,75
<i>Raphanus raphanistrum</i>	0,25	0,25	0,25	0,25	2,50	2,50	1,75	2,25
<i>Sinapis arvensis</i>	0,25	/	0,25	0,16	2,12	2,15	2,25	2,18
<i>Polygonum aviculare</i>	3,15	3,25	3,25	3,22	6,12	6,25	6,20	6,20
<i>Vicia villosa</i>	1,50	1,25	1,15	1,30	3,50	2,75	2,50	2,90
<i>Cirsium arvense</i>	1,00	0,75	0,75	0,83	3,50	3,75	3,25	3,50
<i>Equisetum arvense</i>	/	0,50	/	0,16	/	1,25	0,25	0,50
<i>Apera spica venti</i>	3,25	3,50	3,25	3,33	7,25	7,50	6,25	7,0
Total in 1 m ²	19,2	17,35	15,15	17,20	40,50	42,75	36,60	40,0
Total species	9	9	9		9	10	10	

the experiment grown in conditions of organic production of oat is 40,0 weeds/m².

Comparing the average weediness in both growing systems, apart the years, climatic conditions and variants, and depend of the used agro technical measures, it could be noted that the average weediness of the experiment grown in conditions of organic production (17,20 weeds/m²) is smaller absolutely for 22,8 weeds/m² or relatively for 132,6% from the average weediness of the experiment grown in conditions of conventional production (40,0 weeds/m²).

Considerate that plants in organic production, in all years of examination, in the initial growing stages, had faster vegetative growing, we could note that weed vegetation had more unfavorable growing conditions, and is less represented than the weed vegetation in the experiment with conventional growing system.

CONCLUSIONS

From the results the following conclusions could be made:

In conditions of conventional production of oat ten (10) weed species were represented, and in conditions of organic production nine (9) weed species were represented.

Apart the variants, the average weediness of the experiment grown in conditions of conventional production, in both years of examination is 40,0 weeds/m².

Apart the variants, the average weediness of the experiment grown in conditions of organic production, in both years of examination is 17,2 weeds/m².

The differences that appear in average weediness in both growing systems, are a result of use of different

agrotechnical measures, although the soil and climatic characteristics and the variants are same.

The agricultural measures used in conditions of organic production of cereal mixtures were better alternative solution for decreasing weeds, than those used in conventional production.

REFERENCES

- Benčeva N., T. Georgieva (1998). Ikonomičeski efekt ot vzaimodejstvieto na sroka i gastotata na sejbtata pri zimuvashtia oves. Ikonomika i upravlenie na selskoto stopanstvo, 3, 31—33.
- Bogdanović, M., N. Pržulj (2000): Rezultati ispitivanja jare zobi u Brdskom-planinskom području Republike Srbije, Agroznanje, Poljoprivredni naučno stručni i informativni časopis, Godina 1, broj 3, 84—90, Banja Luka.
- Vasiljević G. (2004): Zrnesti i klubenesti kulturi (Univerzitetski učebnik), Univerzitet „Sv. Kiril i Metodij“, Skopje, Fakultet za zemljodelski nauki i hrana, Skopje.
- Georgieva, T., K. Kostov (1998). Proučvane varhov biologičnите proyavi v rasteža i razvitioto na novi kandidat-sortove zimuvasht oves v usloviata na Centralna Južna Bulgaria (I). Institut po Ečemika — Karnobat, Naučni trudove Tom VII, 313—317, Selskostopanska akademija — Sofia.
- Zinta, G., M. Beidere, I. Skrabule, J. Vigovskis (2004). The first steps in variety testing for organic agriculture in Latvia: oats and potatoes. Proceedings of the First World Conference on Organic Seed. Challenges and Opportunities for Organic Agriculture and the Seed Industry, FAO Headquarters, Rome, Italy.
- Milnar, R., Z. Martinčić-Jerčić (1996): Program implementiranja jare zobi u BiH Institutu, d.d., Zagreb. Agronomski glasnik, 49—61.
- Pržulj, N., V. Momčilović, V. Đurić (1998): Proizvodnja i preprada jecma i ovsu za ljudsku ishranu XIII Savetovanje žito-hleb, 15—16, Novi Sad, Jugoslavija.
- [1] Državeni zavod za statistika na Republika Makedonija (2008): Poljodelstvo, ovojtarstvo i lozarstvo, 2007. Statistički pregledi: Zemljodelstvo. Maj 2008. Skopje.